

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bis(2,4,6-triamino-1,3,5-triazin-1-ium) hydrogen phosphate trihydrate

Xue-Mei Li,^a Si-Si Feng,^{b*} Fan Wang,^a Qi Ma^{a,b} and Miao-Li Zhu^b

^aCollege of Chemistry and Chemical Engineering, Shanxi Datong University, Datong, Shanxi 037009, People's Republic of China, and ^bInstitute of Molecular Science, Key Laboratory of Chemical Biology and Molecular, Engineering of the Education Ministry, Shanxi University, Taiyuan, Shanxi 030006, People's Republic of China
Correspondence e-mail: sffeng@sxu.edu.cn miaoli@sxu.edu.cn


Received 22 November 2009; accepted 20 December 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{N}-\text{C}) = 0.005$ Å; disorder in main residue; R factor = 0.057; wR factor = 0.178; data-to-parameter ratio = 10.2.

In the title hydrated molecular salt, $2\text{C}_3\text{H}_7\text{N}_6^+ \cdot \text{HPO}_4^{2-} \cdot 3\text{H}_2\text{O}$, three of the O atoms of the hydrogen phosphate anion are disordered over two positions, with relative occupancies of 0.763 (1) and 0.237 (1). In the crystal, the components are linked by $\text{N}-\text{H} \cdots \text{N}$, $\text{N}-\text{H} \cdots \text{O}$ and $\text{O}-\text{H} \cdots \text{O}$ hydrogen bonds

Related literature

For related structures, see: Choi *et al.* (2004); Janczak & Perpétuo (2001*a,b,c*, 2002, 2003, 2004); Li *et al.* (2005); Perpétuo & Janczak (2002); Zhang *et al.* (2004).



Experimental

Crystal data

 $2\text{C}_3\text{H}_7\text{N}_6^+ \cdot \text{HPO}_4^{2-} \cdot 3\text{H}_2\text{O}$ $M_r = 404.32$ Triclinic, $P\bar{1}$ $a = 6.767$ (4) Å $b = 10.548$ (7) Å $c = 12.497$ (8) Å $\alpha = 91.865$ (11)° $\beta = 105.609$ (10)° $\gamma = 108.020$ (9)° $V = 810.3$ (9) Å³ $Z = 2$ Mo $K\alpha$ radiation $\mu = 0.24$ mm⁻¹ $T = 296$ K

0.30 × 0.25 × 0.20 mm

Data collection

Bruker SMART 1K CCD

diffractometer

Absorption correction: multi-scan

(SADABS; Bruker, 2000)

 $T_{\min} = 0.933$, $T_{\max} = 0.954$

3823 measured reflections

2803 independent reflections

2027 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.029$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.057$ $wR(F^2) = 0.178$ $S = 1.06$

2803 reflections

274 parameters

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.33$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.53$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{N3}-\text{H3} \cdots \text{O5}^{\text{i}}$	0.77 (4)	2.06 (4)	2.791 (4)	158 (4)
$\text{N4}-\text{H4B} \cdots \text{O4A}^{\text{ii}}$	0.86	2.22	2.998 (5)	151
$\text{N4}-\text{H4A} \cdots \text{N7}^{\text{iii}}$	0.86	2.20	3.059 (5)	177
$\text{N5}-\text{H5B} \cdots \text{O7}$	0.86	1.97	2.822 (5)	170
$\text{N5}-\text{H5A} \cdots \text{O6}^{\text{iv}}$	0.86	2.41	3.258 (5)	167
$\text{N6}-\text{H6B} \cdots \text{O5}^{\text{i}}$	0.86	2.28	3.009 (5)	143
$\text{N6}-\text{H6A} \cdots \text{N9}^{\text{v}}$	0.86	2.17	3.031 (5)	176
$\text{N8}-\text{H8} \cdots \text{O3A}$	0.84 (3)	1.93 (4)	2.742 (7)	164 (3)
$\text{N10}-\text{H10B} \cdots \text{O2A}$	0.86	2.11	2.952 (5)	168
$\text{N10}-\text{H10A} \cdots \text{O2A}^{\text{vi}}$	0.86	1.99	2.744 (5)	146
$\text{N11}-\text{H11B} \cdots \text{O4A}^{\text{vii}}$	0.86	2.53	3.147 (6)	129
$\text{N11}-\text{H11B} \cdots \text{O1}^{\text{viii}}$	0.86	2.33	3.121 (5)	152
$\text{N11}-\text{H11A} \cdots \text{N1}^{\text{v}}$	0.86	2.02	2.884 (4)	178
$\text{N12}-\text{H12B} \cdots \text{O6}^{\text{ix}}$	0.86	2.07	2.840 (5)	149
$\text{N12}-\text{H12A} \cdots \text{N2}^{\text{iii}}$	0.86	2.08	2.935 (4)	172
$\text{O1}-\text{H1} \cdots \text{O3A}^{\text{vii}}$	0.82	1.76	2.535 (5)	157
$\text{O5}-\text{H5D} \cdots \text{O3A}^{\text{i}}$	0.85	1.89	2.739 (5)	177
$\text{O5}-\text{H5C} \cdots \text{O4A}$	0.85	2.00	2.726 (6)	142
$\text{O6}-\text{H6D} \cdots \text{O1}^{\text{ix}}$	0.85	1.97	2.787 (4)	161
$\text{O6}-\text{H6C} \cdots \text{O4A}$	0.85	2.02	2.838 (6)	162
$\text{O7}-\text{H7B} \cdots \text{O1}^{\text{ix}}$	0.85	2.57	3.257 (5)	139
$\text{O7}-\text{H7B} \cdots \text{O2A}^{\text{ix}}$	0.85	2.28	2.958 (6)	137
$\text{O7}-\text{H7A} \cdots \text{O5}^{\text{i}}$	0.85	2.28	3.006 (6)	144

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x, y, z-1$; (iii) $-x+1, -y+2, -z$; (iv) $-x+2, -y+2, -z+1$; (v) $-x, -y+1, -z$; (vi) $-x+1, -y+2, -z+1$; (vii) $-x, -y+1, -z+1$; (viii) $x-1, y, z-1$; (ix) $x+1, y, z$.

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997); software used to prepare material for publication: SHELXTL (Sheldrick, 2008).

This work was supported by the Science Foundation of Shanxi Datong University of China (No.2005k03) and the Doctoral Start-up Foundation of Shanxi University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5248).

References

- Bruker (2000). SMART, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Choi, C. S., Venkatraman, R., Kim, E. H., Hwang, H. S. & Kang, S. K. (2004). *Acta Cryst.* **C60**, o295–o296.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Janczak, J. & Perpétuo, G. J. (2001*a*). *Acta Cryst.* **C57**, 123–125.
Janczak, J. & Perpétuo, G. J. (2001*b*). *Acta Cryst.* **C57**, 1120–1122.

- Janczak, J. & Perpétuo, G. J. (2001c). *Acta Cryst.* **C57**, 1431–1433.
- Janczak, J. & Perpétuo, G. J. (2002). *Acta Cryst.* **C58**, o455–o459.
- Janczak, J. & Perpétuo, G. J. (2003). *Acta Cryst.* **C59**, o349–o352.
- Janczak, J. & Perpétuo, G. J. (2004). *Acta Cryst.* **C60**, o211–o214.
- Li, X.-M., Lu, L.-P., Feng, S.-S., Zhang, H.-M., Qin, S.-D. & Zhu, M.-L. (2005). *Acta Cryst.* **E61**, o811–o813.
- Perpétuo, G. J. & Janczak, J. (2002). *Acta Cryst.* **C58**, o112–o114.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Zhang, J., Kang, Y., Wen, Y.-H., Li, Z.-J., Qin, Y.-Y. & Yao, Y.-G. (2004). *Acta Cryst.* **E60**, o462–o463.

supplementary materials

Acta Cryst. (2010). E66, o239-o240 [doi:10.1107/S1600536809054798]

Bis(2,4,6-triamino-1,3,5-triazin-1-ium) hydrogen phosphate trihydrate

X.-M. Li, S.-S. Feng, F. Wang, Q. Ma and M.-L. Zhu

Comment

In previous work, we reported a compound of 2,4,6-triamino-1,3,5-triazin-1-ium with organic anion linked *via* weak interactions (Li *et al.*, 2005). As continuing interest, here, we present the results of the crystal structure of another ion pair adduct containing monoprotonated melaminium, inorganic anion (hydrogen phosphate), and solvent water, (I, Fig. 1 & Table 1).

The aromatic rings both monoprotonated melaminium exhibit significant distortions from the ideal hexagonal form. Thus, the internal C—N—C angles at the protonated N atoms (C1—N3—C2 and C6—N8—C5) are significantly greater than the other two ring (*i.e.* C1—N2—C2, C1—N1—C3, C5—N7—C4, C6—N9—C4) angles, and the internal N2—C1—N1 and N7—C4—N9 angle, *i.e.* containing only non-protonated N atoms is greater than either of the remaining N—C—N angles containing both protonated and non-protonated N atoms (Table 1). This feature of the structure is similar to our previous report (Li *et al.*, 2005) and the other reported mono-protonated melaminium cations (Janczak & Perpétuo, 2001a,b,c, 2002, 2003, 2004; Perpétuo & Janczak, 2002; Zhang *et al.*, 2004; Choi *et al.*, 2004).

Between charged residues and the water molecules interact extensively by a combination of ionic, H-bonds (Table 2) as well as π - π interactions as shown in Fig 2. Neighboring melaminium residues are interconnected by double N—H \cdots N H-bonds, leading to the formation of layers.

Experimental

Hot solutions of melamine and *meta*-phosphoric acid in a 1:1 molar ratio were mixed and, after allowed the mixture to stand at room temperature for a few days, colourless blocks of (I) were desposited.

Refinement

H atoms attached to N (except N3 and N8) and O1 atoms of (I) were placed in geometrically idealized positions with N—H = 0.86 Å O1—H = 0.82 Å and refined with $U_{\text{iso}}=1.2U_{\text{eq}}$ [N] or $U_{\text{iso}}=1.5U_{\text{eq}}$ [O1]. H atoms attached to O (water) N3 and N8 atom were located from the difference Fourier maps and refined their global $U_{\text{iso}}=1.2U_{\text{eq}}$. The N3—H, N8—H, and O—H distances are in the range 0.768 - 0.851 Å. The O2, O3, and, O4 atoms from hydrogen phosphate anion of crystallization in (I) were found to be disordered and were modelled over two sets of positions using restraints on the anisotropic displacement parameters of these O atoms. The major and minor disorder components had refined occupancies of O2A 0.77 (1), O3A 0.76 (1), O4A 0.76 (1) % and O2B 0.23 (1) O3B 0.24 (1) O4B 0.24 (1)%, respectively.

Figures




Fig. 1. The structure of (I), with displacement ellipsoids drawn at the 30% probability level for non-H atoms.





Fig. 2. The packing of (I) viewed down the *a* axis.

Bis(2,4,6-triamino-1,3,5-triazin-1-ium) hydrogen phosphate trihydrate

Crystal data



$M_r = 404.32$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 6.767\ (4)\ \text{\AA}$

$b = 10.548\ (7)\ \text{\AA}$

$c = 12.497\ (8)\ \text{\AA}$

$\alpha = 91.865\ (11)^\circ$

$\beta = 105.609\ (10)^\circ$

$\gamma = 108.020\ (9)^\circ$

$V = 810.3\ (9)\ \text{\AA}^3$

$Z = 2$

$F(000) = 424$

$D_x = 1.657\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 1970 reflections

$\theta = 2.5\text{--}28.0^\circ$

$\mu = 0.24\ \text{mm}^{-1}$

$T = 296\ \text{K}$

Block, colourless

$0.30 \times 0.25 \times 0.20\ \text{mm}$

Data collection

Bruker SMART 1K CCD
diffractometer

Radiation source: fine-focus sealed tube
graphite

ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2000)

$T_{\min} = 0.933$, $T_{\max} = 0.954$

3823 measured reflections

2803 independent reflections

2027 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

$\theta_{\text{max}} = 25.1^\circ$, $\theta_{\text{min}} = 2.1^\circ$

$h = -7 \rightarrow 8$

$k = -11 \rightarrow 12$

$l = -14 \rightarrow 7$

Refinement

Refinement on F^2

Primary atom site location: structure-invariant direct
methods

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.057$$

$$wR(F^2) = 0.178$$

$$S = 1.06$$

2803 reflections

274 parameters

0 restraints

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0996P)^2 + 0.520P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.003$$

$$\Delta\rho_{\max} = 0.33 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.53 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.5323 (6)	0.7920 (4)	-0.0156 (3)	0.0352 (8)	
C2	0.6740 (6)	0.8584 (4)	0.1714 (3)	0.0359 (8)	
C3	0.4373 (6)	0.6393 (4)	0.1008 (3)	0.0337 (8)	
C4	0.0847 (5)	0.8128 (3)	0.0751 (3)	0.0317 (8)	
C5	0.2296 (6)	0.8597 (4)	0.2627 (3)	0.0372 (8)	
C6	-0.0212 (5)	0.6489 (4)	0.1803 (3)	0.0329 (8)	
N1	0.4191 (5)	0.6664 (3)	-0.0031 (2)	0.0352 (7)	
N2	0.6575 (5)	0.8903 (3)	0.0688 (2)	0.0355 (7)	
N3	0.5651 (5)	0.7323 (3)	0.1889 (3)	0.0363 (7)	
H3	0.577 (6)	0.712 (4)	0.248 (3)	0.029 (10)*	
N4	0.5233 (5)	0.8212 (3)	-0.1176 (3)	0.0436 (8)	
H4A	0.5955	0.9002	-0.1280	0.052*	
H4B	0.4453	0.7613	-0.1741	0.052*	
N5	0.7957 (5)	0.9458 (3)	0.2586 (3)	0.0459 (8)	
H5A	0.8672	1.0258	0.2499	0.055*	
H5B	0.8047	0.9235	0.3249	0.055*	
N6	0.3325 (5)	0.5192 (3)	0.1207 (3)	0.0418 (8)	
H6A	0.2515	0.4577	0.0659	0.050*	
H6B	0.3450	0.5021	0.1886	0.050*	
N7	0.2203 (5)	0.9023 (3)	0.1634 (2)	0.0358 (7)	
N8	0.1115 (5)	0.7333 (3)	0.2726 (3)	0.0374 (7)	
H8	0.126 (5)	0.703 (3)	0.334 (3)	0.021 (8)*	

supplementary materials

N9	-0.0400 (4)	0.6864 (3)	0.0791 (2)	0.0330 (7)	
N10	0.3541 (5)	0.9350 (4)	0.3562 (3)	0.0552 (10)	
H10A	0.4343	1.0157	0.3550	0.066*	
H10B	0.3558	0.9039	0.4191	0.066*	
N11	-0.1322 (5)	0.5270 (3)	0.1931 (3)	0.0420 (8)	
H11A	-0.2178	0.4709	0.1354	0.050*	
H11B	-0.1193	0.5031	0.2591	0.050*	
N12	0.0741 (5)	0.8512 (3)	-0.0246 (2)	0.0379 (7)	
H12A	0.1520	0.9305	-0.0310	0.045*	
H12B	-0.0107	0.7969	-0.0834	0.045*	
P1	0.24796 (15)	0.68587 (9)	0.57240 (7)	0.0335 (3)	
O1	0.0294 (5)	0.6330 (3)	0.6053 (3)	0.0599 (9)	
H1	-0.0482	0.5614	0.5669	0.090*	
O2A	0.2963 (6)	0.8336 (3)	0.5671 (3)	0.0445 (14)	0.767 (9)
O3A	0.2088 (7)	0.6098 (4)	0.4601 (4)	0.0353 (15)	0.757 (14)
O4A	0.4058 (6)	0.6486 (5)	0.6643 (3)	0.0507 (16)	0.765 (11)
O2B	0.176 (2)	0.7686 (13)	0.4856 (10)	0.046 (5)	0.233 (9)
O3B	0.279 (2)	0.5614 (14)	0.5130 (13)	0.040 (5)	0.243 (14)
O4B	0.4481 (19)	0.7567 (17)	0.6642 (10)	0.048 (5)	0.235 (11)
O5	0.4731 (5)	0.4107 (3)	0.6295 (3)	0.0606 (9)	
H5C	0.4405	0.4792	0.6078	0.091*	
H5D	0.5757	0.4063	0.6041	0.091*	
O6	0.8660 (5)	0.7583 (4)	0.7442 (3)	0.0753 (11)	
H6C	0.7299	0.7431	0.7189	0.113*	
H6D	0.8984	0.7038	0.7061	0.113*	
O7	0.8612 (6)	0.8583 (4)	0.4724 (3)	0.0799 (11)	
H7A	0.7320	0.8059	0.4420	0.120*	
H7B	0.9418	0.8093	0.4809	0.120*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0292 (18)	0.040 (2)	0.040 (2)	0.0123 (16)	0.0150 (15)	0.0077 (16)
C2	0.0342 (19)	0.039 (2)	0.042 (2)	0.0175 (16)	0.0177 (16)	0.0081 (17)
C3	0.0292 (18)	0.038 (2)	0.040 (2)	0.0133 (15)	0.0165 (15)	0.0067 (16)
C4	0.0274 (18)	0.039 (2)	0.0343 (19)	0.0133 (15)	0.0146 (15)	0.0090 (15)
C5	0.0281 (18)	0.047 (2)	0.035 (2)	0.0074 (16)	0.0143 (15)	0.0034 (16)
C6	0.0272 (18)	0.044 (2)	0.0321 (19)	0.0140 (16)	0.0127 (14)	0.0088 (15)
N1	0.0342 (17)	0.0360 (17)	0.0369 (17)	0.0086 (13)	0.0161 (13)	0.0076 (13)
N2	0.0348 (16)	0.0349 (17)	0.0402 (17)	0.0088 (13)	0.0193 (13)	0.0082 (13)
N3	0.0397 (18)	0.0396 (18)	0.0358 (19)	0.0147 (14)	0.0188 (15)	0.0119 (15)
N4	0.0468 (19)	0.0413 (18)	0.0375 (18)	0.0038 (15)	0.0161 (14)	0.0071 (14)
N5	0.054 (2)	0.0432 (19)	0.0371 (18)	0.0088 (15)	0.0154 (15)	0.0043 (15)
N6	0.0464 (19)	0.0408 (19)	0.0396 (18)	0.0101 (15)	0.0188 (15)	0.0120 (14)
N7	0.0302 (16)	0.0381 (17)	0.0375 (17)	0.0053 (13)	0.0146 (13)	0.0040 (13)
N8	0.0384 (18)	0.047 (2)	0.0266 (17)	0.0108 (15)	0.0130 (14)	0.0108 (14)
N9	0.0303 (15)	0.0344 (17)	0.0331 (16)	0.0065 (13)	0.0119 (12)	0.0084 (12)
N10	0.048 (2)	0.062 (2)	0.0367 (19)	-0.0071 (17)	0.0125 (16)	-0.0012 (16)

N11	0.0438 (18)	0.0413 (18)	0.0323 (17)	0.0017 (14)	0.0111 (14)	0.0125 (13)
N12	0.0408 (18)	0.0368 (17)	0.0337 (17)	0.0056 (14)	0.0150 (13)	0.0101 (13)
P1	0.0332 (5)	0.0360 (5)	0.0311 (5)	0.0064 (4)	0.0142 (4)	0.0090 (4)
O1	0.0576 (19)	0.0512 (19)	0.068 (2)	-0.0043 (14)	0.0431 (16)	-0.0138 (15)
O2A	0.047 (2)	0.039 (2)	0.047 (3)	0.0066 (17)	0.022 (2)	0.0108 (18)
O3A	0.036 (2)	0.037 (2)	0.033 (2)	0.0065 (16)	0.0155 (18)	0.0097 (17)
O4A	0.058 (3)	0.059 (4)	0.038 (2)	0.028 (2)	0.0091 (17)	0.0085 (18)
O2B	0.060 (9)	0.058 (9)	0.030 (8)	0.032 (7)	0.015 (7)	0.015 (6)
O3B	0.042 (7)	0.045 (8)	0.039 (9)	0.019 (6)	0.015 (6)	-0.006 (6)
O4B	0.039 (7)	0.054 (12)	0.047 (7)	0.012 (6)	0.008 (5)	0.011 (6)
O5	0.073 (2)	0.075 (2)	0.065 (2)	0.0440 (18)	0.0458 (17)	0.0388 (16)
O6	0.065 (2)	0.121 (3)	0.0487 (19)	0.050 (2)	0.0113 (16)	-0.0150 (18)
O7	0.085 (3)	0.086 (3)	0.077 (2)	0.036 (2)	0.025 (2)	0.037 (2)

Geometric parameters (Å, °)

C1—N4	1.311 (5)	N6—H6A	0.8600
C1—N2	1.346 (5)	N6—H6B	0.8600
C1—N1	1.347 (5)	N8—H8	0.84 (3)
C2—N5	1.303 (5)	N10—H10A	0.8600
C2—N2	1.320 (5)	N10—H10B	0.8600
C2—N3	1.363 (5)	N11—H11A	0.8600
C3—N6	1.315 (5)	N11—H11B	0.8600
C3—N1	1.320 (4)	N12—H12A	0.8600
C3—N3	1.347 (5)	N12—H12B	0.8600
C4—N12	1.313 (4)	P1—O4B	1.480 (12)
C4—N7	1.343 (5)	P1—O2B	1.488 (10)
C4—N9	1.350 (4)	P1—O2A	1.498 (4)
C5—N10	1.305 (5)	P1—O4A	1.498 (4)
C5—N7	1.326 (5)	P1—O3A	1.510 (4)
C5—N8	1.355 (5)	P1—O1	1.581 (3)
C6—N11	1.313 (5)	P1—O3B	1.584 (11)
C6—N9	1.323 (4)	O1—H1	0.8200
C6—N8	1.346 (5)	O5—H5C	0.8501
N3—H3	0.77 (4)	O5—H5D	0.8500
N4—H4A	0.8600	O6—H6C	0.8500
N4—H4B	0.8600	O6—H6D	0.8501
N5—H5A	0.8600	O7—H7A	0.8508
N5—H5B	0.8600	O7—H7B	0.8500
N4—C1—N2	116.9 (3)	H6A—N6—H6B	120.0
N4—C1—N1	118.0 (3)	C5—N7—C4	115.4 (3)
N2—C1—N1	125.1 (3)	C6—N8—C5	119.8 (3)
N5—C2—N2	121.1 (3)	C6—N8—H8	118 (2)
N5—C2—N3	118.1 (3)	C5—N8—H8	122 (2)
N2—C2—N3	120.7 (3)	C6—N9—C4	115.6 (3)
N6—C3—N1	120.4 (3)	C5—N10—H10A	120.0
N6—C3—N3	118.3 (3)	C5—N10—H10B	120.0
N1—C3—N3	121.3 (3)	H10A—N10—H10B	120.0
N12—C4—N7	117.0 (3)	C6—N11—H11A	120.0

supplementary materials

N12—C4—N9	116.9 (3)	C6—N11—H11B	120.0
N7—C4—N9	126.1 (3)	H11A—N11—H11B	120.0
N10—C5—N7	122.6 (4)	C4—N12—H12A	120.0
N10—C5—N8	115.9 (3)	C4—N12—H12B	120.0
N7—C5—N8	121.5 (3)	H12A—N12—H12B	120.0
N11—C6—N9	120.4 (3)	O4B—P1—O2B	116.0 (8)
N11—C6—N8	118.1 (3)	O2A—P1—O4A	115.2 (3)
N9—C6—N8	121.5 (3)	O2A—P1—O3A	112.0 (2)
C3—N1—C1	116.4 (3)	O4A—P1—O3A	112.0 (3)
C2—N2—C1	116.6 (3)	O4B—P1—O1	116.8 (5)
C3—N3—C2	119.9 (3)	O2B—P1—O1	97.5 (5)
C3—N3—H3	118 (3)	O2A—P1—O1	107.06 (18)
C2—N3—H3	122 (3)	O4A—P1—O1	102.7 (2)
C1—N4—H4A	120.0	O3A—P1—O1	106.93 (17)
C1—N4—H4B	120.0	O4B—P1—O3B	109.3 (8)
H4A—N4—H4B	120.0	O2B—P1—O3B	108.8 (9)
C2—N5—H5A	120.0	O1—P1—O3B	107.7 (4)
C2—N5—H5B	120.0	P1—O1—H1	109.5
H5A—N5—H5B	120.0	H5C—O5—H5D	107.7
C3—N6—H6A	120.0	H6C—O6—H6D	107.7
C3—N6—H6B	120.0	H7A—O7—H7B	106.2
N6—C3—N1—C1	-179.9 (3)	N10—C5—N7—C4	179.9 (4)
N3—C3—N1—C1	-0.9 (5)	N8—C5—N7—C4	1.0 (5)
N4—C1—N1—C3	177.9 (3)	N12—C4—N7—C5	-179.0 (3)
N2—C1—N1—C3	-1.1 (5)	N9—C4—N7—C5	0.1 (5)
N5—C2—N2—C1	178.3 (3)	N11—C6—N8—C5	179.6 (3)
N3—C2—N2—C1	-1.1 (5)	N9—C6—N8—C5	-0.2 (5)
N4—C1—N2—C2	-176.9 (3)	N10—C5—N8—C6	-180.0 (3)
N1—C1—N2—C2	2.1 (5)	N7—C5—N8—C6	-1.0 (5)
N6—C3—N3—C2	-179.2 (3)	N11—C6—N9—C4	-178.6 (3)
N1—C3—N3—C2	1.8 (5)	N8—C6—N9—C4	1.2 (5)
N5—C2—N3—C3	179.9 (3)	N12—C4—N9—C6	178.0 (3)
N2—C2—N3—C3	-0.8 (5)	N7—C4—N9—C6	-1.2 (5)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N3—H3 \cdots O5 ⁱ	0.77 (4)	2.06 (4)	2.791 (4)	158 (4)
N4—H4B \cdots O4A ⁱⁱ	0.86	2.22	2.998 (5)	151
N4—H4A \cdots N7 ⁱⁱⁱ	0.86	2.20	3.059 (5)	177
N5—H5B \cdots O7	0.86	1.97	2.822 (5)	170
N5—H5A \cdots O6 ^{iv}	0.86	2.41	3.258 (5)	167
N6—H6B \cdots O5 ⁱ	0.86	2.28	3.009 (5)	143
N6—H6A \cdots N9 ^v	0.86	2.17	3.031 (5)	176
N8—H8 \cdots O3A	0.84 (3)	1.93 (4)	2.742 (7)	164 (3)
N10—H10B \cdots O2A	0.86	2.11	2.952 (5)	168
N10—H10A \cdots O2A ^{vi}	0.86	1.99	2.744 (5)	146

N11—H11B···O4A ^{vii}	0.86	2.53	3.147 (6)	129
N11—H11B···O1 ^{vii}	0.86	2.33	3.121 (5)	152
N11—H11A···N1 ^v	0.86	2.02	2.884 (4)	178
N12—H12B···O6 ^{viii}	0.86	2.07	2.840 (5)	149
N12—H12A···N2 ⁱⁱⁱ	0.86	2.08	2.935 (4)	172
O1—H1···O3A ^{vii}	0.82	1.76	2.535 (5)	157
O5—H5D···O3A ⁱ	0.85	1.89	2.739 (5)	177
O5—H5C···O4A	0.85	2.00	2.726 (6)	142
O6—H6D···O1 ^{ix}	0.85	1.97	2.787 (4)	161
O6—H6C···O4A	0.85	2.02	2.838 (6)	162
O7—H7B···O1 ^{ix}	0.85	2.57	3.257 (5)	139
O7—H7B···O2A ^{ix}	0.85	2.28	2.958 (6)	137
O7—H7A···O5 ⁱ	0.85	2.28	3.006 (6)	144

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x, y, z-1$; (iii) $-x+1, -y+2, -z$; (iv) $-x+2, -y+2, -z+1$; (v) $-x, -y+1, -z$; (vi) $-x+1, -y+2, -z+1$; (vii) $-x, -y+1, -z+1$; (viii) $x-1, y, z-1$; (ix) $x+1, y, z$.

Fig. 1




Fig. 2

