

## Bis(1,5-diphenylcarbazonato)dimethanolcobalt(II)

Yanmei Chen,<sup>a</sup> Bin Xu,<sup>a</sup> Shixiong She,<sup>b</sup> Bin Hu<sup>a</sup> and Yahong Li<sup>a,b\*</sup>

<sup>a</sup>Qinghai Institute of Salt Lakes, Chinese Academy of Sciences, Xining 810008, People's Republic of China, and <sup>b</sup>Key Laboratory of Organic Synthesis of Jiangsu Province, College of Chemistry and Chemical Engineering, Suzhou University, Suzhou 215123, People's Republic of China

Correspondence e-mail: liyahong@suda.edu.cn


Received 12 December 2009; accepted 23 December 2009

Key indicators: single-crystal X-ray study;  $T = 296\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$ ;  $R$  factor = 0.042;  $wR$  factor = 0.117; data-to-parameter ratio = 13.1.

The structure of the title compound,  $[\text{Co}(\text{C}_{13}\text{H}_{11}\text{N}_4\text{O})_2(\text{CH}_3\text{OH})_2]$ , is a mononuclear six-coordinated octahedral cobalt(II) complex of  $C_i$  molecular symmetry. The  $\text{Co}^{\text{II}}$  ion is coordinated by two N atoms and two O atoms from two 1,5-biphenylcarbazide ligands, and two O atoms from two methanol molecules. Two diphenylcarbazide ligands and the central  $\text{Co}^{\text{II}}$  ion form the basal plane, with the two methanol molecules located in axial positions. The crystal packing is defined by bifurcated  $\text{O}-\text{H}\cdots\text{N}$  hydrogen bonding and intramolecular  $\text{N}-\text{H}\cdots\text{O}$  interactions.

## Related literature

For the use of biphenylcarbazide for the analytical determination of chromium in biological materials, see: Yarbro & Flaschka (1976). For its coordination modes, see: Feigl (1924); Shafranskii & Mal'kova (1975a,b); Martynova *et al.* (1985); Turkington & Tracy (1958); Deshpande & Jain (1988). For related literature, see: Pankaj & Chauhan (2004); Sollott & Peterson (1969); Cazeneuve (1900a,b).



## Experimental

### Crystal data

$[\text{Co}(\text{C}_{13}\text{H}_{11}\text{N}_4\text{O})_2(\text{CH}_3\text{OH})_2]$   
 $M_r = 601.53$   
Monoclinic,  $P2_1/c$   
 $a = 6.492 (2)\text{ \AA}$   
 $b = 8.926 (3)\text{ \AA}$   
 $c = 25.159 (9)\text{ \AA}$   
 $\beta = 92.372 (6)^\circ$

$V = 1456.7 (9)\text{ \AA}^3$   
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.64\text{ mm}^{-1}$   
 $T = 296\text{ K}$   
 $0.35 \times 0.28 \times 0.27\text{ mm}$

### Data collection

Bruker SMART CCD area-detector diffractometer  
Absorption correction: multi-scan (*SADABS*; Bruker, 2001)  
 $T_{\min} = 0.808$ ,  $T_{\max} = 0.847$

6990 measured reflections  
2564 independent reflections  
2037 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.046$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$   
 $wR(F^2) = 0.117$   
 $S = 1.08$   
2564 reflections  
196 parameters

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.40\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.43\text{ e \AA}^{-3}$

**Table 1**  
Selected bond lengths ( $\text{\AA}$ ).

Co1—O1	2.0263 (18)	Co1—N1	2.193 (2)
Co1—O2	2.214 (2)		

**Table 2**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N4—H4'…O1	0.78 (3)	2.20 (3)	2.587 (3)	111 (2)
O2—H2'…N2 <sup>i</sup>	0.81 (4)	2.11 (4)	2.899 (3)	166 (3)
O2—H2'…N3 <sup>i</sup>	0.81 (4)	2.52 (4)	3.161 (3)	138 (3)

Symmetry code: (i)  $-x + 1, -y + 2, -z + 2$ .

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT* program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors appreciate the financial support of the Hundred Talents Program (2005012) of CAS, the Natural Science Foundation of China (20872105), the 'Qinglan Project' of Jiangsu Province (Bu109805) and the Natural Science Foundation of Qinghai Province (2006-G-105).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: KP2244).

## References

- Bruker (2001). *SMART, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cazeneuve, P. (1900a). *Bull. Soc. Chim. Fr.* **23**, 592–600.
- Cazeneuve, P. (1900b). *C. R. Hebd Séances Acad. Sci.* **130**, 1561–1563.
- Deshpande, S. G. & Jain, S. C. (1988). *Indian J. Chem. Sect. A*, **27**, 552–554.
- Feigl, F. (1924). *Ber. Wien. Akad. IIb*, **133**, 115–132.
- Martynova, T. K., Neverov, V. A., Byushkin, V. N., Shafranskii, V. N. & Malkova, T. A. (1985). *Koord. Khim.*, **11**, 132–135.
- Pankaj, & Chauhan, M. (2004). *Indian J. Chem. Sect. A Inorg. Bio-inorg. Phys. Theor. Anal. Chem.* **43**, 1206–1209.
- Shafranskii, V. N. & Mal'kova, T. A. (1975a). *Zh. Obshch. Khim.* **45**, 1065–1069.
- Shafranskii, V. N. & Mal'kova, T. A. (1975b). *J. Gen. Chem. (USSR)*, **45**, 1051–1054.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sollott, G. P. & Peterson, W. R. (1969). *J. Org. Chem.* **34**, 1506–1508.
- Turkington, R. W. & Tracy, F. M. (1958). *Anal. Chem.* **30**, 1699–1701.
- Yarbro, S. & Flaschka, H. A. (1976). *Microchem. J.* **21**, 415–423.

## **supplementary materials**

*Acta Cryst.* (2010). E66, m114-m115 [doi:10.1107/S1600536809055305]

## Bis(1,5-diphenylcarbazonato)dimethanolcobalt(II)

**Y. Chen, B. Xu, S. She, B. Hu and Y. Li**

### Comment

The diphenylcarbazide is often used for analytical determination of chromium in biological materials (Yarbro *et al.* 1976). As a multidentate ligand, diphenylcarbazide chelates the metal centres by two N atoms (Feigl 1924) or coordinates with the metal ions by O atom in monodentate fashion (Shafranskii *et al.*, 1975a,b; Martynova *et al.*, 1985), whereas the examples of diphenylcarbazide complexes, in which the ligands chelated metal ions bidentately by one O atom and one N atom, were very rare (Turkington *et al.*, 1958; Deshpande *et al.*, 1988). Herein we report the synthesis and crystal structure of such diphenylcarbazide coordinated cobalt complex with Co occupying an inversion centre (Fig. 1 and Table 1). The Packing diagram of **I** viewed down the  $a$  axis (Fig. 2) reveals hydrogen bond interactions (Table 2).

### Experimental

The compound **1** was synthesized by solvothermal reaction. A mixture of diphenylcarbazide (0.0499 g, 0.2 mmol),  $\text{Co}(\text{CH}_3\text{COO})_2 \cdot 4\text{H}_2\text{O}$  (0.0245 g, 0.1 mmol) and  $\text{CH}_3\text{OH} / \text{CH}_3\text{CN}$  ( $v/v = 2:1$ , 2 ml) was sealed in a 5 ml glass tube and heated to 353 K for 48 h. After cooling to room temperature, purple crystals were obtained.

### Refinement

Methyl H atoms were placed in calculated positions with  $\text{C}-\text{H} = 0.96 \text{ \AA}$ , and torsion angles were refined,  $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{C}, \text{O})$ . Other H atoms were placed in calculated positions with  $\text{C}-\text{H} = 0.93$  (aromatic) or  $0.803 \text{ \AA}$  (Imino) and refined in riding mode, with  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ .

### Figures




Fig. 1. Molecular structure showing 50% probability displacement ellipsoids. H atoms are omitted for clarity.




Fig. 2. Packing diagram viewed down the  $a$  axis. Symmetry code corresponds to  $A:-x+2, -y+2, -z+2$ .

# supplementary materials

---

## Bis(1,5-diphenylcarbazono)dimethanolcobalt(II)

### Crystal data

[Co(C <sub>13</sub> H <sub>11</sub> N <sub>4</sub> O) <sub>2</sub> (CH <sub>4</sub> O) <sub>2</sub> ]	$F(000) = 626$
$M_r = 601.53$	$D_x = 1.371 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 2392 reflections
$a = 6.492 (2) \text{ \AA}$	$\theta = 2.4\text{--}24.6^\circ$
$b = 8.926 (3) \text{ \AA}$	$\mu = 0.64 \text{ mm}^{-1}$
$c = 25.159 (9) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 92.372 (6)^\circ$	Block, clear violet
$V = 1456.7 (9) \text{ \AA}^3$	$0.35 \times 0.28 \times 0.27 \text{ mm}$
$Z = 2$	

### Data collection

Bruker SMART CCD area-detector diffractometer	2564 independent reflections
Radiation source: fine-focus sealed tube graphite	2037 reflections with $I > 2\sigma(I)$
phi and $\omega$ scans	$R_{\text{int}} = 0.046$
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2001)	$\theta_{\text{max}} = 25.0^\circ, \theta_{\text{min}} = 1.6^\circ$
$T_{\text{min}} = 0.808, T_{\text{max}} = 0.847$	$h = -7 \rightarrow 7$
6990 measured reflections	$k = -10 \rightarrow 10$
	$l = -20 \rightarrow 29$

### Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.042$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.117$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.08$	$w = 1/[\sigma^2(F_o^2) + (0.0652P)^2 + 0.0401P]$ where $P = (F_o^2 + 2F_c^2)/3$
2564 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
196 parameters	$\Delta\rho_{\text{max}} = 0.40 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.43 \text{ e \AA}^{-3}$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations

between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	1.0000	1.0000	1.0000	0.03524 (19)
C1	0.8530 (5)	0.7374 (3)	1.09426 (11)	0.0528 (7)
H1	0.9758	0.7899	1.0914	0.063*
C2	0.8241 (6)	0.6473 (4)	1.13836 (11)	0.0640 (9)
H2	0.9270	0.6404	1.1651	0.077*
C3	0.6446 (6)	0.5688 (4)	1.14251 (13)	0.0642 (9)
H3	0.6256	0.5087	1.1721	0.077*
C4	0.4933 (5)	0.5785 (4)	1.10342 (14)	0.0720 (10)
H4	0.3721	0.5241	1.1064	0.086*
C5	0.5185 (5)	0.6687 (4)	1.05925 (13)	0.0634 (9)
H5	0.4145	0.6753	1.0328	0.076*
C6	0.7007 (4)	0.7492 (3)	1.05484 (10)	0.0382 (6)
C7	0.6545 (4)	0.9405 (3)	0.93219 (10)	0.0367 (6)
C8	0.4192 (4)	1.0503 (3)	0.80937 (10)	0.0437 (6)
C9	0.2276 (5)	0.9825 (3)	0.80672 (13)	0.0584 (8)
H9	0.1907	0.9157	0.8330	0.070*
C10	0.0906 (6)	1.0150 (4)	0.76451 (14)	0.0710 (10)
H10	-0.0385	0.9697	0.7627	0.085*
C11	0.1441 (6)	1.1139 (4)	0.72520 (13)	0.0719 (10)
H11	0.0509	1.1361	0.6972	0.086*
C12	0.3356 (6)	1.1794 (4)	0.72765 (11)	0.0648 (9)
H12	0.3720	1.2450	0.7009	0.078*
C13	0.4753 (5)	1.1491 (3)	0.76934 (10)	0.0538 (7)
H13	0.6047	1.1940	0.7707	0.065*
C14	0.9161 (6)	1.3027 (4)	1.06416 (16)	0.0849 (12)
H14A	1.0319	1.2726	1.0866	0.127*
H14B	0.8090	1.3405	1.0857	0.127*
H14C	0.9580	1.3797	1.0403	0.127*
N1	0.7412 (3)	0.8471 (2)	1.01157 (8)	0.0359 (5)
N2	0.6030 (3)	0.8460 (2)	0.97330 (8)	0.0391 (5)
N3	0.5120 (3)	0.9360 (3)	0.89297 (8)	0.0411 (5)
N4	0.5597 (4)	1.0247 (3)	0.85196 (9)	0.0463 (6)
O1	0.8187 (3)	1.0213 (2)	0.93292 (7)	0.0438 (5)
O2	0.8423 (3)	1.1797 (2)	1.03493 (9)	0.0570 (6)
H4'	0.660 (5)	1.073 (3)	0.8563 (11)	0.048 (9)*
H2'	0.720 (6)	1.167 (4)	1.0382 (13)	0.084 (12)*

## supplementary materials

---

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Co1	0.0269 (3)	0.0445 (3)	0.0341 (3)	-0.0019 (2)	-0.00083 (18)	-0.0004 (2)
C1	0.0589 (19)	0.0546 (17)	0.0445 (16)	-0.0123 (14)	-0.0024 (14)	0.0054 (13)
C2	0.088 (3)	0.0600 (19)	0.0428 (16)	-0.0077 (18)	-0.0077 (16)	0.0091 (14)
C3	0.086 (3)	0.0534 (18)	0.0543 (19)	-0.0007 (18)	0.0204 (18)	0.0125 (15)
C4	0.057 (2)	0.074 (2)	0.086 (3)	-0.0096 (18)	0.0160 (19)	0.029 (2)
C5	0.0419 (17)	0.076 (2)	0.073 (2)	-0.0062 (16)	0.0013 (15)	0.0258 (17)
C6	0.0402 (14)	0.0368 (13)	0.0382 (14)	0.0022 (11)	0.0080 (11)	-0.0002 (11)
C7	0.0284 (13)	0.0454 (13)	0.0362 (13)	0.0017 (11)	0.0003 (10)	-0.0023 (11)
C8	0.0474 (17)	0.0479 (15)	0.0352 (14)	0.0059 (12)	-0.0044 (12)	-0.0064 (11)
C9	0.064 (2)	0.0593 (19)	0.0508 (18)	-0.0049 (15)	-0.0145 (15)	0.0038 (14)
C10	0.066 (2)	0.078 (2)	0.066 (2)	-0.0084 (18)	-0.0298 (17)	0.0024 (18)
C11	0.087 (3)	0.071 (2)	0.0549 (19)	0.013 (2)	-0.0312 (18)	-0.0007 (18)
C12	0.093 (3)	0.0605 (19)	0.0408 (17)	0.0102 (18)	-0.0044 (16)	0.0036 (14)
C13	0.0610 (19)	0.0583 (18)	0.0420 (15)	0.0018 (15)	0.0001 (13)	-0.0031 (13)
C14	0.058 (2)	0.084 (3)	0.113 (3)	-0.0036 (19)	0.015 (2)	-0.047 (2)
N1	0.0296 (11)	0.0426 (12)	0.0355 (11)	0.0026 (9)	0.0028 (9)	-0.0024 (9)
N2	0.0313 (11)	0.0472 (13)	0.0386 (12)	0.0003 (9)	0.0013 (9)	-0.0009 (10)
N3	0.0342 (12)	0.0524 (13)	0.0365 (12)	-0.0015 (10)	-0.0017 (9)	-0.0010 (10)
N4	0.0397 (14)	0.0592 (16)	0.0394 (13)	-0.0065 (12)	-0.0059 (10)	0.0034 (11)
O1	0.0355 (10)	0.0560 (12)	0.0395 (10)	-0.0074 (9)	-0.0035 (8)	0.0049 (8)
O2	0.0295 (11)	0.0635 (13)	0.0786 (15)	-0.0035 (10)	0.0100 (10)	-0.0232 (11)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Co1—O1 <sup>i</sup>	2.0263 (18)	C8—C9	1.383 (4)
Co1—O1	2.0263 (18)	C8—N4	1.397 (3)
Co1—O2	2.114 (2)	C8—C13	1.398 (4)
Co1—O2 <sup>i</sup>	2.114 (2)	C9—C10	1.387 (4)
Co1—N1 <sup>i</sup>	2.193 (2)	C9—H9	0.9300
Co1—N1	2.193 (2)	C10—C11	1.381 (5)
C1—C6	1.375 (4)	C10—H10	0.9300
C1—C2	1.389 (4)	C11—C12	1.373 (5)
C1—H1	0.9300	C11—H11	0.9300
C2—C3	1.368 (5)	C12—C13	1.384 (4)
C2—H2	0.9300	C12—H12	0.9300
C3—C4	1.363 (5)	C13—H13	0.9300
C3—H3	0.9300	C14—O2	1.395 (4)
C4—C5	1.387 (4)	C14—H14A	0.9600
C4—H4	0.9300	C14—H14B	0.9600
C5—C6	1.392 (4)	C14—H14C	0.9600
C5—H5	0.9300	N1—N2	1.289 (3)
C6—N1	1.429 (3)	N3—N4	1.347 (3)
C7—O1	1.286 (3)	N4—H4'	0.78 (3)
C7—N3	1.325 (3)	O2—H2'	0.81 (4)

C7—N2	1.386 (3)		
O1 <sup>i</sup> —Co1—O1	179.999 (1)	C9—C8—N4	121.6 (3)
O1 <sup>i</sup> —Co1—O2	89.97 (8)	C9—C8—C13	120.1 (3)
O1—Co1—O2	90.03 (8)	N4—C8—C13	118.3 (3)
O1 <sup>i</sup> —Co1—O2 <sup>i</sup>	90.03 (8)	C8—C9—C10	119.5 (3)
O1—Co1—O2 <sup>i</sup>	89.97 (8)	C8—C9—H9	120.2
O2—Co1—O2 <sup>i</sup>	179.997 (1)	C10—C9—H9	120.2
O1 <sup>i</sup> —Co1—N1 <sup>i</sup>	75.34 (7)	C11—C10—C9	120.7 (3)
O1—Co1—N1 <sup>i</sup>	104.66 (7)	C11—C10—H10	119.7
O2—Co1—N1 <sup>i</sup>	88.27 (8)	C9—C10—H10	119.7
O2 <sup>i</sup> —Co1—N1 <sup>i</sup>	91.73 (8)	C12—C11—C10	119.6 (3)
O1 <sup>i</sup> —Co1—N1	104.66 (7)	C12—C11—H11	120.2
O1—Co1—N1	75.34 (7)	C10—C11—H11	120.2
O2—Co1—N1	91.73 (8)	C11—C12—C13	121.0 (3)
O2 <sup>i</sup> —Co1—N1	88.27 (8)	C11—C12—H12	119.5
N1 <sup>i</sup> —Co1—N1	180.0	C13—C12—H12	119.5
C6—C1—C2	120.2 (3)	C12—C13—C8	119.1 (3)
C6—C1—H1	119.9	C12—C13—H13	120.4
C2—C1—H1	119.9	C8—C13—H13	120.4
C3—C2—C1	120.1 (3)	O2—C14—H14A	109.5
C3—C2—H2	120.0	O2—C14—H14B	109.5
C1—C2—H2	120.0	H14A—C14—H14B	109.5
C4—C3—C2	120.2 (3)	O2—C14—H14C	109.5
C4—C3—H3	119.9	H14A—C14—H14C	109.5
C2—C3—H3	119.9	H14B—C14—H14C	109.5
C3—C4—C5	120.6 (3)	N2—N1—C6	114.8 (2)
C3—C4—H4	119.7	N2—N1—Co1	114.78 (15)
C5—C4—H4	119.7	C6—N1—Co1	130.38 (16)
C4—C5—C6	119.5 (3)	N1—N2—C7	111.8 (2)
C4—C5—H5	120.3	C7—N3—N4	112.2 (2)
C6—C5—H5	120.3	N3—N4—C8	121.3 (3)
C1—C6—C5	119.4 (3)	N3—N4—H4'	116 (2)
C1—C6—N1	116.5 (2)	C8—N4—H4'	122 (2)
C5—C6—N1	124.1 (3)	C7—O1—Co1	114.33 (15)
O1—C7—N3	125.5 (2)	C14—O2—Co1	130.91 (19)
O1—C7—N2	123.8 (2)	C14—O2—H2'	112 (3)
N3—C7—N2	110.8 (2)	Co1—O2—H2'	116 (3)
C6—C1—C2—C3	0.6 (5)	O1—Co1—N1—C6	178.9 (2)
C1—C2—C3—C4	0.1 (5)	O2—Co1—N1—C6	-91.5 (2)
C2—C3—C4—C5	-0.5 (6)	O2 <sup>i</sup> —Co1—N1—C6	88.5 (2)
C3—C4—C5—C6	0.4 (5)	N1 <sup>i</sup> —Co1—N1—C6	-89 (10)
C2—C1—C6—C5	-0.8 (4)	C6—N1—N2—C7	-178.5 (2)
C2—C1—C6—N1	178.5 (3)	Co1—N1—N2—C7	1.8 (2)
C4—C5—C6—C1	0.3 (5)	O1—C7—N2—N1	-1.2 (3)
C4—C5—C6—N1	-178.9 (3)	N3—C7—N2—N1	179.1 (2)
N4—C8—C9—C10	177.9 (3)	O1—C7—N3—N4	1.5 (4)

## supplementary materials

---

C13—C8—C9—C10	−0.9 (5)	N2—C7—N3—N4	−178.9 (2)
C8—C9—C10—C11	0.1 (5)	C7—N3—N4—C8	−172.2 (2)
C9—C10—C11—C12	0.7 (5)	C9—C8—N4—N3	−1.8 (4)
C10—C11—C12—C13	−0.8 (5)	C13—C8—N4—N3	177.1 (3)
C11—C12—C13—C8	0.0 (5)	N3—C7—O1—Co1	179.6 (2)
C9—C8—C13—C12	0.9 (4)	N2—C7—O1—Co1	−0.1 (3)
N4—C8—C13—C12	−178.0 (3)	O1 <sup>i</sup> —Co1—O1—C7	129 (3)
C1—C6—N1—N2	174.5 (2)	O2—Co1—O1—C7	−90.99 (18)
C5—C6—N1—N2	−6.3 (4)	O2 <sup>i</sup> —Co1—O1—C7	89.01 (18)
C1—C6—N1—Co1	−5.9 (3)	N1 <sup>i</sup> —Co1—O1—C7	−179.21 (17)
C5—C6—N1—Co1	173.3 (2)	N1—Co1—O1—C7	0.79 (17)
O1 <sup>i</sup> —Co1—N1—N2	178.49 (15)	O1 <sup>i</sup> —Co1—O2—C14	44.4 (3)
O1—Co1—N1—N2	−1.51 (15)	O1—Co1—O2—C14	−135.6 (3)
O2—Co1—N1—N2	88.07 (17)	O2 <sup>i</sup> —Co1—O2—C14	−89 (4)
O2 <sup>i</sup> —Co1—N1—N2	−91.93 (17)	N1 <sup>i</sup> —Co1—O2—C14	−30.9 (3)
N1 <sup>i</sup> —Co1—N1—N2	91 (10)	N1—Co1—O2—C14	149.1 (3)
O1 <sup>i</sup> —Co1—N1—C6	−1.1 (2)		


Symmetry codes: (i)  $-x+2, -y+2, -z+2$ .

### Hydrogen-bond geometry ( $\text{\AA}$ , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N4—H4'…O1	0.78 (3)	2.20 (3)	2.587 (3)	111 (2)
O2—H2'…N2 <sup>ii</sup>	0.81 (4)	2.11 (4)	2.899 (3)	166 (3)
O2—H2'…N3 <sup>ii</sup>	0.81 (4)	2.52 (4)	3.161 (3)	138 (3)

Symmetry codes: (ii)  $-x+1, -y+2, -z+2$ .

Fig. 1



## **supplementary materials**

---

**Fig. 2**

