

(2,2'-Bipyridine- κ^2N,N')bis(N -ethyl- N -methylidithiocarbamato- κ^2S,S')zinc(II)

Noorul Aisyah Abdul Ghafar,^a Ibrahim Baba,^a Bohari M. Yamin^a and Seik Weng Ng^{b*}

^aSchool of Chemical Sciences, Universiti Kebangsaan Malaysia, 43600 Bangi, Malaysia, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 19 January 2010; accepted 20 January 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; disorder in main residue; R factor = 0.030; wR factor = 0.091; data-to-parameter ratio = 18.5.

The complete molecule of the title compound, $[\text{Zn}(\text{C}_4\text{H}_8\text{NS}_2)_2(\text{C}_{10}\text{H}_8\text{N}_2)]$, is generated by crystallographic twofold symmetry, with the Zn atom lying on the rotation axis; the axis also bisects the central C–C bond of the 2,2'-bipyridine molecule. The metal atom is chelated by two S,S' -bidentate dithiocarbamate anions and the N,N' -bidentate heterocycle, resulting in a distorted *cis*-ZnN₂S₄ octahedral geometry. The methyl and ethyl groups of the anion are statistically disordered.

Related literature

For other 2,2'-bipyridine adducts of zinc dithio carbamates, see: Ali *et al.* (2006); Deng *et al.* (2007); Jie & Tiekkink (2002); Lai & Tiekkink (2004); Manohar *et al.* (1998); Thirumaran *et al.* (1999); Yin *et al.* (2004); Zemskova *et al.* (1993).



Experimental

Crystal data

$[\text{Zn}(\text{C}_4\text{H}_8\text{NS}_2)_2(\text{C}_{10}\text{H}_8\text{N}_2)]$

$M_r = 490.02$

Orthorhombic, $Pnaa$
 $a = 16.9478 (7)\text{ \AA}$
 $b = 19.3282 (8)\text{ \AA}$
 $c = 6.6572 (3)\text{ \AA}$
 $V = 2180.70 (16)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 1.52\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.45 \times 0.40 \times 0.35\text{ mm}$

Data collection

Bruker SMART APEX CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.548$, $T_{\max} = 0.618$

13725 measured reflections
2513 independent reflections
2252 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.091$
 $S = 1.04$
2513 reflections
136 parameters

14 restraints
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.35\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.22\text{ e \AA}^{-3}$

Table 1
Selected geometric parameters (\AA , $^\circ$).

Zn1–N2	2.1742 (15)	Zn1–S1	2.5261 (6)
Zn1–S2	2.5259 (5)		
N2–Zn1–N2 ⁱ	75.24 (8)	S2–Zn1–S1	70.884 (17)
Symmetry code: (i) $x, -y + \frac{1}{2}, -z + \frac{1}{2}$			

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank Universiti Kebangsaan Malaysia (UKM-GUP-NBT-08-27-111 and 06-01-02-SF0539) and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5313).

References

- Ali, B. F., Al-Sou'od, K. A., Al-Far, R. & Judeh, Z. (2006). *Struct. Chem.* **17**, 423–429.
- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2000). *SAINT* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Deng, Y.-H., Liu, J., Li, N., Yang, Y.-L. & Ma, H.-W. (2007). *Huaxue Xuebao*, **65**, 2868–2874.
- Jie, Q. & Tiekkink, E. R. T. (2002). *Main Group Met. Chem.* **25**, 317–318.
- Lai, C. S. & Tiekkink, E. R. T. (2004). *Appl. Organomet. Chem.* **18**, 197–198.
- Manohar, A., Venkatachalam, V., Ramalingam, K., Thirumaran, S., Bocelli, G. & Cantoni, A. (1998). *J. Chem. Crystallogr.* **28**, 861–866.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Thirumaran, S., Ramalingam, K., Bocelli, G. & Cantoni, A. (1999). *Polyhedron*, **18**, 925–930.
- Westrip, S. P. (2010). *publCIF*. In preparation.
- Yin, H.-D., Wang, C.-H. & Xing, Q.-J. (2004). *Indian J. Chem. Sect. A*, **43**, 1911–1914.
- Zemskova, S. M., Glinskaya, L. A., Durasov, V. B., Klevtsova, R. F. & Larionov, S. V. (1993). *Zh. Strukt. Khim.* **34**, 155–157.

supplementary materials

Acta Cryst. (2010). E66, m208 [doi:10.1107/S1600536810002606]

(2,2'-Bipyridine- κ^2N,N')bis(*N*-ethyl-*N*-methyldithiocarbamato- κ^2S,S')zinc(II)

N. A. Abdul Ghafar, I. Baba, B. M. Yamin and S. W. Ng

Experimental

Zinc chloride (10 mmol), ethylmethylamine (20 mmol), carbon disulfide (20 mmol), 2,2'-bipyridine and ammonia (10 ml) were reacted in ethanol (30 ml) at 277 K to produce a white solid. This was collected and recrystallized from ethanol to yield colourless blocks of (I).

Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.93 to 0.97 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to 1.2 to 1.5 $U(C)$. The methyl group is disordered with respect to the ethyl group. Both were refined as ethyl groups, but the methyl carbon atoms were refined with 0.5 occupancy each. The carbon–carbon distance was restrained to 1.50±0.01 Å; the anisotropic temperature factors of the half-occupancy atoms were restrained to be nearly isotropic.

Figures




Fig. 1. View of (I) at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius. The disorder is not shown.

(2,2'-Bipyridine- κ^2N,N')bis(*N*-ethyl-*N*- methyldithiocarbamato- κ^2S,S')zinc(II)

Crystal data

[Zn(C ₄ H ₈ NS ₂) ₂ (C ₁₀ H ₈ N ₂)]	$F(000) = 1016$
$M_r = 490.02$	$D_x = 1.493 \text{ Mg m}^{-3}$
Orthorhombic, <i>Pnaa</i>	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ac 2bc	Cell parameters from 6268 reflections
$a = 16.9478 (7) \text{ \AA}$	$\theta = 2.4\text{--}27.5^\circ$
$b = 19.3282 (8) \text{ \AA}$	$\mu = 1.52 \text{ mm}^{-1}$
$c = 6.6572 (3) \text{ \AA}$	$T = 293 \text{ K}$
$V = 2180.70 (16) \text{ \AA}^3$	Block, colorless
$Z = 4$	$0.45 \times 0.40 \times 0.35 \text{ mm}$

supplementary materials

Data collection

Bruker SMART APEX CCD diffractometer	2513 independent reflections
Radiation source: fine-focus sealed tube graphite	2252 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.019$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 2.1^\circ$
$T_{\text{min}} = 0.548, T_{\text{max}} = 0.618$	$h = -20 \rightarrow 22$
13725 measured reflections	$k = -25 \rightarrow 24$
	$l = -5 \rightarrow 8$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.030$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.091$	H-atom parameters constrained
$S = 1.04$	$w = 1/[\sigma^2(F_o^2) + (0.0554P)^2 + 0.6285P]$ where $P = (F_o^2 + 2F_c^2)/3$
2513 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
136 parameters	$\Delta\rho_{\text{max}} = 0.35 \text{ e \AA}^{-3}$
14 restraints	$\Delta\rho_{\text{min}} = -0.22 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Zn1	0.292606 (17)	0.2500	0.2500	0.03998 (12)	
S1	0.27352 (4)	0.14351 (3)	0.03544 (9)	0.05648 (17)	
S2	0.19702 (3)	0.16932 (3)	0.42289 (8)	0.04706 (15)	
N1	0.16951 (11)	0.05783 (10)	0.2003 (3)	0.0545 (4)	
N2	0.39422 (8)	0.21503 (8)	0.4216 (2)	0.0379 (3)	
C1	0.20922 (11)	0.11716 (11)	0.2174 (3)	0.0424 (4)	
C2	0.18288 (18)	0.00855 (16)	0.0390 (5)	0.0884 (10)	0.50
H2A	0.2286	0.0238	-0.0367	0.106*	0.50
H2B	0.1961	-0.0357	0.0990	0.106*	0.50
C3	0.1206 (3)	-0.0020 (3)	-0.0968 (8)	0.0731 (15)	0.50
H3A	0.1326	-0.0407	-0.1819	0.110*	0.50
H3B	0.1136	0.0387	-0.1775	0.110*	0.50
H3C	0.0729	-0.0113	-0.0237	0.110*	0.50
C2'	0.18288 (18)	0.00855 (16)	0.0390 (5)	0.0884 (10)	0.50
H2'A	0.1652	-0.0364	0.0804	0.106*	0.50
H2'B	0.2382	0.0068	0.0082	0.106*	0.50
H2'C	0.1541	0.0227	-0.0781	0.106*	0.50
C4	0.11331 (16)	0.03509 (15)	0.3511 (4)	0.0725 (7)	0.50

H4A	0.1398	0.0296	0.4775	0.087*	0.50
H4B	0.0908	-0.0083	0.3109	0.087*	0.50
H4C	0.0722	0.0690	0.3644	0.087*	0.50
C4'	0.11331 (16)	0.03509 (15)	0.3511 (4)	0.0725 (7)	0.50
H4'A	0.1228	-0.0134	0.3796	0.087*	0.50
H4'B	0.1228	0.0608	0.4739	0.087*	0.50
C5'	0.0321 (4)	0.0434 (5)	0.2957 (14)	0.114 (3)	0.50
H5'A	-0.0004	0.0154	0.3816	0.171*	0.50
H5'B	0.0249	0.0293	0.1587	0.171*	0.50
H5'C	0.0173	0.0911	0.3100	0.171*	0.50
C6	0.39024 (12)	0.18074 (10)	0.5947 (3)	0.0453 (4)	
H6	0.3408	0.1703	0.6470	0.054*	
C7	0.45602 (13)	0.16000 (11)	0.6997 (3)	0.0492 (5)	
H7	0.4511	0.1362	0.8205	0.059*	
C8	0.52930 (13)	0.17520 (11)	0.6222 (3)	0.0512 (5)	
H8	0.5748	0.1617	0.6897	0.061*	
C9	0.53413 (11)	0.21061 (11)	0.4434 (3)	0.0468 (4)	
H9	0.5831	0.2212	0.3882	0.056*	
C10	0.46553 (10)	0.23031 (9)	0.3464 (3)	0.0367 (4)	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.03265 (18)	0.04134 (19)	0.0460 (2)	0.000	0.000	0.00225 (12)
S1	0.0583 (3)	0.0555 (3)	0.0557 (3)	-0.0150 (2)	0.0193 (3)	-0.0093 (2)
S2	0.0409 (3)	0.0566 (3)	0.0437 (3)	-0.0025 (2)	0.00315 (19)	-0.0026 (2)
N1	0.0479 (10)	0.0455 (9)	0.0701 (12)	-0.0080 (8)	0.0102 (9)	-0.0039 (8)
N2	0.0354 (7)	0.0381 (7)	0.0402 (8)	-0.0013 (6)	0.0004 (6)	0.0002 (6)
C1	0.0357 (9)	0.0422 (10)	0.0493 (10)	0.0005 (7)	0.0022 (7)	0.0024 (8)
C2	0.0759 (18)	0.0659 (16)	0.123 (3)	-0.0218 (14)	0.0324 (18)	-0.0391 (18)
C3	0.097 (4)	0.063 (3)	0.059 (3)	0.017 (3)	-0.009 (3)	-0.012 (2)
C2'	0.0759 (18)	0.0659 (16)	0.123 (3)	-0.0218 (14)	0.0324 (18)	-0.0391 (18)
C4	0.0698 (16)	0.0695 (15)	0.0783 (17)	-0.0251 (13)	0.0123 (14)	0.0122 (13)
C4'	0.0698 (16)	0.0695 (15)	0.0783 (17)	-0.0251 (13)	0.0123 (14)	0.0122 (13)
C5'	0.086 (4)	0.124 (6)	0.132 (6)	-0.010 (4)	0.025 (4)	0.026 (5)
C6	0.0455 (10)	0.0459 (10)	0.0444 (10)	-0.0039 (8)	0.0010 (8)	0.0047 (8)
C7	0.0581 (12)	0.0442 (10)	0.0453 (10)	-0.0011 (8)	-0.0068 (9)	0.0067 (9)
C8	0.0490 (11)	0.0497 (11)	0.0549 (12)	0.0086 (9)	-0.0128 (10)	0.0034 (9)
C9	0.0347 (9)	0.0524 (11)	0.0533 (11)	0.0049 (8)	-0.0018 (8)	0.0007 (9)
C10	0.0342 (8)	0.0355 (8)	0.0402 (10)	0.0007 (6)	-0.0008 (7)	-0.0024 (7)

Geometric parameters (\AA , $^\circ$)

Zn1—N2	2.1742 (15)	C3—H3B	0.9600
Zn1—N2 ⁱ	2.1742 (15)	C3—H3C	0.9600
Zn1—S2	2.5259 (5)	C4—H4A	0.9600
Zn1—S2 ⁱ	2.5259 (5)	C4—H4B	0.9600
Zn1—S1	2.5261 (6)	C4—H4C	0.9600

supplementary materials

Zn1—S1 ⁱ	2.5261 (6)	C5'—H5'A	0.9600
S1—C1	1.707 (2)	C5'—H5'B	0.9600
S2—C1	1.712 (2)	C5'—H5'C	0.9600
N1—C1	1.334 (3)	C6—C7	1.376 (3)
N1—C4	1.452 (3)	C6—H6	0.9300
N1—C2	1.453 (3)	C7—C8	1.377 (3)
N2—C6	1.331 (2)	C7—H7	0.9300
N2—C10	1.341 (2)	C8—C9	1.375 (3)
C2—C3	1.405 (5)	C8—H8	0.9300
C2—H2A	0.9700	C9—C10	1.384 (3)
C2—H2B	0.9700	C9—H9	0.9300
C3—H3A	0.9600	C10—C10 ⁱ	1.492 (4)
N2—Zn1—N2 ⁱ	75.24 (8)	C3—C2—H2A	108.1
N2—Zn1—S2	94.40 (4)	N1—C2—H2A	108.1
N2 ⁱ —Zn1—S2	159.93 (4)	C3—C2—H2B	108.1
N2—Zn1—S2 ⁱ	159.93 (4)	N1—C2—H2B	108.1
N2 ⁱ —Zn1—S2 ⁱ	94.40 (4)	H2A—C2—H2B	107.3
S2—Zn1—S2 ⁱ	100.22 (3)	N1—C4—H4A	109.5
N2—Zn1—S1	98.35 (4)	N1—C4—H4B	109.5
N2 ⁱ —Zn1—S1	93.31 (4)	N1—C4—H4C	109.5
S2—Zn1—S1	70.884 (17)	H5'A—C5'—H5'B	109.5
S2 ⁱ —Zn1—S1	99.39 (2)	H5'A—C5'—H5'C	109.5
N2—Zn1—S1 ⁱ	93.31 (4)	H5'B—C5'—H5'C	109.5
N2 ⁱ —Zn1—S1 ⁱ	98.35 (4)	N2—C6—C7	122.96 (18)
S2—Zn1—S1 ⁱ	99.39 (2)	N2—C6—H6	118.5
S2 ⁱ —Zn1—S1 ⁱ	70.884 (17)	C7—C6—H6	118.5
S1—Zn1—S1 ⁱ	165.28 (3)	C8—C7—C6	118.6 (2)
C1—S1—Zn1	85.62 (7)	C8—C7—H7	120.7
C1—S2—Zn1	85.53 (7)	C6—C7—H7	120.7
C1—N1—C4	122.2 (2)	C7—C8—C9	118.97 (19)
C1—N1—C2	123.2 (2)	C7—C8—H8	120.5
C4—N1—C2	114.5 (2)	C9—C8—H8	120.5
C6—N2—C10	118.59 (16)	C8—C9—C10	119.40 (19)
C6—N2—Zn1	124.71 (12)	C8—C9—H9	120.3
C10—N2—Zn1	116.70 (12)	C10—C9—H9	120.3
N1—C1—S2	120.88 (16)	N2—C10—C9	121.49 (18)
N1—C1—S1	121.19 (16)	N2—C10—C10 ⁱ	115.69 (10)
S2—C1—S1	117.93 (12)	C9—C10—C10 ⁱ	122.82 (12)
C3—C2—N1	117.0 (3)		
N2—Zn1—S1—C1	92.98 (8)	C2—N1—C1—S2	-174.1 (2)
N2 ⁱ —Zn1—S1—C1	168.53 (8)	C4—N1—C1—S1	-178.68 (19)
S2—Zn1—S1—C1	1.19 (7)	C2—N1—C1—S1	6.0 (3)
S2 ⁱ —Zn1—S1—C1	-96.46 (7)	Zn1—S2—C1—N1	-178.01 (18)
S1 ⁱ —Zn1—S1—C1	-49.05 (7)	Zn1—S2—C1—S1	1.87 (11)
N2—Zn1—S2—C1	-98.50 (8)	Zn1—S1—C1—N1	178.01 (18)

N2 ⁱ —Zn1—S2—C1	−40.79 (14)	Zn1—S1—C1—S2	−1.87 (11)
S2 ⁱ —Zn1—S2—C1	95.31 (7)	C1—N1—C2—C3	−114.6 (4)
S1—Zn1—S2—C1	−1.18 (7)	C4—N1—C2—C3	69.8 (4)
S1 ⁱ —Zn1—S2—C1	167.40 (7)	C10—N2—C6—C7	−0.3 (3)
N2 ⁱ —Zn1—N2—C6	179.12 (19)	Zn1—N2—C6—C7	−179.41 (16)
S2—Zn1—N2—C6	−18.34 (15)	N2—C6—C7—C8	−0.2 (3)
S2 ⁱ —Zn1—N2—C6	118.46 (16)	C6—C7—C8—C9	0.2 (3)
S1—Zn1—N2—C6	−89.64 (15)	C7—C8—C9—C10	0.3 (3)
S1 ⁱ —Zn1—N2—C6	81.35 (15)	C6—N2—C10—C9	0.7 (3)
N2 ⁱ —Zn1—N2—C10	−0.03 (10)	Zn1—N2—C10—C9	179.93 (14)
S2—Zn1—N2—C10	162.50 (13)	C6—N2—C10—C10 ⁱ	−179.12 (19)
S2 ⁱ —Zn1—N2—C10	−60.7 (2)	Zn1—N2—C10—C10 ⁱ	0.1 (3)
S1—Zn1—N2—C10	91.21 (13)	C8—C9—C10—N2	−0.7 (3)
S1 ⁱ —Zn1—N2—C10	−97.80 (13)	C8—C9—C10—C10 ⁱ	179.1 (2)
C4—N1—C1—S2	1.2 (3)		

Symmetry codes: (i) $x, -y+1/2, -z+1/2$.

supplementary materials

Fig. 1

