

1-(2,6-Dichlorobenzoyl)-3-(3-nitrophenyl)thiourea dimethylformamide solvate

Min Li^{a*} and Dongxiao Hou^b

^aDepartment of Chemistry, Changzhi University, Shanxi, People's Republic of China, and ^bInstitute of Applied Chemistry, Shanxi University, Shanxi, People's Republic of China

Correspondence e-mail: limin0081.student@sina.com


Received 1 December 2009; accepted 22 December 2009

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.046; wR factor = 0.139; data-to-parameter ratio = 13.3.

In the title compound, $\text{C}_{14}\text{H}_9\text{Cl}_2\text{N}_3\text{O}_3\text{S} \cdot \text{C}_3\text{H}_7\text{NO}$, the two aromatic rings enclose a dihedral angle of $32.93(12)^\circ$. The thiourea molecule exists in its thione form in the solid state with typical $\text{C}=\text{S}$ and $\text{C}-\text{N}$ bond lengths. In the crystal, $\text{N}-\text{H} \cdots \text{O}$ hydrogen bonds exist between the thiourea and carbonyl groups on the same and neighboring molecules. In addition, each dimethylformamide solvate molecule forms a hydrogen bond to one N atom of the thiourea group.

Related literature

For general background to the use of thiourea and urea derivatives in the development of agrochemicals and pharmacological agents, see: Darlington *et al.* (1996); Dowding & Leeds (1971); Sasse *et al.* (1969). For bond lengths in other substituted thioureas, see: Khawar Rauf *et al.* (2006a,*b,c*, 2007, 2009). For previously reported $\text{C}=\text{S}$ distances, see: Bailey *et al.* (1997).



Experimental

Crystal data

$\text{C}_{14}\text{H}_9\text{Cl}_2\text{N}_3\text{O}_3\text{S} \cdot \text{C}_3\text{H}_7\text{NO}$
 $M_r = 443.30$
Triclinic, $P\bar{1}$

$a = 8.507(5)\text{ \AA}$
 $b = 10.240(5)\text{ \AA}$
 $c = 12.414(8)\text{ \AA}$

$\alpha = 70.40(4)^\circ$
 $\beta = 81.74(5)^\circ$
 $\gamma = 87.98(4)^\circ$
 $V = 1008(1)\text{ \AA}^3$
 $Z = 2$

Mo $K\alpha$ radiation
 $\mu = 0.46\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.30 \times 0.20 \times 0.20\text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1997)
 $(SADABS)$: Sheldrick, 1997)
 $T_{\min} = 0.875$, $T_{\max} = 0.914$

4551 measured reflections
3413 independent reflections
2691 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.139$
 $S = 1.11$
3413 reflections

256 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.30\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.42\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
N2—H2 \cdots O1 ⁱ	0.86	2.47	3.182 (3)	141
N2—H2 \cdots O1	0.86	1.99	2.675 (3)	136
N1—H1 \cdots O4 ⁱⁱ	0.86	1.96	2.787 (3)	161

Symmetry codes: (i) $-x + 1, -y + 1, -z + 2$; (ii) $-x + 1, -y + 2, -z + 1$.

Data collection: *SMART* (Siemens, 1996); cell refinement: *SAINT* (Siemens, 1996); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: EZ2196).

References

- Bailey, P. J., Grant, K. J. & Parsons, S. (1997). *Acta Cryst. C53*, 247–248.
Darlington, A., Vishnevetskia, K. & Blake, T. T. (1996). *Physiol. Plant.* **97**, 217–222.
Dowding, J. & Leeds, W. G. (1971). Ger. Patent No. 2 040 580.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Khawar Rauf, M., Badshah, A. & Bolte, M. (2006a). *Acta Cryst. E62*, o1859–o1860.
Khawar Rauf, M., Badshah, A. & Bolte, M. (2006b). *Acta Cryst. E62*, o2221–o2222.
Khawar Rauf, M., Badshah, A. & Bolte, M. (2006c). *Acta Cryst. E62*, o2444–o2445.
Khawar Rauf, M., Bolte, M. & Badshah, A. (2009). *Acta Cryst. E65*, o142.
Khawar Rauf, M., Badshah, A., Bolte, M. & Ahmad, I. (2007). *Acta Cryst. E63*, o1073–o1075.
Sasse, K., Barden, R., Eue, L. & Hack, H. (1969). S. Afr. Patent No. 900 256.
Sheldrick, G. M. (1997). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
Siemens (1996). *SMART* and *SAINT*. Siemens Analytical X-ray Instruments Inc., Madison Wisconsin, USA.

supplementary materials

Acta Cryst. (2010). E66, o280 [doi:10.1107/S1600536809055056]

1-(2,6-Dichlorobenzoyl)-3-(3-nitrophenyl)thiourea dimethylformamide solvate

M. Li and D. Hou

Comment

Earlier studies have shown that thiourea and urea derivatives have played an important role in developing agrochemicals and pharmacological agents (Dowding & Leeds, 1971; Sasse *et al.*, 1969; Darlington *et al.*, 1996). As part of our interest in N,N'-disubstituted thioureas, we now report the crystal structure of the title compound (I).

The N—C bonds in (I), see Fig. 1, differ significantly from one another but are short in comparison with the typical value for an N—C single bond (1.479 Å). Owing to the introduction of the C=O electron-acceptor group the adjacent C—S bond length [1.652 (2) Å] is shorter than previously reported C=S distances (1.710 (7) Å) (Bailey, *et al.*, 1997). These distances are similar to those usually found in other substituted thioureas (Khawar Rauf *et al.*, 2006a, 2006b, 2006c, 2007, 2009). The dihedral angle between the aromatic rings is 32.93 (12)°, and the corresponding angles with the thiourea plane are 83.52 (7)° for the C2—C7 ring and 50.61 (7)° for the C9—C14 ring. The thiocarbonyl and carbonyl groups are almost coplanar.

Inter- and intramolecular N—H···O hydrogen bonds exist between the thiourea N—H atoms and carbonyl-O atoms. In addition, each dimethylformamide solvate molecule also has a hydrogen bond to the N of the thiourea groups (2.787 (3) Å; Table 1, Fig. 2).

Experimental

Freshly prepared 2,6-dichlorobenzoylisothiocyanate (2.32 g, 10 mmol) was added to dimethylformamide (30 ml) and stirred for 2 minutes. Afterwards neat 3-nitroaniline (1.38 g, 10 mmol) was added and the resulting mixture was stirred for 1 h. The reaction mixture was then poured into an ice-water mixture and stirred well. The solid product was separated and washed with deionized water and purified by recrystallization from methanol/CH₂Cl₂ (1:1 v/v) to give fine crystals of the title compound (I), with an overall yield of 85%.

Refinement

All H atoms were positioned geometrically, with C—H = 0.96–0.98 Å, and refined as riding, allowing for free rotation of the methyl groups. The $U_{\text{iso}}(\text{H})$ values were set at 1.5 $U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. View of a molecule of (I) showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are omitted for clarity.

supplementary materials




Fig. 2. Partial packing view of (I). Displacement ellipsoids are drawn at the 30% probability level. Hydrogen bonds are shown as dashed lines.

1-(2,6-Dichlorobenzoyl)-3-(3-nitrophenyl)thiourea dimethylformamide solvate

Crystal data

C ₁₄ H ₉ Cl ₂ N ₃ O ₃ S·C ₃ H ₇ NO	Z = 2
M _r = 443.30	F(000) = 456
Triclinic, P <bar{1}< bar=""></bar{1}<>	D _x = 1.460 Mg m ⁻³
a = 8.507 (5) Å	Mo K α radiation, λ = 0.71073 Å
b = 10.240 (5) Å	Cell parameters from 1797 reflections
c = 12.414 (8) Å	θ = 2.4–25.9°
α = 70.40 (4)°	μ = 0.46 mm ⁻¹
β = 81.74 (5)°	T = 293 K
γ = 87.98 (4)°	Block, yellow
V = 1008 (1) Å ³	0.30 × 0.20 × 0.20 mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer	3413 independent reflections
Radiation source: fine-focus sealed tube graphite	2691 reflections with $I > 2\sigma(I)$
phi and ω scans	$R_{\text{int}} = 0.021$
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1997)	$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 2.4^\circ$
$T_{\text{min}} = 0.875$, $T_{\text{max}} = 0.914$	$h = -10 \rightarrow 9$
4551 measured reflections	$k = -12 \rightarrow 12$
	$l = -14 \rightarrow 13$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.046$	H-atom parameters constrained
$wR(F^2) = 0.139$	$w = 1/[\sigma^2(F_o^2) + (0.0842P)^2 + 0.0267P]$
$S = 1.11$	where $P = (F_o^2 + 2F_c^2)/3$
3413 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
256 parameters	$\Delta\rho_{\text{max}} = 0.30 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.42 \text{ e \AA}^{-3}$
	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001xF_c^2\lambda^3/\sin(2\theta)]^{1/4}$

Primary atom site location: structure-invariant direct methods Extinction coefficient: 0.082 (8)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.05945 (7)	0.84634 (6)	0.96546 (5)	0.0570 (3)
Cl1	0.47677 (12)	0.70890 (8)	0.61611 (6)	0.0911 (3)
Cl2	0.67130 (10)	0.96356 (8)	0.88967 (7)	0.0831 (3)
N1	0.3397 (2)	0.80330 (17)	0.86720 (15)	0.0445 (4)
H1	0.3226	0.8832	0.8187	0.053*
N2	0.2412 (2)	0.62210 (17)	1.03144 (15)	0.0481 (5)
H2	0.3272	0.5811	1.0161	0.058*
N3	-0.0870 (3)	0.6004 (2)	1.39465 (16)	0.0594 (5)
O1	0.52143 (19)	0.63011 (15)	0.90012 (14)	0.0577 (5)
O2	-0.0391 (3)	0.7149 (2)	1.38288 (17)	0.0829 (6)
O3	-0.1862 (2)	0.5355 (2)	1.47485 (16)	0.0832 (6)
C1	0.4823 (2)	0.7462 (2)	0.84468 (18)	0.0433 (5)
C2	0.5907 (3)	0.8398 (2)	0.74414 (18)	0.0466 (5)
C3	0.6004 (3)	0.8284 (2)	0.6353 (2)	0.0588 (6)
C4	0.7002 (4)	0.9109 (3)	0.5417 (2)	0.0786 (9)
H4	0.7032	0.9025	0.4692	0.094*
C5	0.7950 (4)	1.0058 (3)	0.5583 (3)	0.0857 (10)
H5	0.8647	1.0609	0.4962	0.103*
C6	0.7900 (3)	1.0218 (3)	0.6639 (3)	0.0788 (9)
H6	0.8553	1.0868	0.6734	0.095*
C7	0.6859 (3)	0.9394 (2)	0.7564 (2)	0.0569 (6)
C8	0.2176 (3)	0.7497 (2)	0.95824 (18)	0.0426 (5)
C9	0.1355 (2)	0.5494 (2)	1.13216 (18)	0.0444 (5)
C10	0.0778 (3)	0.6112 (2)	1.21291 (18)	0.0477 (5)
H10	0.1048	0.7024	1.2021	0.057*
C11	-0.0212 (3)	0.5334 (2)	1.31033 (18)	0.0496 (5)
C12	-0.0602 (3)	0.3960 (2)	1.3312 (2)	0.0579 (6)
H12	-0.1252	0.3452	1.3981	0.070*
C13	0.0003 (3)	0.3373 (2)	1.2498 (2)	0.0624 (7)
H13	-0.0241	0.2452	1.2617	0.075*
C14	0.0970 (3)	0.4129 (2)	1.1505 (2)	0.0536 (6)

supplementary materials

H14	0.1364	0.3718	1.0958	0.064*
N4	0.6148 (2)	0.7230 (2)	0.23093 (16)	0.0561 (5)
O4	0.6593 (2)	0.91813 (16)	0.26969 (14)	0.0645 (5)
C16	0.6896 (3)	0.8409 (2)	0.2126 (2)	0.0549 (6)
H16	0.7720	0.8675	0.1514	0.066*
C17	0.6543 (4)	0.6418 (3)	0.1547 (2)	0.0737 (8)
H17A	0.7418	0.6852	0.0969	0.110*
H17B	0.5639	0.6365	0.1181	0.110*
H17C	0.6834	0.5500	0.1989	0.110*
C18	0.4832 (4)	0.6777 (3)	0.3233 (3)	0.0856 (9)
H18A	0.4517	0.7523	0.3519	0.128*
H18B	0.5151	0.6003	0.3847	0.128*
H18C	0.3954	0.6503	0.2946	0.128*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0485 (4)	0.0507 (4)	0.0612 (4)	0.0082 (3)	0.0081 (3)	-0.0122 (3)
Cl1	0.1307 (8)	0.0806 (5)	0.0656 (5)	-0.0230 (5)	-0.0006 (4)	-0.0318 (4)
Cl2	0.0883 (6)	0.0810 (5)	0.0918 (6)	-0.0026 (4)	-0.0312 (4)	-0.0362 (4)
N1	0.0443 (10)	0.0372 (8)	0.0438 (10)	0.0025 (7)	0.0030 (8)	-0.0068 (7)
N2	0.0432 (10)	0.0420 (9)	0.0505 (11)	0.0018 (8)	0.0079 (8)	-0.0104 (8)
N3	0.0614 (13)	0.0695 (14)	0.0425 (11)	0.0034 (11)	-0.0032 (10)	-0.0143 (10)
O1	0.0545 (9)	0.0450 (8)	0.0562 (9)	0.0103 (7)	0.0075 (7)	-0.0012 (7)
O2	0.1064 (16)	0.0775 (13)	0.0678 (12)	-0.0047 (12)	0.0060 (11)	-0.0351 (10)
O3	0.0753 (13)	0.1017 (15)	0.0586 (11)	-0.0029 (11)	0.0196 (10)	-0.0198 (11)
C1	0.0445 (12)	0.0407 (11)	0.0425 (11)	0.0002 (9)	-0.0002 (9)	-0.0132 (9)
C2	0.0427 (11)	0.0406 (11)	0.0476 (12)	0.0038 (9)	0.0040 (9)	-0.0078 (9)
C3	0.0645 (15)	0.0500 (13)	0.0525 (14)	0.0020 (11)	0.0107 (11)	-0.0128 (11)
C4	0.090 (2)	0.0636 (16)	0.0582 (16)	0.0079 (16)	0.0252 (15)	-0.0043 (13)
C5	0.073 (2)	0.0604 (17)	0.089 (2)	0.0000 (15)	0.0355 (17)	0.0011 (16)
C6	0.0540 (16)	0.0512 (14)	0.113 (3)	-0.0089 (12)	0.0091 (16)	-0.0105 (15)
C7	0.0453 (13)	0.0490 (12)	0.0693 (16)	-0.0002 (10)	-0.0003 (11)	-0.0136 (11)
C8	0.0456 (12)	0.0386 (10)	0.0419 (11)	-0.0041 (9)	0.0010 (9)	-0.0139 (9)
C9	0.0387 (11)	0.0455 (11)	0.0425 (11)	-0.0026 (9)	0.0003 (9)	-0.0082 (9)
C10	0.0470 (12)	0.0460 (11)	0.0469 (12)	-0.0045 (9)	-0.0015 (10)	-0.0126 (10)
C11	0.0458 (12)	0.0577 (13)	0.0407 (12)	-0.0010 (10)	-0.0015 (10)	-0.0119 (10)
C12	0.0570 (14)	0.0557 (13)	0.0482 (13)	-0.0066 (11)	0.0026 (11)	-0.0036 (11)
C13	0.0654 (16)	0.0449 (12)	0.0651 (15)	-0.0113 (11)	0.0052 (13)	-0.0076 (11)
C14	0.0552 (14)	0.0456 (12)	0.0558 (14)	-0.0004 (10)	0.0015 (11)	-0.0153 (10)
N4	0.0623 (12)	0.0489 (11)	0.0509 (11)	-0.0033 (9)	-0.0015 (9)	-0.0110 (9)
O4	0.0862 (13)	0.0446 (8)	0.0560 (10)	0.0051 (8)	-0.0006 (9)	-0.0123 (8)
C16	0.0620 (15)	0.0470 (12)	0.0465 (13)	0.0028 (11)	-0.0039 (11)	-0.0052 (10)
C17	0.094 (2)	0.0662 (16)	0.0650 (17)	-0.0082 (15)	-0.0123 (15)	-0.0254 (13)
C18	0.077 (2)	0.0725 (18)	0.096 (2)	-0.0153 (15)	0.0155 (17)	-0.0218 (16)

Geometric parameters (\AA , $^\circ$)

S1—C8	1.652 (2)	C6—H6	0.9300
-------	-----------	-------	--------

Cl1—C3	1.738 (3)	C9—C10	1.381 (3)
Cl2—C7	1.740 (3)	C9—C14	1.383 (3)
N1—C1	1.362 (3)	C10—C11	1.384 (3)
N1—C8	1.394 (3)	C10—H10	0.9300
N1—H1	0.8600	C11—C12	1.386 (3)
N2—C8	1.345 (3)	C12—C13	1.373 (4)
N2—C9	1.423 (3)	C12—H12	0.9300
N2—H2	0.8600	C13—C14	1.382 (3)
N3—O2	1.210 (3)	C13—H13	0.9300
N3—O3	1.225 (3)	C14—H14	0.9300
N3—C11	1.472 (3)	N4—C16	1.319 (3)
O1—C1	1.219 (3)	N4—C18	1.448 (4)
C1—C2	1.502 (3)	N4—C17	1.455 (3)
C2—C7	1.384 (3)	O4—C16	1.226 (3)
C2—C3	1.385 (3)	C16—H16	0.9300
C3—C4	1.376 (4)	C17—H17A	0.9600
C4—C5	1.369 (5)	C17—H17B	0.9600
C4—H4	0.9300	C17—H17C	0.9600
C5—C6	1.368 (5)	C18—H18A	0.9600
C5—H5	0.9300	C18—H18B	0.9600
C6—C7	1.387 (4)	C18—H18C	0.9600
C1—N1—C8	128.75 (17)	C14—C9—N2	118.5 (2)
C1—N1—H1	115.6	C9—C10—C11	118.0 (2)
C8—N1—H1	115.6	C9—C10—H10	121.0
C8—N2—C9	125.37 (18)	C11—C10—H10	121.0
C8—N2—H2	117.3	C10—C11—C12	122.7 (2)
C9—N2—H2	117.3	C10—C11—N3	118.1 (2)
O2—N3—O3	123.3 (2)	C12—C11—N3	119.2 (2)
O2—N3—C11	118.9 (2)	C13—C12—C11	117.8 (2)
O3—N3—C11	117.8 (2)	C13—C12—H12	121.1
O1—C1—N1	124.13 (19)	C11—C12—H12	121.1
O1—C1—C2	121.96 (19)	C12—C13—C14	121.0 (2)
N1—C1—C2	113.91 (18)	C12—C13—H13	119.5
C7—C2—C3	117.4 (2)	C14—C13—H13	119.5
C7—C2—C1	121.7 (2)	C13—C14—C9	120.1 (2)
C3—C2—C1	121.0 (2)	C13—C14—H14	119.9
C4—C3—C2	122.5 (3)	C9—C14—H14	119.9
C4—C3—C11	119.1 (2)	C16—N4—C18	119.9 (2)
C2—C3—C11	118.37 (18)	C16—N4—C17	121.3 (2)
C5—C4—C3	118.2 (3)	C18—N4—C17	118.7 (2)
C5—C4—H4	120.9	O4—C16—N4	125.3 (2)
C3—C4—H4	120.9	O4—C16—H16	117.3
C6—C5—C4	121.8 (3)	N4—C16—H16	117.3
C6—C5—H5	119.1	N4—C17—H17A	109.5
C4—C5—H5	119.1	N4—C17—H17B	109.5
C5—C6—C7	119.0 (3)	H17A—C17—H17B	109.5
C5—C6—H6	120.5	N4—C17—H17C	109.5
C7—C6—H6	120.5	H17A—C17—H17C	109.5
C2—C7—C6	121.2 (3)	H17B—C17—H17C	109.5

supplementary materials


C2—C7—Cl2	119.17 (18)	N4—C18—H18A	109.5
C6—C7—Cl2	119.6 (2)	N4—C18—H18B	109.5
N2—C8—N1	115.74 (18)	H18A—C18—H18B	109.5
N2—C8—S1	126.52 (17)	N4—C18—H18C	109.5
N1—C8—S1	117.74 (15)	H18A—C18—H18C	109.5
C10—C9—C14	120.3 (2)	H18B—C18—H18C	109.5
C10—C9—N2	121.05 (19)		
C8—N1—C1—O1	4.9 (4)	C9—N2—C8—S1	-2.9 (3)
C8—N1—C1—C2	-174.8 (2)	C1—N1—C8—N2	-2.8 (3)
O1—C1—C2—C7	-96.1 (3)	C1—N1—C8—S1	177.82 (18)
N1—C1—C2—C7	83.7 (3)	C8—N2—C9—C10	-50.6 (3)
O1—C1—C2—C3	83.2 (3)	C8—N2—C9—C14	132.3 (2)
N1—C1—C2—C3	-97.1 (2)	C14—C9—C10—C11	-1.3 (3)
C7—C2—C3—C4	0.3 (3)	N2—C9—C10—C11	-178.34 (19)
C1—C2—C3—C4	-178.9 (2)	C9—C10—C11—C12	2.0 (3)
C7—C2—C3—Cl1	-177.59 (18)	C9—C10—C11—N3	-178.0 (2)
C1—C2—C3—Cl1	3.2 (3)	O2—N3—C11—C10	-8.1 (3)
C2—C3—C4—C5	1.2 (4)	O3—N3—C11—C10	173.1 (2)
Cl1—C3—C4—C5	179.1 (2)	O2—N3—C11—C12	171.9 (2)
C3—C4—C5—C6	-1.4 (4)	O3—N3—C11—C12	-6.9 (3)
C4—C5—C6—C7	0.0 (4)	C10—C11—C12—C13	-1.4 (4)
C3—C2—C7—C6	-1.7 (3)	N3—C11—C12—C13	178.7 (2)
C1—C2—C7—C6	177.5 (2)	C11—C12—C13—C14	0.1 (4)
C3—C2—C7—Cl2	176.95 (17)	C12—C13—C14—C9	0.6 (4)
C1—C2—C7—Cl2	-3.8 (3)	C10—C9—C14—C13	0.1 (4)
C5—C6—C7—C2	1.6 (4)	N2—C9—C14—C13	177.2 (2)
C5—C6—C7—Cl2	-177.1 (2)	C18—N4—C16—O4	1.4 (4)
C9—N2—C8—N1	177.71 (19)	C17—N4—C16—O4	176.9 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	$D\cdots A$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
N2—H2 \cdots O1 ⁱ	0.86	2.47	3.182 (3)	141.
N2—H2 \cdots O1	0.86	1.99	2.675 (3)	136.
N1—H1 \cdots O4 ⁱⁱ	0.86	1.96	2.787 (3)	161.

Symmetry codes: (i) $-x+1, -y+1, -z+2$; (ii) $-x+1, -y+2, -z+1$.

Fig. 1



supplementary materials

Fig. 2

