

(E)-2-[(4-Fluorophenyl)iminomethyl]-5-methoxyphenol

Ciğdem Albayrak,^{a*} Arzu Özek,^b Başak Koşar,^a Mustafa Odabaşoğlu^c and Orhan Büyükgüngör^b

^aFaculty of Education, Sinop University, Sinop, Turkey, ^bDepartment of Physics, Ondokuz Mayıs University, TR-55139 Samsun, Turkey, and ^cChemistry Programme, Denizli Higher Vocational School, Pamukkale University, TR-20159 Denizli, Turkey
Correspondence e-mail: calbayrak@sinop.edu.tr


Received 4 January 2010; accepted 5 January 2010

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.035; wR factor = 0.104; data-to-parameter ratio = 8.3.

In the molecule of the title compound, $\text{C}_{14}\text{H}_{12}\text{FNO}_2$, the aromatic rings are oriented at a dihedral angle of $48.17(1)^\circ$. An intramolecular $\text{O}-\text{H}\cdots\text{N}$ hydrogen bond results in the formation of a six-membered ring. The title molecule is a phenol-imine tautomer, as evidenced by the $\text{C}-\text{O}$ [1.351 (3) Å], $\text{C}-\text{N}$ [1.282 (3) Å], and $\text{C}-\text{C}$ [1.416 (3)–1.445 (3) Å] bond lengths. In the crystal, molecules are linked by intermolecular $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

The present work is part of a structural study of Schiff bases, see: Özek *et al.* (2007); Odabaşoğlu *et al.* (2007); Albayrak *et al.* (2005). For related structures, see: Özek *et al.* (2007, 2009).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{12}\text{FNO}_2$	$V = 602.49(6)\text{ \AA}^3$
$M_r = 245.25$	$Z = 2$
Monoclinic, Pc	Mo $K\alpha$ radiation
$a = 13.1806(7)\text{ \AA}$	$\mu = 0.10\text{ mm}^{-1}$
$b = 7.1785(5)\text{ \AA}$	$T = 296\text{ K}$
$c = 6.4297(3)\text{ \AA}$	$0.68 \times 0.48 \times 0.17\text{ mm}$
$\beta = 97.967(4)^\circ$	

Data collection

Stoe IPDS II diffractometer	6287 measured reflections
Absorption correction: integration (<i>X-RED32</i> ; Stoe & Cie, 2002)	1399 independent reflections
$T_{\min} = 0.932$, $T_{\max} = 0.985$	1273 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.104$	$\Delta\rho_{\max} = 0.20\text{ e \AA}^{-3}$
$S = 1.09$	$\Delta\rho_{\min} = -0.11\text{ e \AA}^{-3}$
1399 reflections	
168 parameters	
3 restraints	

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg1$ and $Cg2$ are the centroids of $\text{C}1-\text{C}6$ and $\text{C}9-\text{C}14$ rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1 \cdots N1	0.82 (2)	1.87 (2)	2.615 (3)	150 (3)
C6—H6 \cdots Cg1 ⁱ	0.93	2.73	3.4363	133
C11—H11 \cdots Cg2 ⁱⁱ	0.93	2.93	3.6414	134
C14—H14 \cdots Cg2 ⁱⁱⁱ	0.93	2.91	3.6076	133

Symmetry codes: (i) $x, -y, z + \frac{1}{2}$; (ii) $x, -y + 1, z + \frac{1}{2}$; (iii) $x, -y, z - \frac{1}{2}$.

Data collection: *X-AREA* (Stoe & Cie, 2002); cell refinement: *X-AREA*; data reduction: *X-RED32* (Stoe & Cie, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors acknowledge the Faculty of Arts and Sciences, Ondokuz Mayıs University, Turkey, for the use of the Stoe IPDS II diffractometer (purchased under grant F.279 of the University Research Fund).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5163).

References

- Albayrak, Ç., Odabaşoğlu, M. & Büyükgüngör, O. (2005). *Acta Cryst. E61*, o423–o424.
- Farrugia, L. J. (1997). *J. Appl. Cryst. 30*, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst. 32*, 837–838.
- Odabaşoğlu, M., Büyükgüngör, O., Narayana, B., Vijeshi, A. M. & Yathirajan, H. S. (2007). *Acta Cryst. E63*, o1916–o1918.
- Özek, A., Albayrak, Ç. & Büyükgüngör, O. (2009). *Acta Cryst. E65*, o2153.
- Özek, A., Albayrak, Ç., Odabaşoğlu, M. & Büyükgüngör, O. (2007). *Acta Cryst. C63*, o177–o180.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Stoe & Cie (2002). *X-AREA* (Version 1.18) and *X-RED32* (Version 1.04). Stoe & Cie, Darmstadt, Germany.

supplementary materials

Acta Cryst. (2010). E66, o315 [doi:10.1107/S1600536810000474]

(E)-2-[(4-Fluorophenyl)iminomethyl]-5-methoxyphenol

Ç. Albayrak, A. Özek, B. Kosar, M. Odabasoglu and O. Büyükgüngör

Comment

The present work is part of a structural study of Schiff bases (Özek *et al.*, 2009; Özek *et al.*, 2007) and we report here the structure of (E)-2-(4-Fluorophenylimino)methyl-5-methoxyphenol, (I).

The *ortho*-hydroxy Schiff Bases that show tautomerism by the intramolecular proton transfer from an oxygen atom to the neighboring nitrogen atom are important compounds. These compounds can exist in three different structures as enol, keto or zwitterionic forms in the solid state. The title compound (I) consists of two aromatic rings (C1 to C6 and C9 to C14), and an imino frame (C9—N1—C8—C1). In (E)-2-(4-Fluorophenylimino)methyl- 5-methoxyphenol which adopts an E configuration about the C=N double bond, dihedral angle between the aromatic rings is 48.17 (1) °. The H atom in title compound (I) is located on atom O1, thus the phenol-imine tautomer is favored over the keto-amine form, as indicated by the C2—O1, C8—N1, C1—C8 and C1—C2 bond lengths (Fig. 1 and Table 2). The O1—C2 bond length of 1.351 (2) Å indicates single-bond character, whereas the N1—C8 bond length of 1.283 (2) Å indicates double-bond character. A similar work was observed for X-ray crystal and computational structural study of (E)-2-[(4-bromophenyl)iminomethyl]-4-methoxyphenol [C—O=1.358 (4) Å, C—N= 1.287 (4) Å, Özek *et al.*, 2007].

It is known that Schiff bases may exhibit thermochromism or photochromism, depending on the planarity or non-planarity of the molecule, respectively. Therefore, one can expect photochromic properties in (I) caused by non-planarity of the molecules; the dihedral angle between rings A(C1—C6) and B ring (C9—C14) is 48.17 (1) °. The intramolecular O—H···N hydrogen bond (Table 1) results in the formation of six-membered ring and it generates an S(6) ring motif. The O1···N1 distance of 2.614 (2) Å is comparable to those observed for analogous hydrogen bonds in "Three (E)-2-[(bromophenyl)iminomethyl]-4-methoxyphenols" [2.603 (2) Å, 2.638 (7) Å, 2.577 (4) Å; Özek *et al.*, 2007]. In the crystal structure, C—H···π interactions exist (Table 1) (Fig. 2).

Experimental

The compound (E)-2-(4-Fluorophenylimino)methyl-5-methoxyphenol was prepared by reflux a mixture of a solution containing 4-methoxysalicylaldehyde (0.5 g 3.3 mmol) in 20 ml e ethanol and a solution containing 4-fluoroaniline (0.37 g 3.3 mmol) in 20 ml e ethanol. The reaction mixture was stirred for 1 h under reflux. The crystals of (E)-2-(4-Fluorophenylimino)methyl-5-methoxyphenol suitable for X-ray analysis were obtained from ethanol by slow evaporation (yield % 82; m.p. 368–369 K).

Refinement

All H atoms except the hydroxyl H atom (which was freely refined) were refined using riding model with C—H distances of 0.96 Å for the methyl group and 0.93 Å for other H atoms. The displacement parameters of these H atoms were fixed at 1.2 U_{eq} of their parent carbon atom or 1.5 U_{eq} for the methyl group. The absolute structure could not be determined, and 1150 Friedel pairs were averaged before the last refinement.

supplementary materials

Figures




Fig. 1. A view of (I), with the atom-numbering scheme. Dashed line indicates intramolecular hydrogen bond.




Fig. 2. A partial packing diagram for (I), with C—H···Cg bonds shown as dashed lines. Cg1 and Cg2 are the centroids of C1—C6 and C9—C14 rings, respectively. Symmetry codes: (i) $x, -y, z + 1/2$; (ii) $x, -y + 1, z + 1/2$; (iii) $x, -y, z - 1/2$.

(E)-2-[(4-Fluorophenyl)iminomethyl]-5-methoxyphenol

Crystal data

C ₁₄ H ₁₂ FNO ₂	$F(000) = 256$
$M_r = 245.25$	$D_x = 1.352 \text{ Mg m}^{-3}$
Monoclinic, $P\bar{c}$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: P -2yc	Cell parameters from 11108 reflections
$a = 13.1806 (7) \text{ \AA}$	$\theta = 1.6\text{--}28.0^\circ$
$b = 7.1785 (5) \text{ \AA}$	$\mu = 0.10 \text{ mm}^{-1}$
$c = 6.4297 (3) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 97.967 (4)^\circ$	Plate, yellow
$V = 602.49 (6) \text{ \AA}^3$	$0.68 \times 0.48 \times 0.17 \text{ mm}$
$Z = 2$	

Data collection

Stoe IPDS II diffractometer	1399 independent reflections
Radiation source: fine-focus sealed tube	1273 reflections with $I > 2\sigma(I)$
plane graphite	$R_{\text{int}} = 0.037$
Detector resolution: 6.67 pixels mm^{-1}	$\theta_{\text{max}} = 27.6^\circ, \theta_{\text{min}} = 2.8^\circ$
ω -scan rotation method	$h = -17 \rightarrow 17$
Absorption correction: integration (<i>X-RED32</i> ; Stoe & Cie, 2002)	$k = -9 \rightarrow 9$
$T_{\text{min}} = 0.932, T_{\text{max}} = 0.985$	$l = -8 \rightarrow 8$
6287 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.035$	H atoms treated by a mixture of independent and constrained refinement

$wR(F^2) = 0.104$	$w = 1/[\sigma^2(F_o^2) + (0.0672P)^2 + 0.0142P]$
$S = 1.09$	where $P = (F_o^2 + 2F_c^2)/3$
1399 reflections	$(\Delta/\sigma)_{\max} < 0.001$
168 parameters	$\Delta\rho_{\max} = 0.20 \text{ e } \text{\AA}^{-3}$
3 restraints	$\Delta\rho_{\min} = -0.11 \text{ e } \text{\AA}^{-3}$
	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = k F_c [1 + 0.001 x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.022 (7)

Special details

Experimental. 237 frames, detector distance = 100 mm

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.67815 (17)	0.7844 (3)	0.5679 (3)	0.0430 (5)
C2	0.68722 (16)	0.7118 (3)	0.3666 (3)	0.0452 (5)
C3	0.78232 (18)	0.6912 (3)	0.3009 (4)	0.0473 (5)
H3	0.7878	0.6379	0.1711	0.057*
C4	0.86907 (16)	0.7504 (3)	0.4299 (3)	0.0448 (5)
C5	0.86142 (16)	0.8247 (3)	0.6292 (3)	0.0482 (5)
H5	0.9199	0.8648	0.7153	0.058*
C6	0.76785 (16)	0.8378 (3)	0.6958 (3)	0.0472 (5)
H6	0.7637	0.8834	0.8297	0.057*
C7	0.9809 (2)	0.6644 (5)	0.1854 (5)	0.0735 (7)
H7A	0.9579	0.5374	0.1792	0.088*
H7B	1.0525	0.6683	0.1720	0.088*
H7C	0.9429	0.7339	0.0730	0.088*
C8	0.58033 (17)	0.7980 (3)	0.6444 (3)	0.0465 (5)
H8	0.5785	0.8356	0.7822	0.056*
C9	0.40316 (17)	0.7582 (3)	0.6157 (4)	0.0456 (5)
C10	0.39603 (19)	0.6839 (3)	0.8123 (4)	0.0539 (5)
H10	0.4544	0.6382	0.8942	0.065*
C11	0.3028 (2)	0.6775 (4)	0.8871 (4)	0.0606 (6)
H11	0.2976	0.6270	1.0184	0.073*
C12	0.21810 (19)	0.7470 (4)	0.7637 (5)	0.0598 (6)
C13	0.22135 (19)	0.8207 (4)	0.5678 (4)	0.0607 (6)

supplementary materials

H13	0.1625	0.8667	0.4877	0.073*
C14	0.31454 (17)	0.8245 (4)	0.4932 (4)	0.0522 (5)
H14	0.3185	0.8717	0.3599	0.063*
N1	0.49593 (14)	0.7598 (2)	0.5277 (3)	0.0485 (5)
O1	0.60343 (14)	0.6613 (3)	0.2332 (3)	0.0631 (5)
O2	0.96531 (13)	0.7433 (3)	0.3800 (3)	0.0561 (4)
F1	0.12629 (15)	0.7419 (3)	0.8381 (4)	0.0922 (6)
H1	0.553 (2)	0.676 (4)	0.294 (5)	0.079 (10)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0418 (10)	0.0416 (11)	0.0453 (12)	-0.0015 (8)	0.0053 (8)	-0.0006 (8)
C2	0.0420 (11)	0.0492 (11)	0.0433 (12)	-0.0009 (8)	0.0020 (9)	-0.0016 (8)
C3	0.0479 (11)	0.0539 (12)	0.0406 (10)	-0.0011 (9)	0.0075 (8)	-0.0044 (9)
C4	0.0433 (11)	0.0459 (10)	0.0459 (12)	0.0014 (8)	0.0083 (9)	0.0045 (9)
C5	0.0439 (11)	0.0547 (11)	0.0443 (11)	-0.0041 (8)	-0.0003 (9)	-0.0031 (9)
C6	0.0493 (12)	0.0510 (11)	0.0403 (11)	-0.0007 (8)	0.0034 (9)	-0.0047 (8)
C7	0.0540 (14)	0.109 (2)	0.0612 (15)	-0.0001 (14)	0.0216 (11)	-0.0148 (14)
C8	0.0456 (11)	0.0472 (11)	0.0467 (11)	0.0012 (8)	0.0068 (9)	-0.0022 (8)
C9	0.0427 (11)	0.0453 (11)	0.0492 (12)	-0.0012 (8)	0.0073 (9)	-0.0028 (8)
C10	0.0521 (12)	0.0561 (12)	0.0527 (13)	0.0052 (10)	0.0051 (10)	0.0028 (10)
C11	0.0668 (16)	0.0630 (13)	0.0541 (13)	-0.0032 (12)	0.0158 (12)	0.0027 (11)
C12	0.0453 (13)	0.0678 (14)	0.0692 (17)	-0.0077 (10)	0.0184 (12)	-0.0090 (12)
C13	0.0435 (12)	0.0705 (15)	0.0663 (17)	-0.0007 (11)	0.0008 (11)	-0.0035 (12)
C14	0.0461 (12)	0.0593 (12)	0.0503 (13)	-0.0003 (9)	0.0041 (9)	0.0013 (10)
N1	0.0415 (10)	0.0532 (10)	0.0509 (11)	0.0016 (8)	0.0063 (8)	-0.0004 (8)
O1	0.0438 (8)	0.0922 (12)	0.0515 (9)	-0.0082 (8)	0.0007 (7)	-0.0194 (9)
O2	0.0420 (8)	0.0746 (12)	0.0529 (9)	-0.0021 (7)	0.0108 (7)	-0.0033 (8)
F1	0.0568 (10)	0.1257 (16)	0.1005 (15)	-0.0092 (10)	0.0334 (9)	-0.0025 (11)

Geometric parameters (\AA , °)

C1—C6	1.397 (3)	C8—N1	1.282 (3)
C1—C2	1.416 (3)	C8—H8	0.9300
C1—C8	1.445 (3)	C9—C10	1.387 (3)
C2—O1	1.351 (3)	C9—C14	1.399 (3)
C2—C3	1.385 (3)	C9—N1	1.417 (3)
C3—C4	1.383 (3)	C10—C11	1.381 (3)
C3—H3	0.9300	C10—H10	0.9300
C4—O2	1.352 (3)	C11—C12	1.371 (4)
C4—C5	1.404 (3)	C11—H11	0.9300
C5—C6	1.364 (3)	C12—F1	1.362 (3)
C5—H5	0.9300	C12—C13	1.372 (4)
C6—H6	0.9300	C13—C14	1.379 (3)
C7—O2	1.414 (3)	C13—H13	0.9300
C7—H7A	0.9600	C14—H14	0.9300
C7—H7B	0.9600	O1—H1	0.824 (19)
C7—H7C	0.9600		

C6—C1—C2	117.86 (19)	N1—C8—C1	121.93 (19)
C6—C1—C8	120.22 (19)	N1—C8—H8	119.0
C2—C1—C8	121.89 (18)	C1—C8—H8	119.0
O1—C2—C3	118.2 (2)	C10—C9—C14	119.1 (2)
O1—C2—C1	120.96 (19)	C10—C9—N1	122.6 (2)
C3—C2—C1	120.86 (18)	C14—C9—N1	118.2 (2)
C4—C3—C2	119.5 (2)	C11—C10—C9	120.4 (2)
C4—C3—H3	120.3	C11—C10—H10	119.8
C2—C3—H3	120.3	C9—C10—H10	119.8
O2—C4—C3	124.8 (2)	C12—C11—C10	118.6 (3)
O2—C4—C5	114.78 (19)	C12—C11—H11	120.7
C3—C4—C5	120.4 (2)	C10—C11—H11	120.7
C6—C5—C4	119.69 (19)	F1—C12—C11	118.6 (3)
C6—C5—H5	120.2	F1—C12—C13	118.4 (3)
C4—C5—H5	120.2	C11—C12—C13	123.0 (2)
C5—C6—C1	121.6 (2)	C12—C13—C14	118.0 (2)
C5—C6—H6	119.2	C12—C13—H13	121.0
C1—C6—H6	119.2	C14—C13—H13	121.0
O2—C7—H7A	109.5	C13—C14—C9	120.8 (2)
O2—C7—H7B	109.5	C13—C14—H14	119.6
H7A—C7—H7B	109.5	C9—C14—H14	119.6
O2—C7—H7C	109.5	C8—N1—C9	119.66 (17)
H7A—C7—H7C	109.5	C2—O1—H1	108 (3)
H7B—C7—H7C	109.5	C4—O2—C7	118.77 (19)
C6—C1—C2—O1	-178.6 (2)	C14—C9—C10—C11	-0.7 (3)
C8—C1—C2—O1	3.5 (3)	N1—C9—C10—C11	-176.9 (2)
C6—C1—C2—C3	1.3 (3)	C9—C10—C11—C12	-0.4 (4)
C8—C1—C2—C3	-176.6 (2)	C10—C11—C12—F1	-179.5 (2)
O1—C2—C3—C4	176.8 (2)	C10—C11—C12—C13	0.9 (4)
C1—C2—C3—C4	-3.1 (3)	F1—C12—C13—C14	-179.7 (2)
C2—C3—C4—O2	-177.68 (19)	C11—C12—C13—C14	-0.1 (4)
C2—C3—C4—C5	2.3 (3)	C12—C13—C14—C9	-1.1 (4)
O2—C4—C5—C6	-179.79 (19)	C10—C9—C14—C13	1.5 (3)
C3—C4—C5—C6	0.2 (3)	N1—C9—C14—C13	177.9 (2)
C4—C5—C6—C1	-2.1 (3)	C1—C8—N1—C9	174.01 (17)
C2—C1—C6—C5	1.3 (3)	C10—C9—N1—C8	-40.8 (3)
C8—C1—C6—C5	179.26 (19)	C14—C9—N1—C8	143.0 (2)
C6—C1—C8—N1	176.2 (2)	C3—C4—O2—C7	-2.2 (3)
C2—C1—C8—N1	-5.9 (3)	C5—C4—O2—C7	177.8 (2)

Hydrogen-bond geometry (Å, °)

Cg1 and Cg2 are the centroids of C1—C6 and C9—C14 rings, respectively.

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1···N1	0.82 (2)	1.87 (2)	2.615 (3)	150 (3)
C6—H6···Cg1 ⁱ	0.93	2.73	3.4363	133.
C11—H11···Cg2 ⁱⁱ	0.93	2.93	3.6414	134.
C14—H14···Cg2 ⁱⁱⁱ	0.93	2.91	3.6076	133.

supplementary materials

Symmetry codes: (i) $x, -y, z+1/2$; (ii) $x, -y+1, z+1/2$; (iii) $x, -y, z-1/2$.

Fig. 1




Fig. 2

