

## 2-Chloro-N'-(5-hydroxy-2-nitrobenzylidene)benzohydrazide

Cong-Shan Zhou and Tao Yang\*

College of Chemistry and Chemical Engineering, Hunan Institute of Science and Technology, Yueyang, Hunan 414006, People's Republic of China  
Correspondence e-mail: zhoucongsh@gmail.com


Received 10 January 2010; accepted 11 January 2010

Key indicators: single-crystal X-ray study;  $T = 298\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ;  $R$  factor = 0.037;  $wR$  factor = 0.103; data-to-parameter ratio = 14.3.

The molecule of the title Schiff base compound,  $\text{C}_{14}\text{H}_{10}\text{ClN}_3\text{O}_4$ , exists in a *trans* configuration with respect to the acyclic  $\text{C}=\text{N}$  bond. The dihedral angle between the two benzene rings is  $62.37(9)^\circ$ . An intramolecular  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bond is observed. In the crystal structure, adjacent molecules are linked into a ribbon along  $[1\bar{1}0]$  by  $\text{O}-\text{H}\cdots\text{O}$  and  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds.

### Related literature

For the biological properties of Schiff bases, see: Mohamed *et al.* (2009); Ritter *et al.* (2009); Bagihalli *et al.* (2008). For related structures, see: Fun *et al.* (2008); Shafiq *et al.* (2009); Goh *et al.* (2010); Zhou *et al.* (2009); Zhou & Yang (2009).



### Experimental

#### Crystal data

$\text{C}_{14}\text{H}_{10}\text{ClN}_3\text{O}_4$	$\gamma = 96.127(2)^\circ$
$M_r = 319.70$	$V = 694.64(4)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 7.2490(2)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 9.4719(3)\text{ \AA}$	$\mu = 0.30\text{ mm}^{-1}$
$c = 10.4749(4)\text{ \AA}$	$T = 298\text{ K}$
$\alpha = 100.623(2)^\circ$	$0.17 \times 0.15 \times 0.15\text{ mm}$
$\beta = 97.433(2)^\circ$	

#### Data collection

Bruker SMART 1000 CCD area-detector diffractometer  
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.951$ ,  $T_{\max} = 0.957$

4097 measured reflections  
2900 independent reflections  
2332 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.016$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$   
 $wR(F^2) = 0.103$   
 $S = 1.03$   
2900 reflections  
203 parameters  
1 restraint

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.20\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.25\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}2-\text{H}2\cdots\text{O}2^i$	0.89 (1)	2.09 (1)	2.9591 (18)	165 (2)
$\text{O}4-\text{H}4\cdots\text{O}1^{ii}$	0.82	1.85	2.6708 (17)	176
$\text{C}7-\text{H}7\cdots\text{O}2$	0.93	2.22	2.817 (2)	122

Symmetry codes: (i)  $-x + 1, -y + 1, -z + 1$ ; (ii)  $-x, -y + 2, -z + 1$ .

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

The authors acknowledge the Hunan Provincial Natural Science Foundation of China (grant No. 09J6022) and the Scientific Research Fund of Hunan Provincial Education Department (grant No. 08B031), China, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI5018).

### References

- Bagihalli, G. B., Avaji, P. G., Patil, S. A. & Badami, P. S. (2008). *Eur. J. Med. Chem.* **43**, 2639–2649.
- Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Fun, H.-K., Chantrapromma, S., Jana, S., Hazra, A. & Goswami, S. (2008). *Acta Cryst. E* **64**, o175–o176.
- Goh, J. H., Fun, H.-K., Vinayaka, A. C. & Kalluraya, B. (2010). *Acta Cryst. E* **66**, o24.
- Mohamed, G. G., Omar, M. M. & Ibrahim, A. A. (2009). *Eur. J. Med. Chem.* **44**, 4801–4812.
- Ritter, E., Przybylski, P., Brzezinski, B. & Bartl, F. (2009). *Curr. Org. Chem.* **13**, 241–249.
- Shafiq, Z., Yaqub, M., Tahir, M. N., Hussain, A. & Iqbal, M. S. (2009). *Acta Cryst. E* **65**, o2501.
- Sheldrick, G. M. (1996). *SADABS*. University of Goettingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Zhou, C.-S., Hou, H.-Y. & Yang, T. (2009). *Z. Kristallogr. New Cryst. Struct.* **224**, 37–38.
- Zhou, C.-S. & Yang, T. (2009). *Z. Kristallogr. New Cryst. Struct.* **224**, 39–40.

## **supplementary materials**

Acta Cryst. (2010). E66, o365 [doi:10.1107/S1600536810001303]

## 2-Chloro-N'-(5-hydroxy-2-nitrobenzylidene)benzohydrazide

C.-S. Zhou and T. Yang

### Comment

Schiff bases usually possess excellent biological properties, such as antibacterial, antimicrobial, and antitumor (Mohamed *et al.*, 2009; Ritter *et al.*, 2009; Bagihalli *et al.*, 2008). Recently, a large number of Schiff bases derived from the reaction of aldehydes with benzohydrazides have been reported (Fun *et al.*, 2008; Shafiq *et al.*, 2009; Goh *et al.*, 2010). In this paper, the crystal structure of the title new Schiff base derived from the condensing of 5-hydroxy-2-nitrobenzaldehyde with 2-chlorobenzohydrazide in methanol is reported.

Bond lengths in the title molecule (Fig. 1) are comparable to those observed in related structures (Zhou *et al.*, 2009; Zhou & Yang, 2009). The molecule exists in a *trans* configuration with respect to the acyclic C=N bond. The dihedral angle between the two benzene rings is 62.37 (9) $^{\circ}$ . An intramolecular C—H $\cdots$ O hydrogen bond is observed.

In the crystal structure, intermolecular N—H $\cdots$ O and O—H $\cdots$ O hydrogen bonds link adjacent molecules into a ribbon along [1 $\bar{1}$ 0] (Table 1 and Fig. 2).

### Experimental

5-Hydroxy-2-nitrobenzaldehyde (1.0 mmol, 167.1 mg) and 2-chlorobenzohydrazide (1.0 mmol, 170.0 mg) were dissolved in a methanol solution (30 ml). The mixture was stirred for 30 min at room temperature. The resulting solution was left in air for a few days, yielding colourless block-shaped crystals.

### Refinement

Atom H2 was located in a difference map and refined with the N—H distance restrained to 0.90 (1) Å. The remaining H atoms were positioned geometrically [C—H = 0.93 Å and O—H = 0.82 Å] and refined using a riding model, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  and  $1.5U_{\text{eq}}(\text{O})$ .

### Figures




Fig. 1. The molecular structure of the title compound, with displacement ellipsoids drawn at the 30% probability level.

## supplementary materials

---




Fig. 2. Part of the crystal packing of the title compound, viewed along the  $a$  axis. Hydrogen bonds are shown as dashed lines.

### 2-Chloro-N<sup>1</sup>-(5-hydroxy-2-nitrobenzylidene)benzohydrazide

#### Crystal data

C <sub>14</sub> H <sub>10</sub> ClN <sub>3</sub> O <sub>4</sub>	Z = 2
$M_r = 319.70$	$F(000) = 328$
Triclinic, $P\bar{1}$	$D_x = 1.528 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 7.2490 (2) \text{ \AA}$	Cell parameters from 1937 reflections
$b = 9.4719 (3) \text{ \AA}$	$\theta = 2.6\text{--}28.4^\circ$
$c = 10.4749 (4) \text{ \AA}$	$\mu = 0.30 \text{ mm}^{-1}$
$\alpha = 100.623 (2)^\circ$	$T = 298 \text{ K}$
$\beta = 97.433 (2)^\circ$	Block, colourless
$\gamma = 96.127 (2)^\circ$	$0.17 \times 0.15 \times 0.15 \text{ mm}$
$V = 694.64 (4) \text{ \AA}^3$	

#### Data collection

Bruker SMART 1000 CCD area-detector diffractometer	2900 independent reflections
Radiation source: fine-focus sealed tube graphite	2332 reflections with $I > 2\sigma(I)$
$\omega$ scans	$R_{\text{int}} = 0.016$
Absorption correction: multi-scan ( <i>SADABS</i> ; Sheldrick, 1996)	$\theta_{\text{max}} = 27.0^\circ, \theta_{\text{min}} = 2.0^\circ$
$T_{\text{min}} = 0.951, T_{\text{max}} = 0.957$	$h = -8 \rightarrow 9$
4097 measured reflections	$k = -11 \rightarrow 12$
	$l = -11 \rightarrow 13$

#### Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.037$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.103$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.03$	$w = 1/[\sigma^2(F_o^2) + (0.0433P)^2 + 0.2231P]$
2900 reflections	where $P = (F_o^2 + 2F_c^2)/3$
203 parameters	$(\Delta/\sigma)_{\max} = 0.001$
1 restraint	$\Delta\rho_{\max} = 0.20 \text{ e \AA}^{-3}$
	$\Delta\rho_{\min} = -0.25 \text{ e \AA}^{-3}$

### Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor wR and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\text{sigma}(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.21511 (7)	0.78685 (7)	0.05703 (5)	0.06308 (19)
N1	0.2273 (2)	0.77176 (15)	0.47983 (14)	0.0378 (3)
N2	0.3525 (2)	0.78387 (15)	0.39232 (15)	0.0387 (3)
N3	0.3042 (2)	0.46714 (16)	0.72546 (17)	0.0469 (4)
O1	0.30056 (19)	1.01035 (13)	0.37374 (13)	0.0487 (3)
O2	0.3739 (2)	0.42683 (16)	0.62639 (16)	0.0618 (4)
O3	0.3681 (2)	0.44819 (19)	0.83292 (17)	0.0740 (5)
O4	-0.33418 (18)	0.74516 (14)	0.71890 (14)	0.0501 (3)
H4	-0.3206	0.8188	0.6885	0.075*
C1	0.1112 (2)	0.63648 (17)	0.62866 (16)	0.0356 (4)
C2	0.1367 (2)	0.54064 (17)	0.71438 (17)	0.0370 (4)
C3	0.0078 (3)	0.51237 (18)	0.79603 (18)	0.0427 (4)
H3	0.0290	0.4480	0.8518	0.051*
C4	-0.1514 (3)	0.57918 (19)	0.79491 (18)	0.0418 (4)
H4A	-0.2407	0.5575	0.8471	0.050*
C5	-0.1775 (2)	0.67960 (18)	0.71488 (17)	0.0381 (4)
C6	-0.0480 (2)	0.70698 (18)	0.63260 (17)	0.0372 (4)
H6	-0.0679	0.7737	0.5790	0.045*
C7	0.2412 (2)	0.66589 (18)	0.53720 (18)	0.0396 (4)
H7	0.3336	0.6068	0.5214	0.047*
C8	0.3786 (2)	0.90324 (17)	0.34225 (16)	0.0342 (4)
C9	0.5138 (2)	0.89383 (17)	0.24509 (17)	0.0352 (4)
C10	0.4520 (2)	0.84446 (18)	0.11239 (18)	0.0391 (4)

## supplementary materials

---

C11	0.5744 (3)	0.8381 (2)	0.02143 (19)	0.0480 (5)
H11	0.5302	0.8060	-0.0677	0.058*
C12	0.7634 (3)	0.8801 (2)	0.0653 (2)	0.0551 (5)
H12	0.8473	0.8770	0.0051	0.066*
C13	0.8291 (3)	0.9268 (2)	0.1973 (2)	0.0598 (6)
H13	0.9571	0.9537	0.2260	0.072*
C14	0.7048 (3)	0.9337 (2)	0.2875 (2)	0.0497 (5)
H14	0.7495	0.9651	0.3767	0.060*
H2	0.421 (3)	0.713 (2)	0.373 (2)	0.080*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl1	0.0402 (3)	0.0935 (4)	0.0521 (3)	0.0016 (2)	0.0072 (2)	0.0100 (3)
N1	0.0386 (8)	0.0387 (7)	0.0418 (8)	0.0101 (6)	0.0200 (6)	0.0102 (6)
N2	0.0396 (8)	0.0373 (7)	0.0475 (8)	0.0143 (6)	0.0243 (7)	0.0128 (6)
N3	0.0455 (9)	0.0406 (8)	0.0619 (11)	0.0148 (7)	0.0141 (8)	0.0196 (7)
O1	0.0609 (8)	0.0432 (7)	0.0561 (8)	0.0249 (6)	0.0327 (7)	0.0191 (6)
O2	0.0643 (10)	0.0625 (9)	0.0756 (10)	0.0356 (7)	0.0338 (8)	0.0261 (8)
O3	0.0729 (11)	0.0925 (12)	0.0715 (11)	0.0404 (9)	0.0110 (9)	0.0382 (9)
O4	0.0463 (7)	0.0567 (8)	0.0628 (9)	0.0224 (6)	0.0302 (6)	0.0280 (7)
C1	0.0368 (9)	0.0324 (8)	0.0399 (9)	0.0056 (6)	0.0127 (7)	0.0077 (7)
C2	0.0363 (9)	0.0340 (8)	0.0438 (9)	0.0088 (7)	0.0106 (7)	0.0103 (7)
C3	0.0490 (10)	0.0395 (9)	0.0454 (10)	0.0083 (8)	0.0141 (8)	0.0176 (8)
C4	0.0428 (10)	0.0447 (9)	0.0441 (10)	0.0073 (7)	0.0198 (8)	0.0149 (8)
C5	0.0382 (9)	0.0376 (8)	0.0419 (9)	0.0089 (7)	0.0143 (7)	0.0086 (7)
C6	0.0394 (9)	0.0371 (8)	0.0411 (9)	0.0100 (7)	0.0154 (7)	0.0138 (7)
C7	0.0382 (9)	0.0384 (9)	0.0489 (10)	0.0135 (7)	0.0182 (8)	0.0134 (8)
C8	0.0334 (8)	0.0372 (8)	0.0349 (9)	0.0093 (7)	0.0112 (7)	0.0078 (7)
C9	0.0366 (9)	0.0319 (8)	0.0410 (9)	0.0083 (6)	0.0163 (7)	0.0079 (7)
C10	0.0363 (9)	0.0426 (9)	0.0426 (10)	0.0083 (7)	0.0141 (7)	0.0124 (7)
C11	0.0512 (11)	0.0585 (11)	0.0399 (10)	0.0129 (9)	0.0199 (9)	0.0123 (8)
C12	0.0499 (12)	0.0622 (12)	0.0601 (13)	0.0098 (9)	0.0338 (10)	0.0110 (10)
C13	0.0373 (10)	0.0655 (13)	0.0723 (15)	-0.0002 (9)	0.0225 (10)	-0.0031 (11)
C14	0.0407 (10)	0.0555 (11)	0.0484 (11)	0.0057 (8)	0.0121 (8)	-0.0047 (9)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Cl1—C10	1.7346 (18)	C4—C5	1.391 (2)
N1—C7	1.267 (2)	C4—H4A	0.93
N1—N2	1.3814 (18)	C5—C6	1.390 (2)
N2—C8	1.337 (2)	C6—H6	0.93
N2—H2	0.891 (10)	C7—H7	0.93
N3—O3	1.213 (2)	C8—C9	1.500 (2)
N3—O2	1.228 (2)	C9—C10	1.381 (2)
N3—C2	1.465 (2)	C9—C14	1.387 (3)
O1—C8	1.2268 (19)	C10—C11	1.381 (2)
O4—C5	1.353 (2)	C11—C12	1.378 (3)
O4—H4	0.82	C11—H11	0.93

C1—C6	1.395 (2)	C12—C13	1.376 (3)
C1—C2	1.399 (2)	C12—H12	0.93
C1—C7	1.470 (2)	C13—C14	1.385 (3)
C2—C3	1.384 (2)	C13—H13	0.93
C3—C4	1.374 (2)	C14—H14	0.93
C3—H3	0.93		
C7—N1—N2	114.29 (13)	C1—C6—H6	119.3
C8—N2—N1	121.18 (13)	N1—C7—C1	120.03 (14)
C8—N2—H2	119.8 (16)	N1—C7—H7	120.0
N1—N2—H2	119.0 (16)	C1—C7—H7	120.0
O3—N3—O2	122.61 (16)	O1—C8—N2	123.19 (14)
O3—N3—C2	118.25 (16)	O1—C8—C9	123.39 (14)
O2—N3—C2	119.14 (16)	N2—C8—C9	113.42 (13)
C5—O4—H4	109.5	C10—C9—C14	118.64 (15)
C6—C1—C2	116.75 (14)	C10—C9—C8	121.08 (15)
C6—C1—C7	119.39 (14)	C14—C9—C8	120.28 (16)
C2—C1—C7	123.86 (15)	C9—C10—C11	121.74 (17)
C3—C2—C1	122.08 (15)	C9—C10—Cl1	119.70 (12)
C3—C2—N3	115.99 (15)	C11—C10—Cl1	118.56 (15)
C1—C2—N3	121.91 (15)	C12—C11—C10	118.78 (18)
C4—C3—C2	120.15 (15)	C12—C11—H11	120.6
C4—C3—H3	119.9	C10—C11—H11	120.6
C2—C3—H3	119.9	C13—C12—C11	120.62 (17)
C3—C4—C5	119.29 (15)	C13—C12—H12	119.7
C3—C4—H4A	120.4	C11—C12—H12	119.7
C5—C4—H4A	120.4	C12—C13—C14	120.08 (19)
O4—C5—C6	122.39 (15)	C12—C13—H13	120.0
O4—C5—C4	117.36 (15)	C14—C13—H13	120.0
C6—C5—C4	120.23 (15)	C13—C14—C9	120.11 (19)
C5—C6—C1	121.40 (15)	C13—C14—H14	119.9
C5—C6—H6	119.3	C9—C14—H14	119.9

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N2—H2···O2 <sup>i</sup>	0.89 (1)	2.09 (1)	2.9591 (18)	165 (2)
O4—H4···O1 <sup>ii</sup>	0.82	1.85	2.6708 (17)	176
C7—H7···O2	0.93	2.22	2.817 (2)	122

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $-x, -y+2, -z+1$ .

## supplementary materials

---

Fig. 1




Fig. 2

