

2-Carboxy-1-(3-nitrophenyl)-ethanaminium perchlorate

Wen-Xian Liang

Ordered Matter Science Research Center, College of Chemistry and Chemical Engineering, Southeast University, Nanjing 210096, People's Republic of China
Correspondence e-mail: lwx927lh@163.com


Received 28 November 2009; accepted 4 December 2009

Key indicators: single-crystal X-ray study; $T = 93\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.030; wR factor = 0.083; data-to-parameter ratio = 15.0.

In the title compound, $\text{C}_9\text{H}_{11}\text{N}_2\text{O}_4^+\cdot\text{ClO}_4^-$, the organic cations form centrosymmetric dimers *via* a pair of $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds between the carboxyl groups. In the crystal, $\text{N}-\text{H}\cdots\text{O}$ interactions between the protonated amine group and the perchlorate anions and the nitro group connect the components into a two-dimensional network parallel to (001).

Related literature

For methods of preparation of β -amino acids, see: Cohen *et al.* (2002); Qu *et al.* (2004).



Experimental

Crystal data

$\text{C}_9\text{H}_{11}\text{N}_2\text{O}_4^+\cdot\text{ClO}_4^-$	$c = 11.8234(11)\text{ \AA}$
$M_r = 310.65$	$\alpha = 94.973(5)^\circ$
Triclinic, $\bar{P}1$	$\beta = 99.093(4)^\circ$
$a = 7.4737(10)\text{ \AA}$	$\gamma = 115.574(9)^\circ$
$b = 7.7676(8)\text{ \AA}$	$V = 602.04(12)\text{ \AA}^3$

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.36\text{ mm}^{-1}$

$T = 93\text{ K}$
 $0.45 \times 0.30 \times 0.15\text{ mm}$

Data collection

Rigaku SCXmini diffractometer
Absorption correction: multi-scan
(*CrystalClear*; Rigaku, 2005)
 $T_{\min} = 0.882$, $T_{\max} = 0.950$

6395 measured reflections
2737 independent reflections
1732 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.049$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.083$
 $S = 1.08$
2737 reflections

183 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.41\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.38\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O3-H3 \cdots O4 ⁱ	0.82	1.82	2.6381 (14)	172
N2-H2D \cdots O5 ⁱⁱ	0.89	2.30	3.0542 (17)	142
N2-H2D \cdots O1 ⁱⁱⁱ	0.89	2.26	2.9200 (17)	130
N2-H2C \cdots O8 ^{iv}	0.89	2.06	2.9159 (17)	162
N2-H2C \cdots O4	0.89	2.45	2.9317 (16)	114
N2-H2B \cdots O7	0.89	2.57	3.1795 (18)	126
N2-H2B \cdots O6	0.89	2.09	2.9720 (17)	172

Symmetry codes: (i) $-x + 2, -y, -z$; (ii) $x + 1, y, z$; (iii) $x + 1, y + 1, z$; (iv) $-x + 1, -y + 1, -z$.

Data collection: *CrystalClear* (Rigaku 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *PRPKAPPA* (Ferguson, 1999).

This work was supported by the Technical Fund Financing Projects (No. 9207042464 and 9207041482) from Southeast University to Zhi-Rong Qu.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2243).

References

- Cohen, J. H., Abdel-Magid, A. F., Almond, H. R. Jr & Maryanoff, C. A. (2002). *Tetrahedron Lett.* **43**, 1977–1981.
- Ferguson, G. (1999). *PRPKAPPA*. University of Guelph, Canada.
- Qu, Z.-R., Zhao, H., Wang, Y.-P., Wang, X.-S., Ye, Q., Li, Y.-H., Xiong, R.-G., Abrahams, B. F., Liu, Z.-G. & Xue, Z.-L. (2004). *Chem. Eur. J.* **10**, 54–60.
- Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2010). E66, o389 [doi:10.1107/S1600536809052167]

2-Carboxy-1-(3-nitrophenyl)ethanaminium perchlorate

W.-X. Liang

Comment

β -Amino acids are important molecules due to their pharmacological properties. Recently, there have been an increased interest in the enantiomeric preparation of β -amino acids as precursors for the synthesis of novel biologically active compounds (Cohen *et al.*, 2002). In addition, β -amino acids are attractive ligands for use in the generation of polar coordination polymers, especially when one considers the ferroelectric compounds (Qu *et al.*, 2004).

The molecular structure of the title compound $C_9H_{11}N_2O_4^+ \cdot ClO_4^-$ is shown in Fig. 1. The crystal is stabilized by intermolecular hydrogen bonds of N—H \cdots O, C—H \cdots O, O—H \cdots O type (Table 1) that connect neighbouring cations and anions, resulting in a two-dimensional network shown in Fig. 2).

Experimental

Under nitrogen protection, 3-nitrobenzaldehyde (4.53 g, 30 mmol), malonic acid (5.0 g, 48 mmol) and ammonium acetate (6.0 g, 78 mmol) were added to a flask and refluxed for 12 h yielding a white precipitate. After being cooled to room temperature, the solution was filtered and the 3-amino-3-(3-nitrophenyl)propanoic acid was obtained, it was dissolved in ethanol and perchloric acid. After slowly evaporating over a period of 3 d, colorless prism crystals of the title compound suitable for diffraction studies were isolated.

Refinement

Positional parameters of all H atoms were calculated geometrically and H atoms were allowed to ride on their parent atoms, with C—H = 0.93 to 0.97 Å, N—H = 0.89 Å, O—H = 0.82 Å, and with $U_{iso}(H) = 1.2 U_{eq}(C)$ or $1.5 U_{eq}(N,O)$.

Figures




Fig. 1. Cation and anion in the title compound with the displacement ellipsoids drawn at the 30% probability level. Intramolecular hydrogen bonds are shown as dashed lines.

supplementary materials




Fig. 2. Packing diagram of the title compound showing hydrogen bonding interactions. Hydrogen bonds are shown as dashed lines.

2-Carboxy-1-(3-nitrophenyl)ethanaminium perchlorate

Crystal data

$\text{C}_9\text{H}_{11}\text{N}_2\text{O}_4^+\cdot\text{ClO}_4^-$	$Z = 2$
$M_r = 310.65$	$F(000) = 320$
Triclinic, $P\bar{1}$	$D_x = 1.713 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 7.4737 (10) \text{ \AA}$	Cell parameters from 1721 reflections
$b = 7.7676 (8) \text{ \AA}$	$\theta = 3.1\text{--}27.5^\circ$
$c = 11.8234 (11) \text{ \AA}$	$\mu = 0.36 \text{ mm}^{-1}$
$\alpha = 94.973 (5)^\circ$	$T = 93 \text{ K}$
$\beta = 99.093 (4)^\circ$	Prism, colorless
$\gamma = 115.574 (9)^\circ$	$0.45 \times 0.30 \times 0.15 \text{ mm}$
$V = 602.04 (12) \text{ \AA}^3$	

Data collection

Rigaku SCXmini diffractometer	2737 independent reflections
Radiation source: fine-focus sealed tube graphite	1732 reflections with $I > 2\sigma(I)$
Detector resolution: 13.6612 pixels mm^{-1}	$R_{\text{int}} = 0.049$
ω and φ scan	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 2.9^\circ$
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku, 2005)	$h = -9 \rightarrow 9$
$T_{\text{min}} = 0.882, T_{\text{max}} = 0.950$	$k = -10 \rightarrow 10$
6395 measured reflections	$l = -15 \rightarrow 15$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.030$	H-atom parameters constrained
$wR(F^2) = 0.083$	$w = 1/[\sigma^2(F_o^2) + (0.0372P)^2 + 0.395P]$
	where $P = (F_o^2 + 2F_c^2)/3$

$S = 1.08$	$(\Delta/\sigma)_{\text{max}} = 0.001$
2737 reflections	$\Delta\rho_{\text{max}} = 0.41 \text{ e \AA}^{-3}$
183 parameters	$\Delta\rho_{\text{min}} = -0.38 \text{ e \AA}^{-3}$
0 restraints	Extinction correction: <i>SHELXL</i>
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.0014 (2)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.03107 (18)	-0.28766 (17)	0.32513 (10)	0.0261 (3)
O2	0.04653 (17)	-0.16514 (17)	0.49951 (10)	0.0222 (2)
O3	0.98215 (17)	-0.04224 (17)	0.14192 (9)	0.0188 (2)
H3	1.0329	-0.0530	0.0868	0.028*
O4	0.82886 (16)	0.08503 (15)	0.02079 (9)	0.0166 (2)
N1	0.11075 (18)	-0.15634 (18)	0.40993 (11)	0.0163 (3)
N2	0.74545 (19)	0.36296 (18)	0.15908 (10)	0.0155 (2)
H2D	0.8712	0.4242	0.2025	0.023*
H2B	0.6791	0.4320	0.1712	0.023*
H2C	0.7504	0.3487	0.0843	0.023*
C1	0.2944 (2)	0.0167 (2)	0.40370 (12)	0.0136 (3)
C2	0.3868 (2)	0.1673 (2)	0.49778 (12)	0.0142 (3)
H2A	0.3309	0.1630	0.5630	0.017*
C3	0.5666 (2)	0.3255 (2)	0.49154 (12)	0.0150 (3)
H3A	0.6328	0.4284	0.5536	0.018*
C4	0.6474 (2)	0.3303 (2)	0.39298 (12)	0.0146 (3)
H4	0.7681	0.4360	0.3899	0.018*
C5	0.5485 (2)	0.1772 (2)	0.29839 (12)	0.0137 (3)
C6	0.3711 (2)	0.0183 (2)	0.30398 (12)	0.0139 (3)
H6	0.3046	-0.0853	0.2423	0.017*
C7	0.6363 (2)	0.1664 (2)	0.19179 (12)	0.0142 (3)
H7	0.5242	0.0828	0.1264	0.017*
C8	0.7776 (2)	0.0740 (2)	0.21606 (12)	0.0146 (3)
H8A	0.8887	0.1565	0.2808	0.018*
H8B	0.7035	-0.0492	0.2399	0.018*
C9	0.8654 (2)	0.0398 (2)	0.11572 (12)	0.0136 (3)

supplementary materials

Cl1	0.33694 (5)	0.50865 (5)	0.14980 (3)	0.01593 (11)
O5	0.18796 (17)	0.43639 (18)	0.21983 (10)	0.0261 (3)
O7	0.31799 (18)	0.35386 (17)	0.06549 (11)	0.0270 (3)
O8	0.3095 (2)	0.6550 (2)	0.09212 (10)	0.0302 (3)
O6	0.53816 (16)	0.59598 (16)	0.22457 (10)	0.0209 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0238 (6)	0.0184 (6)	0.0237 (6)	-0.0008 (5)	0.0036 (5)	0.0009 (5)
O2	0.0208 (5)	0.0255 (6)	0.0254 (6)	0.0112 (5)	0.0135 (5)	0.0098 (5)
O3	0.0227 (5)	0.0280 (6)	0.0151 (5)	0.0186 (5)	0.0074 (4)	0.0048 (4)
O4	0.0196 (5)	0.0206 (5)	0.0144 (5)	0.0130 (4)	0.0051 (4)	0.0031 (4)
N1	0.0138 (6)	0.0171 (6)	0.0197 (6)	0.0082 (5)	0.0035 (5)	0.0062 (5)
N2	0.0186 (6)	0.0164 (6)	0.0148 (5)	0.0099 (5)	0.0060 (5)	0.0034 (5)
C1	0.0119 (6)	0.0146 (7)	0.0169 (6)	0.0079 (5)	0.0031 (5)	0.0049 (5)
C2	0.0155 (6)	0.0176 (7)	0.0140 (6)	0.0105 (6)	0.0056 (5)	0.0047 (5)
C3	0.0162 (6)	0.0136 (7)	0.0156 (6)	0.0083 (5)	0.0016 (5)	-0.0001 (5)
C4	0.0121 (6)	0.0124 (6)	0.0193 (7)	0.0050 (5)	0.0045 (5)	0.0032 (5)
C5	0.0153 (6)	0.0153 (7)	0.0145 (6)	0.0098 (5)	0.0054 (5)	0.0032 (5)
C6	0.0148 (6)	0.0140 (6)	0.0145 (6)	0.0086 (5)	0.0017 (5)	0.0012 (5)
C7	0.0144 (6)	0.0135 (6)	0.0146 (6)	0.0061 (5)	0.0047 (5)	0.0007 (5)
C8	0.0167 (6)	0.0153 (6)	0.0148 (6)	0.0084 (5)	0.0073 (5)	0.0039 (5)
C9	0.0122 (6)	0.0119 (6)	0.0157 (6)	0.0045 (5)	0.0041 (5)	0.0010 (5)
Cl1	0.01637 (17)	0.01607 (18)	0.01561 (17)	0.00828 (13)	0.00303 (13)	-0.00028 (13)
O5	0.0183 (5)	0.0306 (6)	0.0253 (6)	0.0070 (5)	0.0087 (5)	-0.0011 (5)
O7	0.0272 (6)	0.0203 (6)	0.0294 (6)	0.0086 (5)	0.0082 (5)	-0.0080 (5)
O8	0.0498 (8)	0.0372 (7)	0.0175 (5)	0.0332 (6)	0.0043 (5)	0.0059 (5)
O6	0.0167 (5)	0.0202 (5)	0.0231 (5)	0.0072 (4)	0.0013 (4)	0.0023 (4)

Geometric parameters (\AA , $^\circ$)

O1—N1	1.2288 (17)	C3—H3A	0.9300
O2—N1	1.2255 (17)	C4—C5	1.400 (2)
O3—C9	1.3011 (18)	C4—H4	0.9300
O3—H3	0.8200	C5—C6	1.384 (2)
O4—C9	1.2296 (18)	C5—C7	1.5217 (18)
N1—C1	1.4670 (18)	C6—H6	0.9300
N2—C7	1.5075 (18)	C7—C8	1.519 (2)
N2—H2D	0.8900	C7—H7	0.9800
N2—H2B	0.8900	C8—C9	1.5000 (18)
N2—H2C	0.8900	C8—H8A	0.9700
C1—C2	1.384 (2)	C8—H8B	0.9700
C1—C6	1.3889 (19)	Cl1—O7	1.4354 (11)
C2—C3	1.394 (2)	Cl1—O5	1.4396 (12)
C2—H2A	0.9300	Cl1—O8	1.4455 (12)
C3—C4	1.391 (2)	Cl1—O6	1.4486 (11)
C9—O3—H3	109.5	C5—C6—C1	118.60 (13)

O2—N1—O1	123.17 (13)	C5—C6—H6	120.7
O2—N1—C1	118.90 (12)	C1—C6—H6	120.7
O1—N1—C1	117.91 (12)	N2—C7—C8	111.06 (11)
C7—N2—H2D	109.5	N2—C7—C5	111.99 (11)
C7—N2—H2B	109.5	C8—C7—C5	108.58 (11)
H2D—N2—H2B	109.5	N2—C7—H7	108.4
C7—N2—H2C	109.5	C8—C7—H7	108.4
H2D—N2—H2C	109.5	C5—C7—H7	108.4
H2B—N2—H2C	109.5	C9—C8—C7	115.16 (12)
C2—C1—C6	123.03 (13)	C9—C8—H8A	108.5
C2—C1—N1	119.16 (12)	C7—C8—H8A	108.5
C6—C1—N1	117.78 (13)	C9—C8—H8B	108.5
C1—C2—C3	117.80 (13)	C7—C8—H8B	108.5
C1—C2—H2A	121.1	H8A—C8—H8B	107.5
C3—C2—H2A	121.1	O4—C9—O3	125.13 (13)
C4—C3—C2	120.38 (13)	O4—C9—C8	122.98 (13)
C4—C3—H3A	119.8	O3—C9—C8	111.90 (12)
C2—C3—H3A	119.8	O7—C11—O5	110.51 (7)
C3—C4—C5	120.49 (13)	O7—C11—O8	110.15 (7)
C3—C4—H4	119.8	O5—C11—O8	109.25 (8)
C5—C4—H4	119.8	O7—C11—O6	109.10 (7)
C6—C5—C4	119.70 (13)	O5—C11—O6	109.02 (7)
C6—C5—C7	116.93 (12)	O8—C11—O6	108.78 (8)
C4—C5—C7	123.14 (12)		
O2—N1—C1—C2	−1.71 (19)	C7—C5—C6—C1	175.40 (12)
O1—N1—C1—C2	179.73 (13)	C2—C1—C6—C5	0.3 (2)
O2—N1—C1—C6	176.20 (12)	N1—C1—C6—C5	−177.56 (12)
O1—N1—C1—C6	−2.37 (19)	C6—C5—C7—N2	148.40 (12)
C6—C1—C2—C3	−0.9 (2)	C4—C5—C7—N2	−37.23 (18)
N1—C1—C2—C3	176.88 (12)	C6—C5—C7—C8	−88.61 (15)
C1—C2—C3—C4	0.5 (2)	C4—C5—C7—C8	85.76 (16)
C2—C3—C4—C5	0.6 (2)	N2—C7—C8—C9	−60.66 (16)
C3—C4—C5—C6	−1.2 (2)	C5—C7—C8—C9	175.79 (12)
C3—C4—C5—C7	−175.45 (13)	C7—C8—C9—O4	0.9 (2)
C4—C5—C6—C1	0.8 (2)	C7—C8—C9—O3	−179.00 (12)

Hydrogen-bond geometry (Å, °)


D—H···A	D—H	H···A	D···A	D—H···A
C8—H8B···O6 ⁱ	0.97	2.46	3.3742 (19)	157
C8—H8A···O2 ⁱⁱ	0.97	2.56	3.3053 (19)	134
C6—H6···O8 ⁱ	0.93	2.59	3.4263 (19)	151
C4—H4···O1 ⁱⁱⁱ	0.93	2.48	3.3858 (19)	163
O3—H3···O4 ^{iv}	0.82	1.82	2.6381 (14)	172
N2—H2D···O5 ^v	0.89	2.30	3.0542 (17)	142
N2—H2D···O1 ⁱⁱⁱ	0.89	2.26	2.9200 (17)	130
N2—H2C···O8 ^{vi}	0.89	2.06	2.9159 (17)	162

supplementary materials

N2—H2C···O4	0.89	2.45	2.9317 (16)	114
N2—H2B···O7	0.89	2.57	3.1795 (18)	126
N2—H2B···O6	0.89	2.09	2.9720 (17)	172

Symmetry codes: (i) $x, y-1, z$; (ii) $-x+1, -y, -z+1$; (iii) $x+1, y+1, z$; (iv) $-x+2, -y, -z$; (v) $x+1, y, z$; (vi) $-x+1, -y+1, -z$.

Fig. 1



supplementary materials

Fig. 2

