

5-Chlorobenzothiazole-2-spiro-3'-indolin-2'-one

Mehmet Akkurt,^{a*} Selvi Karaca,^a Görkem Ermut,^b Nilgün Karal^b and Orhan Büyükgüngör^c^aDepartment of Physics, Faculty of Arts and Sciences, Erciyes University, 38039 Kayseri, Turkey, ^bDepartment of Pharmaceutical Chemistry, Faculty of Pharmacy, İstanbul University, 34116 Beyazıt-İstanbul, Turkey, and ^cDepartment of Physics, Faculty of Arts and Sciences, Ondokuz Mayıs University, 55139 Samsun, Turkey
Correspondence e-mail: akkurt@erciyes.edu.tr


Received 16 December 2009; accepted 11 January 2010

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.033; wR factor = 0.080; data-to-parameter ratio = 14.7.

The title compound, $\text{C}_{14}\text{H}_9\text{ClN}_2\text{OS}$, crystallizes with two unique molecules, *A* and *B*, in the asymmetric unit. The five-membered rings of the benzothiazole groups in both molecules adopt an envelope conformation [puckering parameters: $q_2 = 0.242$ (1) Å and $\varphi_2 = 217.5$ (4)° for *A*, and $q_2 = 0.234$ (1) Å and $\varphi_2 = 37.7$ (4)° for *B*]. The five-membered rings of the indolinone groups in both molecules are also not planar, with a twisted conformation [puckering parameters are $q_2 = 0.112$ (2) Å and $\varphi_2 = 126.3$ (8)° for *A*, and $q_2 = 0.108$ (2) Å and $\varphi_2 = 306.4$ (9)° for *B*]. In the crystal structure, there are intermolecular $\text{N}-\text{H}\cdots\text{O}$, $\text{N}-\text{H}\cdots\text{S}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen-bonding interactions, forming the layers propagating normal to *c*.

Related literature

For general background to and applications of 1*H*-indole-2,3-dione derivatives, see: Alam & Nawwar (2002); Cho *et al.* (2008); Da-Silva *et al.* (2001); Dandia *et al.* (1990); Hall *et al.* (2009); Joshi *et al.* (1990); Kumar *et al.* (2008); Quenelle *et al.* (2006); Vine *et al.* (2007, 2009); Čaleta *et al.* (2009). For bond-length data, see: Allen *et al.* (1987). For puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{14}\text{H}_9\text{ClN}_2\text{OS}$
 $M_r = 288.75$
 Monoclinic, $P2_1/c$
 $a = 12.8421$ (6) Å
 $b = 9.1159$ (3) Å
 $c = 22.1553$ (9) Å
 $\beta = 97.051$ (3)°
 $V = 2574.1$ (2) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.45$ mm⁻¹
 $T = 295$ K
 $0.77 \times 0.49 \times 0.19$ mm

Data collection

Stoe IPDS 2 diffractometer
 Absorption correction: integration
 (*X-RED32*; Stoe & Cie, 2002)
 $T_{\min} = 0.723$, $T_{\max} = 0.919$
 23976 measured reflections
 5271 independent reflections
 3987 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.080$
 $S = 1.01$
 5271 reflections
 359 parameters
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.17$ e Å⁻³
 $\Delta\rho_{\min} = -0.19$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1A \cdots O2 ⁱ	0.90 (3)	1.95 (3)	2.840 (2)	171 (2)
N2—H2A \cdots S1 ⁱⁱ	0.90 (2)	2.61 (2)	3.506 (1)	177 (2)
N3—H3A \cdots O1 ⁱⁱⁱ	0.89 (2)	1.99 (2)	2.867 (2)	166 (2)
N4—H4A \cdots S2 ^{iv}	0.89 (2)	2.63 (2)	3.511 (1)	176 (2)
C3—H3 \cdots O1 ^v	0.93	2.53	3.418 (2)	161

Symmetry codes: (i) $x, y - 1, z$; (ii) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $x, y + 1, z$; (iv) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (v) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *X-AREA* (Stoe & Cie, 2002); cell refinement: *X-AREA*; data reduction: *X-RED32* (Stoe & Cie, 2002); program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors acknowledge the Faculty of Arts and Sciences, Ondokuz Mayıs University, Turkey, for the use of the Stoe IPDS 2 diffractometer (purchased under grant F.279 of the University Research Fund).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IM2170).

References

- Alam, Y. A. & Nawwar, G. A. M. (2002). *Heteroat. Chem.* **13**, 207–210.
 Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
 Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
 Čaleta, I., Kralj, M., Marjanović, M., Bertoša, B., Tomić, S., Pavlović, G., Pavelić, K. & Karminski-Zamola, G. (2009). *J. Med. Chem.* **52**, 1744–1756.
 Cho, Y., Loerger, T. R. & Sacchettini, J. C. (2008). *J. Med. Chem.* **51**, 5984–5992.
 Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
 Dandia, A., Khanna, S. & Joshi, K. C. (1990). *J. Indian Chem. Soc.* **67**, 824–826.

- Da-Silva, J. F. M., Garden, S. J. & Pinto, A. C. (2001). *J. Braz. Chem. Soc.* **12**, 273–324.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Hall, M. D., Salam, N. K., Hellawell, J. L., Fales, H. M., Kensler, C. B., Ludwig, J. A., Szakács, G., Hibbs, D. E. & Gottesman, M. M. (2009). *J. Med. Chem.* **52**, 3191–3204.
- Joshi, K. C., Dandia, A. & Khanna, S. (1990). *Indian J. Chem. Soc. Sect. B*, **29**, 824–829.
- Kumar, R. R., Perumal, S., Senthilkumar, P., Yogeewari, P. & Sriram, D. (2008). *J. Med. Chem.* **51**, 5731–5735.
- Quenelle, D. C., Keith, K. A. & Kern, K. E. R. (2006). *Antiviral Res.* **71**, 24–30.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Stoe & Cie (2002). *X-AREA* and *X-RED32*. Stoe & Cie, Darmstadt, Germany.
- Vine, K. L., Locke, J. M., Ranson, M., Pyne, S. G. & Bremner, J. B. (2007). *Bioorg. Med. Chem.* **15**, 931–938.
- Vine, K. L., Matesic, L., Locke, J. M., Ranson, M. & Skropeta, D. (2009). *Anti-Cancer Agents Med. Chem.* **9**, 397–414.

supplementary materials

Acta Cryst. (2010). E66, o399-o400 [doi:10.1107/S1600536810001285]

5-Chlorobenzothiazole-2-spiro-3'-indolin-2'-one

M. Akkurt, S. Karaca, G. Ermut, N. Karali and O. Büyükgüngör

Comment

1*H*-Indole-2,3-dione is a synthetically versatile molecule which has led to an array of derivatives displaying a broad spectrum of biological properties including anticancer, antiviral, antituberculosis and antibacterial activities (Vine *et al.*, 2007, 2009; Quenelle *et al.*, 2006). Investigation of the structure-activity relationships in 2-indolinones revealed that cyclization to thiazolines and spiroindolinones are associated with increased activity against a range of human cancer cell lines, various bacteria and viruses (Hall *et al.*, 2009; Kumar *et al.*, 2008). A large number of 2-arylbenzothiazoles have been prepared because of their wide pharmacological potential. This important class of compounds has significant anticancer and antimicrobial properties (Ćaleta *et al.*, 2009; Cho *et al.*, 2008). The reactivity of 1*H*-indole-2,3-dione towards 2-aminothiophenol has been the subject of a number of reports and some of the products obtained are quite interesting. The first results reported that 1*H*-indole-2,3-dione furnished benzothiazinone, indolobenzothiazide and spiro benzothiazole when the reaction was carried out in dry xylene in the presence of anhydrous zinc chloride under reflux. On the other hand, the reaction of 1-methyl-1*H*-indole-2,3-dione with 2-amino thiophenol under the same conditions furnished solely the spiro compound (Joshi *et al.*, 1990; Dandia *et al.*, 1990; Da-Silva *et al.*, 2001). In addition, there is one report on the reaction of 1*H*-indole-2,3-dione with 2-aminothiophenol in ethanol yielding a single spirobenzothiazole (Alam & Nawwar, 2002). Promoted by the above observations and in continuation of our study on the indolinone derivatives, we synthesized the title compound (**3**) by incorporating the benzothiazole moiety. Thus spectroscopic and X-ray diffraction studies were carried out on (**3**) to determine the spiro benzothiazole structure.

Fig. 1 shows the two crystallographically independent molecules in the asymmetric unit. Bond lengths in both molecules are within normal ranges (Allen *et al.*, 1987). The five-membered rings S1/N2/C8/C9/C14 and S2/N4/C22/C23/C28 of the benzothiazole groups in both molecules A and B [A: S1/C11/O1/N1/N2/C1–C14 and B: S2/C12/O2/N3/N4/C15–C28] adopt an envelope conformation with atom C8 at the flap for molecule A [puckering parameters are $q_2 = 0.242$ (1) Å and $\varphi_2 = 217.5$ (4)° (Cremer & Pople, 1975)] and with atom C22 at the flap for molecule B [puckering parameters are $q_2 = 0.234$ (1) Å and $\varphi_2 = 37.7$ (4)°]. The five-membered rings N1/C1/C2/C7/C8 and N3/C15/C20–C22 of the indolinone groups in both molecules A and B also are not planar, with twisted C7–C8 and C21–C22 bonds, respectively, [puckering parameters are $q_2 = 0.112$ (2) Å and $\varphi_2 = 126.3$ (8)° for A, and $q_2 = 0.108$ (2) Å and $\varphi_2 = 306.4$ (9)° for B].

The torsion angles N1–C7–C8–N2, C2–C8–N2–C14 in A and N3–C21–C22–N4, C20–C22–N4–C28 in B are 141.0 (1)°, 148.5 (1)° and -140.3 (1)°, -147.8 (1)°, respectively. Thus, they adopt +anti-clinal (+*ac*) and -anti-clinal (-*ac*) conformations, for molecules A and B, respectively.

The crystal packing is stabilized by intermolecular N–H···O, N–H···S and C–H···O hydrogen bonding interactions, forming the layers of molecules which are parallel to the (001) planes (Table 1 and Fig. 2).

Experimental

To a solution of 1*H*-indole-2,3-dione **1** (3.5 mmol) in absolute ethanol (15 ml) was added 2-aminothiophenol **2** (3.5 mmol). The mixture was heated under reflux for 5 h. The solid thus obtained (**3**) was filtered, dried and recrystallized from ethanol (Alam & Nawwar, 2002). Yield: 86%; m.p.: 514 K; IR (KBr) ν (cm⁻¹): 3281, 3149 (N—H), 1728(C=O); ¹H-NMR (DMSO-d₆, 500 MHz) δ (p.p.m.): 6.50 (1*H*, d, *J* = 0.96 Hz, C₁₃—H), 6.62 (1*H*, dd, *J* = 8.30, 2.44 Hz, C₁₁—H), 6.84 (1*H*, d, *J* = 7.81 Hz, C₆—H), 7.03 (1*H*, d, *J* = 8.30 Hz, C₁₀—H), 7.05 (1*H*, dt, *J* = 7.81 Hz, C₄—H), 7.29 (1*H*, dt, *J* = 7.81 Hz, C₅—H), 7.55 (1*H*, d, *J* = 2.92 Hz, C₃—H), 7.56 (1*H*, s, N₂—H), 10.39 (1*H*, s, N₁—H); ¹³C-NMR (HSQC) (125 MHz)(DMSO-d₆/TMS) δ (p.p.m.): 75.64 (C₈), 108.40 (C₁₃), 118.52 (C₁₁), 110.88 (C₆), 122.70 (C₁₀), 123.28 (C₄), 123.99 (C₂), 126.38 (C₃), 129.86(C₉), 130.92 (C₁₂), 131.43 (C₅), 142.03 (C₁), 149.43 (C₁₄), 176.53 (C₇). MS (ESI+) *m/z* (%): 289 (MH⁺, 35), 287 (100). Analysis calculated for C₁₄H₉ClN₂OS: C 58.23, H 3.14, N 9.70%. Found: C 58.06, H 3.14, N 9.52%.

Refinement

H atoms bound to N atoms were located from a difference Fourier map and refined freely. H atoms bound to C atoms were positioned geometrically with C—H = 0.93 Å and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The title compound (**3**), with the atom-numbering scheme, intramolecular H-bonds and 50% probability displacement ellipsoids.




Fig. 2. The packing and hydrogen bonding of the title compound (**3**) down the *b* axis. H atoms not involved in hydrogen bonding have been omitted.

5-Chlorobenzothiazole-2-spiro-3'-indolin-2'-one

Crystal data

C₁₄H₉ClN₂OS

$M_r = 288.75$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 12.8421$ (6) Å

$F(000) = 1184$

$D_x = 1.490$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 33214 reflections

$\theta = 1.6$ – 28.0°

$b = 9.1159 (3) \text{ \AA}$
 $c = 22.1553 (9) \text{ \AA}$
 $\beta = 97.051 (3)^\circ$
 $V = 2574.1 (2) \text{ \AA}^3$
 $Z = 8$

$\mu = 0.45 \text{ mm}^{-1}$
 $T = 295 \text{ K}$
 Prism, yellow
 $0.77 \times 0.49 \times 0.19 \text{ mm}$

Data collection

Stoe IPDS 2
 diffractometer
 Radiation source: sealed X-ray tube, 12 x 0.4 mm
 long-fine focus
 plane graphite
 Detector resolution: 6.67 pixels mm⁻¹
 ω scans
 Absorption correction: integration
 (X-RED32; Stoe & Cie, 2002)
 $T_{\min} = 0.723$, $T_{\max} = 0.919$
 23976 measured reflections

5271 independent reflections
 3987 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$
 $\theta_{\max} = 26.5^\circ$, $\theta_{\min} = 1.9^\circ$
 $h = -14 \rightarrow 16$
 $k = -11 \rightarrow 11$
 $l = -27 \rightarrow 27$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.080$
 $S = 1.01$
 5271 reflections
 359 parameters
 0 restraints

Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0445P)^2 + 0.0864P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.17 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.19 \text{ e \AA}^{-3}$

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating $-R$ -factor-obs *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

supplementary materials

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	−0.21534 (5)	0.20753 (8)	0.44750 (3)	0.0897 (3)
S1	0.14151 (3)	0.34132 (4)	0.29015 (2)	0.0398 (1)
O1	0.15538 (9)	−0.02243 (11)	0.29802 (5)	0.0468 (4)
N1	0.20968 (12)	0.04018 (14)	0.20602 (7)	0.0459 (5)
N2	−0.02266 (11)	0.17509 (14)	0.25719 (6)	0.0386 (4)
C1	0.17319 (13)	0.13376 (17)	0.15826 (7)	0.0439 (5)
C2	0.09339 (13)	0.22368 (16)	0.17481 (7)	0.0407 (5)
C3	0.04229 (16)	0.32123 (18)	0.13402 (8)	0.0528 (6)
C4	0.0736 (2)	0.3287 (2)	0.07608 (9)	0.0684 (8)
C5	0.1542 (2)	0.2413 (2)	0.06049 (9)	0.0682 (8)
C6	0.20529 (17)	0.1426 (2)	0.10113 (9)	0.0581 (7)
C7	0.15275 (12)	0.05484 (15)	0.25322 (7)	0.0384 (5)
C8	0.08181 (12)	0.19188 (15)	0.24007 (7)	0.0371 (4)
C9	0.04814 (13)	0.31467 (15)	0.34106 (7)	0.0400 (5)
C10	0.04669 (15)	0.3771 (2)	0.39753 (8)	0.0538 (6)
C11	−0.03608 (18)	0.3447 (2)	0.43046 (9)	0.0603 (7)
C12	−0.11329 (15)	0.2499 (2)	0.40578 (8)	0.0541 (6)
C13	−0.11335 (13)	0.18672 (17)	0.34914 (8)	0.0446 (5)
C14	−0.03182 (12)	0.22070 (15)	0.31617 (7)	0.0366 (4)
Cl2	0.71172 (5)	0.55718 (7)	0.05591 (3)	0.0844 (2)
S2	0.35504 (3)	0.42746 (4)	0.21381 (2)	0.0400 (1)
O2	0.34292 (10)	0.79114 (12)	0.20783 (5)	0.0470 (4)
N3	0.29057 (12)	0.72689 (14)	0.30029 (7)	0.0456 (4)
N4	0.52018 (11)	0.59219 (14)	0.24673 (6)	0.0388 (4)
C15	0.32736 (13)	0.63104 (16)	0.34761 (8)	0.0441 (5)
C16	0.29684 (17)	0.6217 (2)	0.40483 (9)	0.0589 (7)
C17	0.3485 (2)	0.5204 (2)	0.44442 (9)	0.0689 (8)
C18	0.4273 (2)	0.4313 (2)	0.42740 (9)	0.0665 (8)
C19	0.45735 (15)	0.44084 (18)	0.36933 (8)	0.0505 (6)
C20	0.40570 (13)	0.54115 (16)	0.32944 (7)	0.0406 (5)
C21	0.34576 (12)	0.71277 (15)	0.25252 (7)	0.0381 (5)
C22	0.41620 (12)	0.57500 (14)	0.26426 (7)	0.0363 (4)
C23	0.44832 (13)	0.45380 (15)	0.16277 (7)	0.0405 (5)
C24	0.44918 (16)	0.3916 (2)	0.10620 (8)	0.0535 (6)
C25	0.53209 (18)	0.4225 (2)	0.07331 (9)	0.0616 (7)
C26	0.60934 (15)	0.5167 (2)	0.09761 (8)	0.0535 (6)
C27	0.60988 (13)	0.58121 (17)	0.15439 (8)	0.0443 (5)
C28	0.52871 (12)	0.54685 (15)	0.18761 (7)	0.0368 (4)
H1A	0.257 (2)	−0.032 (3)	0.2053 (10)	0.081 (7)*
H2A	−0.0558 (16)	0.091 (2)	0.2451 (8)	0.056 (5)*
H3	−0.01150	0.38030	0.14480	0.0630*
H4	0.04000	0.39320	0.04750	0.0820*
H5	0.17450	0.24920	0.02170	0.0820*
H6	0.25940	0.08410	0.09040	0.0700*
H10	0.10020	0.43990	0.41350	0.0650*

H11	-0.03920	0.38640	0.46850	0.0720*
H13	-0.16670	0.12320	0.33360	0.0530*
H3A	0.2402 (19)	0.794 (2)	0.3016 (9)	0.070 (6)*
H4A	0.5540 (16)	0.674 (2)	0.2583 (8)	0.050 (5)*
H16	0.24380	0.68090	0.41650	0.0710*
H17	0.32980	0.51190	0.48350	0.0830*
H18	0.46040	0.36450	0.45510	0.0800*
H19	0.51040	0.38170	0.35770	0.0610*
H24	0.39510	0.32980	0.09020	0.0640*
H25	0.53510	0.37990	0.03540	0.0740*
H27	0.66300	0.64540	0.16960	0.0530*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0863 (5)	0.1113 (5)	0.0817 (4)	-0.0106 (4)	0.0518 (4)	0.0006 (3)
S1	0.0346 (2)	0.0328 (2)	0.0528 (2)	-0.0029 (2)	0.0087 (2)	-0.0030 (2)
O1	0.0451 (7)	0.0386 (5)	0.0585 (7)	0.0065 (5)	0.0138 (6)	0.0070 (5)
N1	0.0440 (8)	0.0377 (7)	0.0591 (9)	0.0076 (6)	0.0185 (7)	-0.0010 (6)
N2	0.0303 (7)	0.0369 (6)	0.0492 (8)	-0.0010 (5)	0.0073 (6)	-0.0045 (6)
C1	0.0438 (9)	0.0372 (7)	0.0527 (9)	-0.0045 (7)	0.0143 (7)	-0.0059 (7)
C2	0.0425 (9)	0.0345 (7)	0.0462 (9)	-0.0026 (6)	0.0095 (7)	-0.0031 (6)
C3	0.0586 (12)	0.0463 (9)	0.0536 (10)	0.0033 (8)	0.0075 (9)	0.0017 (7)
C4	0.0886 (18)	0.0634 (12)	0.0530 (11)	-0.0042 (11)	0.0074 (11)	0.0111 (9)
C5	0.0888 (17)	0.0686 (12)	0.0511 (11)	-0.0163 (12)	0.0241 (11)	-0.0036 (9)
C6	0.0643 (13)	0.0544 (10)	0.0604 (11)	-0.0060 (9)	0.0272 (10)	-0.0081 (9)
C7	0.0326 (8)	0.0307 (7)	0.0530 (9)	-0.0008 (6)	0.0092 (7)	-0.0022 (6)
C8	0.0336 (8)	0.0315 (7)	0.0468 (8)	0.0016 (6)	0.0080 (7)	-0.0016 (6)
C9	0.0371 (9)	0.0345 (7)	0.0487 (9)	0.0003 (6)	0.0060 (7)	0.0005 (6)
C10	0.0572 (12)	0.0538 (9)	0.0505 (10)	-0.0064 (9)	0.0072 (8)	-0.0098 (8)
C11	0.0680 (14)	0.0663 (12)	0.0488 (10)	0.0010 (10)	0.0163 (9)	-0.0085 (9)
C12	0.0537 (12)	0.0582 (10)	0.0543 (10)	0.0052 (9)	0.0225 (9)	0.0079 (8)
C13	0.0369 (9)	0.0421 (8)	0.0561 (10)	0.0020 (7)	0.0109 (8)	0.0049 (7)
C14	0.0320 (8)	0.0329 (7)	0.0453 (8)	0.0044 (6)	0.0063 (6)	0.0016 (6)
Cl2	0.0786 (4)	0.1071 (4)	0.0763 (4)	-0.0002 (3)	0.0450 (3)	0.0068 (3)
S2	0.0353 (2)	0.0323 (2)	0.0531 (2)	-0.0031 (2)	0.0084 (2)	-0.0037 (2)
O2	0.0452 (7)	0.0385 (6)	0.0590 (7)	0.0070 (5)	0.0134 (6)	0.0055 (5)
N3	0.0422 (8)	0.0373 (6)	0.0605 (9)	0.0076 (6)	0.0194 (7)	-0.0010 (6)
N4	0.0308 (7)	0.0364 (6)	0.0499 (8)	-0.0023 (5)	0.0081 (6)	-0.0070 (6)
C15	0.0448 (10)	0.0357 (7)	0.0535 (9)	-0.0046 (7)	0.0133 (7)	-0.0041 (7)
C16	0.0678 (13)	0.0531 (10)	0.0604 (11)	-0.0056 (9)	0.0269 (10)	-0.0083 (9)
C17	0.0906 (17)	0.0673 (12)	0.0523 (11)	-0.0088 (12)	0.0234 (11)	0.0001 (9)
C18	0.0848 (17)	0.0620 (11)	0.0517 (11)	0.0016 (11)	0.0048 (10)	0.0102 (9)
C19	0.0544 (11)	0.0429 (8)	0.0534 (10)	0.0025 (7)	0.0037 (8)	0.0004 (7)
C20	0.0404 (9)	0.0350 (7)	0.0470 (9)	-0.0031 (6)	0.0084 (7)	-0.0042 (6)
C21	0.0313 (8)	0.0311 (7)	0.0526 (9)	-0.0008 (6)	0.0080 (7)	-0.0029 (6)
C22	0.0319 (8)	0.0294 (7)	0.0481 (8)	0.0012 (6)	0.0069 (7)	-0.0025 (6)
C23	0.0382 (9)	0.0343 (7)	0.0495 (9)	0.0023 (6)	0.0075 (7)	-0.0002 (6)

supplementary materials

C24	0.0593 (12)	0.0509 (9)	0.0502 (9)	-0.0062 (8)	0.0062 (8)	-0.0075 (8)
C25	0.0736 (15)	0.0652 (12)	0.0482 (10)	0.0052 (10)	0.0169 (10)	-0.0062 (9)
C26	0.0538 (11)	0.0563 (10)	0.0538 (10)	0.0076 (8)	0.0204 (9)	0.0078 (8)
C27	0.0360 (9)	0.0429 (8)	0.0551 (10)	0.0027 (7)	0.0102 (7)	0.0051 (7)
C28	0.0324 (8)	0.0312 (7)	0.0469 (8)	0.0044 (6)	0.0054 (6)	0.0013 (6)

Geometric parameters (Å, °)

C11—C12	1.737 (2)	C11—C12	1.377 (3)
C12—C26	1.737 (2)	C12—C13	1.381 (2)
S1—C9	1.7609 (17)	C13—C14	1.383 (2)
S1—C8	1.8617 (15)	C3—H3	0.9300
S2—C22	1.8600 (15)	C4—H4	0.9300
S2—C23	1.7619 (17)	C5—H5	0.9300
O1—C7	1.2142 (18)	C6—H6	0.9300
O2—C21	1.2179 (18)	C10—H10	0.9300
N1—C7	1.354 (2)	C11—H11	0.9300
N1—C1	1.395 (2)	C13—H13	0.9300
N2—C14	1.390 (2)	C15—C16	1.375 (3)
N2—C8	1.446 (2)	C15—C20	1.395 (2)
N1—H1A	0.90 (3)	C16—C17	1.385 (3)
N2—H2A	0.901 (19)	C17—C18	1.386 (3)
N3—C15	1.402 (2)	C18—C19	1.391 (3)
N3—C21	1.350 (2)	C19—C20	1.383 (2)
N4—C28	1.391 (2)	C20—C22	1.499 (2)
N4—C22	1.444 (2)	C21—C22	1.551 (2)
N3—H3A	0.89 (2)	C23—C28	1.396 (2)
N4—H4A	0.885 (19)	C23—C24	1.377 (2)
C1—C2	1.396 (2)	C24—C25	1.391 (3)
C1—C6	1.381 (3)	C25—C26	1.371 (3)
C2—C8	1.500 (2)	C26—C27	1.388 (2)
C2—C3	1.376 (2)	C27—C28	1.384 (2)
C3—C4	1.393 (3)	C16—H16	0.9300
C4—C5	1.383 (3)	C17—H17	0.9300
C5—C6	1.381 (3)	C18—H18	0.9300
C7—C8	1.553 (2)	C19—H19	0.9300
C9—C14	1.398 (2)	C24—H24	0.9300
C9—C10	1.377 (2)	C25—H25	0.9300
C10—C11	1.393 (3)	C27—H27	0.9300
C11...C12 ⁱ	3.6085 (10)	C13...O2 ^{vii}	3.206 (2)
C12...C11 ⁱⁱ	3.6085 (10)	C13...C21 ^{vii}	3.520 (2)
C12...H5 ⁱⁱⁱ	2.9700	C13...C3 ^{vii}	3.463 (2)
S1...O1	3.3241 (11)	C14...O1	3.3303 (18)
S1...N1	3.4896 (14)	C15...S1	3.6805 (16)
S1...N2 ^{iv}	3.5063 (14)	C15...C25 ^{ix}	3.551 (3)
S1...C15	3.6805 (16)	C16...C25 ^{ix}	3.510 (3)
S1...S2	3.4836 (6)	C19...C27 ^v	3.414 (2)

S2...N4 ^v	3.5113 (14)	C21...C13 ^{iv}	3.520 (2)
S2...S1	3.4836 (6)	C25...C16 ^v	3.510 (3)
S2...O2	3.3208 (12)	C25...C15 ^v	3.551 (3)
S2...N3	3.4923 (14)	C25...C25 ⁱⁱⁱ	3.546 (3)
S2...C1	3.6649 (16)	C27...C19 ^{ix}	3.414 (2)
S1...H2A ^{iv}	2.606 (19)	C27...C7 ^{ix}	3.465 (2)
S1...H27 ^v	3.1200	C27...O1 ^{ix}	3.211 (2)
S2...H13 ^{iv}	3.0800	C28...O2	3.3339 (19)
S2...H4A ^v	2.628 (19)	C6...H24	3.0100
O1...N2	2.9632 (18)	C7...H3A ^{vi}	2.787 (19)
O1...N3 ^{vi}	2.8666 (18)	C7...H27 ^v	2.8700
O1...S1	3.3241 (11)	C16...H10	3.0500
O1...C3 ^{vii}	3.418 (2)	C21...H1A ^{viii}	2.74 (3)
O1...C14	3.3303 (18)	C21...H13 ^{iv}	2.9200
O1...C27 ^v	3.211 (2)	C23...H4A ^v	3.095 (18)
O2...C28	3.3339 (19)	C25...H25 ⁱⁱⁱ	3.0500
O2...N1 ^{viii}	2.8402 (18)	H1A...C21 ^{vi}	2.74 (3)
O2...N4	2.9550 (18)	H1A...O2 ^{vi}	1.95 (3)
O2...S2	3.3208 (12)	H2A...S1 ^{vii}	2.606 (19)
O2...C13 ^{iv}	3.206 (2)	H2A...H13	2.5800
O1...H3A ^{vi}	1.99 (2)	H3...O1 ^{iv}	2.5300
O1...H27 ^v	2.8100	H3A...C7 ^{viii}	2.787 (19)
O1...H3 ^{vii}	2.5300	H3A...O1 ^{viii}	1.99 (2)
O2...H1A ^{viii}	1.95 (3)	H4A...S2 ^{ix}	2.628 (19)
O2...H13 ^{iv}	2.7900	H4A...C23 ^{ix}	3.095 (18)
O2...H19 ^{ix}	2.6500	H4A...H27	2.5600
N1...S1	3.4896 (14)	H5...C12 ⁱⁱⁱ	2.9700
N1...O2 ^{vi}	2.8402 (18)	H10...C16	3.0500
N2...S1 ^{vii}	3.5063 (14)	H13...O2 ^{vii}	2.7900
N2...O1	2.9632 (18)	H13...C21 ^{vii}	2.9200
N3...O1 ^{viii}	2.8666 (18)	H13...S2 ^{vii}	3.0800
N3...S2	3.4923 (14)	H13...H2A	2.5800
N4...O2	2.9550 (18)	H19...O2 ^v	2.6500
N4...S2 ^{ix}	3.5113 (14)	H24...C6	3.0100
C1...S2	3.6649 (16)	H25...C25 ⁱⁱⁱ	3.0500
C3...C13 ^{iv}	3.463 (2)	H27...H4A	2.5600
C3...O1 ^{iv}	3.418 (2)	H27...S1 ^{ix}	3.1200
C6...C11 ^{vii}	3.495 (3)	H27...O1 ^{ix}	2.8100
C7...C27 ^v	3.465 (2)	H27...C7 ^{ix}	2.8700
C11...C6 ^{iv}	3.495 (3)		
C8—S1—C9	91.02 (7)	C5—C6—H6	121.00

supplementary materials

C22—S2—C23	90.97 (7)	C9—C10—H10	121.00
C1—N1—C7	111.26 (14)	C11—C10—H10	120.00
C8—N2—C14	113.78 (13)	C10—C11—H11	121.00
C7—N1—H1A	121.2 (15)	C12—C11—H11	120.00
C1—N1—H1A	126.9 (14)	C12—C13—H13	121.00
C8—N2—H2A	115.6 (13)	C14—C13—H13	121.00
C14—N2—H2A	116.3 (12)	C16—C15—C20	121.82 (16)
C15—N3—C21	111.45 (14)	N3—C15—C16	128.34 (16)
C22—N4—C28	113.79 (13)	N3—C15—C20	109.83 (15)
C21—N3—H3A	122.9 (13)	C15—C16—C17	117.16 (19)
C15—N3—H3A	125.6 (13)	C16—C17—C18	121.83 (19)
C28—N4—H4A	115.5 (12)	C17—C18—C19	120.67 (18)
C22—N4—H4A	116.8 (13)	C18—C19—C20	117.82 (17)
N1—C1—C2	110.43 (14)	C15—C20—C19	120.69 (15)
N1—C1—C6	128.38 (16)	C15—C20—C22	108.09 (13)
C2—C1—C6	121.19 (15)	C19—C20—C22	131.22 (15)
C3—C2—C8	131.49 (15)	O2—C21—N3	127.96 (14)
C1—C2—C8	107.61 (13)	O2—C21—C22	124.75 (14)
C1—C2—C3	120.89 (15)	N3—C21—C22	107.29 (12)
C2—C3—C4	117.94 (18)	S2—C22—C20	110.41 (10)
C3—C4—C5	120.76 (18)	S2—C22—C21	106.87 (10)
C4—C5—C6	121.53 (19)	S2—C22—N4	104.66 (10)
C1—C6—C5	117.68 (19)	C20—C22—C21	102.05 (12)
O1—C7—C8	125.24 (14)	N4—C22—C20	118.49 (13)
O1—C7—N1	127.63 (14)	N4—C22—C21	113.99 (12)
N1—C7—C8	107.13 (12)	S2—C23—C28	110.98 (11)
S1—C8—C2	110.54 (10)	C24—C23—C28	121.39 (16)
N2—C8—C2	118.57 (13)	S2—C23—C24	127.60 (13)
N2—C8—C7	114.05 (12)	C23—C24—C25	119.06 (17)
S1—C8—N2	104.39 (10)	C24—C25—C26	118.99 (18)
C2—C8—C7	102.19 (12)	C12—C26—C25	118.86 (15)
S1—C8—C7	106.72 (10)	C12—C26—C27	118.18 (14)
S1—C9—C14	110.91 (11)	C25—C26—C27	122.96 (18)
C10—C9—C14	121.38 (16)	C26—C27—C28	117.77 (15)
S1—C9—C10	127.68 (13)	N4—C28—C27	125.83 (14)
C9—C10—C11	119.00 (17)	C23—C28—C27	119.79 (14)
C10—C11—C12	118.93 (18)	N4—C28—C23	114.33 (14)
C11—C12—C13	122.87 (18)	C15—C16—H16	121.00
C11—C12—C11	118.80 (15)	C17—C16—H16	121.00
C11—C12—C13	118.33 (14)	C16—C17—H17	119.00
C12—C13—C14	118.07 (15)	C18—C17—H17	119.00
N2—C14—C13	125.90 (14)	C17—C18—H18	120.00
C9—C14—C13	119.73 (14)	C19—C18—H18	120.00
N2—C14—C9	114.29 (14)	C18—C19—H19	121.00
C4—C3—H3	121.00	C20—C19—H19	121.00
C2—C3—H3	121.00	C23—C24—H24	120.00
C3—C4—H4	120.00	C25—C24—H24	120.00
C5—C4—H4	120.00	C24—C25—H25	121.00
C6—C5—H5	119.00	C26—C25—H25	121.00

C4—C5—H5	119.00	C26—C27—H27	121.00
C1—C6—H6	121.00	C28—C27—H27	121.00
C9—S1—C8—C7	-101.67 (11)	O1—C7—C8—S1	75.43 (17)
C8—S1—C9—C10	171.57 (16)	C10—C9—C14—N2	175.71 (15)
C8—S1—C9—C14	-10.32 (12)	C14—C9—C10—C11	0.4 (3)
C9—S1—C8—N2	19.41 (10)	C10—C9—C14—C13	-1.3 (2)
C9—S1—C8—C2	147.97 (11)	S1—C9—C14—C13	-179.58 (12)
C23—S2—C22—C21	102.41 (11)	S1—C9—C10—C11	178.36 (14)
C23—S2—C22—C20	-147.38 (11)	S1—C9—C14—N2	-2.55 (16)
C22—S2—C23—C28	9.92 (12)	C9—C10—C11—C12	0.7 (3)
C23—S2—C22—N4	-18.82 (10)	C10—C11—C12—C11	179.01 (15)
C22—S2—C23—C24	-172.01 (16)	C10—C11—C12—C13	-1.0 (3)
C1—N1—C7—O1	169.93 (15)	C11—C12—C13—C14	0.1 (3)
C1—N1—C7—C8	-10.37 (17)	C11—C12—C13—C14	-179.91 (13)
C7—N1—C1—C6	-174.55 (17)	C12—C13—C14—C9	1.0 (2)
C7—N1—C1—C2	4.51 (19)	C12—C13—C14—N2	-175.62 (15)
C8—N2—C14—C9	19.25 (18)	C16—C15—C20—C19	-1.5 (3)
C8—N2—C14—C13	-163.92 (14)	C16—C15—C20—C22	177.60 (16)
C14—N2—C8—S1	-25.03 (14)	N3—C15—C20—C19	177.41 (15)
C14—N2—C8—C2	-148.54 (13)	N3—C15—C20—C22	-3.54 (18)
C14—N2—C8—C7	91.06 (15)	C20—C15—C16—C17	1.0 (3)
C15—N3—C21—C22	10.02 (17)	N3—C15—C16—C17	-177.60 (18)
C21—N3—C15—C16	174.40 (17)	C15—C16—C17—C18	-0.3 (3)
C15—N3—C21—O2	-169.98 (16)	C16—C17—C18—C19	0.0 (3)
C21—N3—C15—C20	-4.37 (19)	C17—C18—C19—C20	-0.3 (3)
C22—N4—C28—C23	-18.74 (18)	C18—C19—C20—C15	1.1 (3)
C28—N4—C22—C20	147.82 (13)	C18—C19—C20—C22	-177.75 (17)
C22—N4—C28—C27	164.02 (14)	C15—C20—C22—S2	-104.54 (13)
C28—N4—C22—C21	-92.09 (15)	C15—C20—C22—C21	8.79 (16)
C28—N4—C22—S2	24.31 (14)	C19—C20—C22—S2	74.4 (2)
N1—C1—C6—C5	177.63 (18)	C15—C20—C22—N4	134.86 (14)
N1—C1—C2—C3	-177.35 (15)	C19—C20—C22—N4	-46.2 (2)
C2—C1—C6—C5	-1.3 (3)	C19—C20—C22—C21	-172.30 (17)
C6—C1—C2—C8	-177.18 (16)	O2—C21—C22—C20	168.65 (15)
N1—C1—C2—C8	3.69 (18)	O2—C21—C22—N4	39.7 (2)
C6—C1—C2—C3	1.8 (3)	N3—C21—C22—C20	-11.35 (16)
C1—C2—C8—S1	104.15 (13)	N3—C21—C22—S2	104.59 (12)
C3—C2—C8—N2	45.7 (2)	N3—C21—C22—N4	-140.30 (13)
C3—C2—C8—C7	172.06 (17)	O2—C21—C22—S2	-75.41 (17)
C1—C2—C8—N2	-135.45 (14)	C28—C23—C24—C25	0.0 (3)
C1—C2—C8—C7	-9.13 (16)	S2—C23—C28—N4	2.57 (16)
C3—C2—C8—S1	-74.7 (2)	C24—C23—C28—N4	-175.64 (15)
C1—C2—C3—C4	-0.8 (3)	C24—C23—C28—C27	1.8 (2)
C8—C2—C3—C4	177.86 (17)	S2—C23—C28—C27	180.00 (12)
C2—C3—C4—C5	-0.5 (3)	S2—C23—C24—C25	-177.89 (14)
C3—C4—C5—C6	0.9 (3)	C23—C24—C25—C26	-1.5 (3)
C4—C5—C6—C1	0.0 (3)	C24—C25—C26—C27	1.2 (3)
N1—C7—C8—C2	11.79 (16)	C24—C25—C26—C12	-179.04 (15)
N1—C7—C8—N2	140.99 (13)	C12—C26—C27—C28	-179.19 (13)

supplementary materials

N1—C7—C8—S1	-104.29 (12)	C25—C26—C27—C28	0.5 (3)
O1—C7—C8—C2	-168.49 (15)	C26—C27—C28—C23	-2.0 (2)
O1—C7—C8—N2	-39.3 (2)	C26—C27—C28—N4	175.09 (15)

Symmetry codes: (i) $x-1, -y+1/2, z+1/2$; (ii) $x+1, -y+1/2, z-1/2$; (iii) $-x+1, -y+1, -z$; (iv) $-x, y+1/2, -z+1/2$; (v) $-x+1, y-1/2, -z+1/2$; (vi) $x, y-1, z$; (vii) $-x, y-1/2, -z+1/2$; (viii) $x, y+1, z$; (ix) $-x+1, y+1/2, -z+1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1A \cdots O2 ^{vi}	0.90 (3)	1.95 (3)	2.840 (2)	171 (2)
N2—H2A \cdots S1 ^{vii}	0.90 (2)	2.61 (2)	3.506 (1)	177 (2)
N3—H3A \cdots O1 ^{viii}	0.89 (2)	1.99 (2)	2.867 (2)	166 (2)
N4—H4A \cdots S2 ^{ix}	0.89 (2)	2.63 (2)	3.511 (1)	176 (2)
C3—H3 \cdots O1 ^{iv}	0.93	2.53	3.418 (2)	161

Symmetry codes: (vi) $x, y-1, z$; (vii) $-x, y-1/2, -z+1/2$; (viii) $x, y+1, z$; (ix) $-x+1, y+1/2, -z+1/2$; (iv) $-x, y+1/2, -z+1/2$.

Fig. 1




Fig. 2

