

1-(1-Carboxymethyl-1,4-anhydro-2,3-O-isopropylidene- α -D-erythrofuranosyl)-thymine

G. M. J. Lenagh-Snow,^a S. F. Jenkinson,^{a*} A. J. Stewart,^b
G. W. J. Fleet^a and D. J. Watkin^c

^aDepartment of Organic Chemistry, Chemistry Research Laboratory, University of Oxford, Mansfield Road, Oxford OX1 3TA, England, ^bIdenix Pharmaceuticals, Inc., 60 Hampshire Street, Cambridge, MA 02139, USA, and ^cDepartment of Chemical Crystallography, Chemistry Research Laboratory, University of Oxford, Mansfield Road, Oxford OX1 3TA, England

Correspondence e-mail: sarah.jenkinson@chem.ox.ac.uk


Received 11 January 2010; accepted 13 January 2010

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}-\text{C}) = 0.006 \text{ \AA}$; R factor = 0.047; wR factor = 0.114; data-to-parameter ratio = 7.4.

X-Ray crystallography unequivocally determined the stereochemistry of the thymine base in the title compound, $C_{14}\text{H}_{18}\text{N}_2\text{O}_7$. The absolute stereochemistry was determined from the use of D-ribose as the starting material. There are two independent molecules in the asymmetric unit ($Z' = 2$) which exist as $\text{N}-\text{H} \cdots \text{O}$ hydrogen-bonded pairs in the crystal structure.

Related literature

The title compound was obtained during studies on the synthesis of the 5-carbon analogue of psicofuranine, a naturally occurring nucleoside. For related literature on psicofuranine, see: Schroeder & Hoeksema (1959); Smith *et al.* (1973); Garrett (1960). For anomeric bromination see: Probert *et al.* (2005); Smith *et al.* (1999). For the extinction correction, see: Larson (1970).



Experimental

Crystal data

$C_{14}\text{H}_{18}\text{N}_2\text{O}_7$

$M_r = 326.31$

Monoclinic, $P2_1$
 $a = 7.8937 (5) \text{ \AA}$
 $b = 13.3471 (10) \text{ \AA}$
 $c = 14.9208 (10) \text{ \AA}$
 $\beta = 103.565 (4)^\circ$
 $V = 1528.17 (18) \text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.12 \text{ mm}^{-1}$
 $T = 150 \text{ K}$
 $0.40 \times 0.20 \times 0.03 \text{ mm}$

Data collection

Nonius KappaCCD diffractometer
Absorption correction: multi-scan
(DENZO/SCALEPACK;
Otwinowski & Minor, 1997)
 $T_{\min} = 0.83$, $T_{\max} = 1.00$

9120 measured reflections
3090 independent reflections
2453 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.065$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.114$
 $S = 0.95$
3090 reflections
416 parameters

1 restraint
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.34 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.31 \text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
N26—H261 \cdots O1	0.88	1.93	2.791 (6)	164
N3—H31 \cdots O24	0.88	2.01	2.863 (6)	165

Data collection: COLLECT (Nonius, 2001); cell refinement: DENZO/SCALEPACK (Otwinowski & Minor, 1997); data reduction: DENZO/SCALEPACK; program(s) used to solve structure: SIR92 (Altomare *et al.*, 1994); program(s) used to refine structure: CRYSTALS (Betteridge *et al.*, 2003); molecular graphics: CAMERON (Watkin *et al.*, 1996); software used to prepare material for publication: CRYSTALS.

We would like to thank the Chemical Crystallography department and ALT at Oxford University for use of the diffractometers.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2979).

References

- Altomare, A., Casciaro, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
- Betteridge, P. W., Carruthers, J. R., Cooper, R. I., Prout, K. & Watkin, D. J. (2003). *J. Appl. Cryst.* **36**, 1487.
- Garrett, E. R. (1960). *J. Am. Chem. Soc.* **82**, 827–832.
- Larson, A. C. (1970). *Crystallographic Computing*, edited by F. R. Ahmed, S. R. Hall & C. P. Huber, pp. 291–294. Copenhagen: Munksgaard.
- Nonius (2001). COLLECT. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Probert, M. R., Watkin, D. J., Stewart, A. J., Storer, R. & Fleet, G. W. J. (2005). *Acta Cryst.* **E61**, o1718–o1720.
- Schroeder, W. & Hoeksema, H. (1959). *J. Am. Chem. Soc.* **81**, 1767–1768.
- Smith, M. D., Long, D. D., Martín, A., Campbell, N., Blériot, Y. & Fleet, G. W. J. (1999). *Synlett*, **7**, 1151–1154.
- Smith, C. G., Poutsiaka, J. W. & Schreiber, E. C. (1973). *J. Int. Med. Res.* **1**, 489–503.
- Watkin, D. J., Prout, C. K. & Pearce, L. J. (1996). CAMERON. Chemical Crystallography Laboratory, Oxford, England.

supplementary materials

Acta Cryst. (2010). E66, o404 [doi:10.1107/S1600536810001704]

1-(1-Carboxymethyl-1,4-anhydro-2,3-O-isopropylidene- α -D-erythrofuranosyl)thymine

G. M. J. Lenagh-Snow, S. F. Jenkinson, A. J. Stewart, G. W. J. Fleet and D. J. Watkin

Comment

Nucleosides are a powerful class of anti-viral and anti-bacterial agents. Psicofuranine **1** (Fig. 1) is a naturally occurring nucleoside with a branch at the anomeric position of the sugar (Schroeder & Hoeksema, 1959). It has potent anti-bacterial and anti-tumour activity but is cardiotoxic in man (Smith *et al.*, 1973). Psicofuranine is also unstable in acidic and basic conditions with the *N*-glycosidic bond readily undergoing hydrolytic cleavage (Garrett, 1960). During studies on the synthesis of the 5-carbon analogue of psicofuranine **2** the ester **4** was synthesized. Anomeric radical bromination (Smith *et al.*, 1999) gave rise to a single isolable bromide **5** (Probert *et al.*, 2005) which on displacement with silylated thymine gave a single nucleoside product. The stereochemistry at the anomeric position of the sugar was firmly established by X-ray crystallography and the structure was confirmed as **6** in which the thymine is in the α rather than the desired β position.

There are two crystallographically distinct molecules in the asymmetric unit which are related by a pseudo 2-fold rotation axis (Fig 2). When the two molecules are mapped they show good overlap (Fig. 3) with RMS deviations of 0.1055 on the positions, 0.082 for the bonds and 3.8892 for the torsion angles. These two molecules form hydrogen bonded pairs in the crystal structure (Fig. 4, Fig. 5). In both cases the central nitrogen (N3, N26) between the two carbonyls of the thymine acts as the donor but hydrogen bonds are formed to different carbonyls of the two thymine rings. The absolute stereochemistry was determined from the use of D-ribose as the starting material. Only classical hydrogen bonding was considered.

Experimental

The title compound was recrystallized by diffusion from a mixture of methanol and acetone: m.p. 457–458 K; $[\alpha]_D^{25}$ -235.2 (*c*, 0.84 in CHCl₃).

Refinement

In the absence of significant anomalous scattering, Friedel pairs were merged and the absolute configuration was assigned from the use of D-ribose as the starting material.

The H atoms were all located in a difference map, but those attached to carbon atoms were repositioned geometrically. The H atoms were initially refined with soft restraints on the bond lengths and angles to regularize their geometry (C—H in the range 0.93–0.98, N—H in the range 0.86–0.89 N—H to 0.86 O—H = 0.82 Å) and $U_{\text{iso}}(\text{H})$ (in the range 1.2–1.5 times U_{eq} of the parent atom), after which the positions were refined with riding constraints.

supplementary materials

Figures



Fig. 1. Synthetic Scheme




Fig. 2. The asymmetric unit of the title compound with displacement ellipsoids drawn at the 50% probability level. H atoms are shown as spheres of arbitrary radius.




Fig. 3. Overlay of the two molecules in the asymmetric unit.




Fig. 4. Hydrogen bonded dimer repeating unit. Hydrogen bonds are shown by dotted lines.




Fig. 5. Packing diagram projected along the a -axis. Hydrogen bonds are shown by dotted lines.

1-(1-Carboxymethyl-1,4-anhydro-2,3-O-isopropylidene- α -D- erythofuranosyl)thymine

Crystal data

$C_{14}H_{18}N_2O_7$

$F(000) = 688$

$M_r = 326.31$

$D_x = 1.418 \text{ Mg m}^{-3}$

Monoclinic, $P2_1$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Hall symbol: P 2yb

Cell parameters from 2763 reflections

$a = 7.8937 (5) \text{ \AA}$

$\theta = 5\text{--}26^\circ$

$b = 13.3471 (10) \text{ \AA}$

$\mu = 0.12 \text{ mm}^{-1}$

$c = 14.9208 (10) \text{ \AA}$

$T = 150 \text{ K}$

$\beta = 103.565 (4)^\circ$ Plate, colourless
 $V = 1528.17 (18) \text{ \AA}^3$ $0.40 \times 0.20 \times 0.03 \text{ mm}$
 $Z = 4$

Data collection

Nonius KappaCCD diffractometer 2453 reflections with $I > 2\sigma(I)$
graphite $R_{\text{int}} = 0.065$
 ω scans $\theta_{\text{max}} = 26.1^\circ, \theta_{\text{min}} = 5.2^\circ$
Absorption correction: multi-scan (DENZO/SCALEPACK; Otwinowski & Minor, 1997) $h = -9 \rightarrow 9$
 $T_{\text{min}} = 0.83, T_{\text{max}} = 1.00$ $k = -14 \rightarrow 16$
9120 measured reflections $l = -18 \rightarrow 18$
3090 independent reflections

Refinement

Refinement on F^2 Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full H-atom parameters constrained
 $R[F^2 > 2\sigma(F^2)] = 0.047$ Method = Modified Sheldrick $w = 1/\sigma^2(F^2) + (0.04P)^2 + 0.5P]$, where $P = (\max(F_o^2, 0) + 2F_c^2)/3$
 $wR(F^2) = 0.114$ $(\Delta/\sigma)_{\text{max}} = 0.0002$
 $S = 0.95$ $\Delta\rho_{\text{max}} = 0.34 \text{ e \AA}^{-3}$
3090 reflections $\Delta\rho_{\text{min}} = -0.31 \text{ e \AA}^{-3}$
416 parameters Extinction correction: Larson (1970), Equation 22
1 restraint Extinction coefficient: 590 (70)
Primary atom site location: structure-invariant direct methods

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.2931 (4)	0.3816 (2)	0.81712 (19)	0.0363
C2	0.3274 (5)	0.3347 (3)	0.8903 (3)	0.0308
N3	0.3044 (5)	0.3728 (3)	0.9715 (2)	0.0346
C4	0.3343 (6)	0.3220 (3)	1.0559 (3)	0.0348
O5	0.3097 (4)	0.3655 (2)	1.1245 (2)	0.0414
C6	0.3964 (5)	0.2203 (3)	1.0536 (3)	0.0336
C7	0.4220 (5)	0.1836 (3)	0.9739 (3)	0.0321
N8	0.3912 (4)	0.2388 (2)	0.8940 (2)	0.0290
C9	0.4110 (5)	0.1982 (3)	0.8058 (3)	0.0292
O10	0.5057 (4)	0.10739 (19)	0.82495 (18)	0.0320
C11	0.4357 (5)	0.0365 (3)	0.7525 (3)	0.0348
C12	0.2451 (5)	0.0597 (3)	0.7235 (3)	0.0343

supplementary materials

C13	0.2338 (5)	0.1728 (3)	0.7383 (3)	0.0320
O14	0.0903 (4)	0.1860 (2)	0.7798 (2)	0.0374
C15	0.0163 (5)	0.0881 (3)	0.7891 (3)	0.0356
O16	0.1544 (4)	0.0203 (2)	0.7876 (2)	0.0362
C17	-0.1363 (6)	0.0703 (3)	0.7081 (3)	0.0455
C18	-0.0325 (6)	0.0826 (4)	0.8799 (3)	0.0469
C19	0.5238 (5)	0.2679 (3)	0.7618 (3)	0.0324
O20	0.5064 (4)	0.2774 (2)	0.6798 (2)	0.0417
O21	0.6543 (4)	0.3076 (2)	0.8269 (2)	0.0380
C22	0.7785 (6)	0.3686 (4)	0.7933 (3)	0.0443
C23	0.4312 (7)	0.1600 (4)	1.1418 (3)	0.0465
O24	0.2166 (4)	0.5799 (2)	0.93916 (19)	0.0374
C25	0.1273 (5)	0.6136 (3)	0.8657 (3)	0.0313
N26	0.0825 (5)	0.5521 (2)	0.7891 (2)	0.0323
C27	-0.0261 (5)	0.5758 (3)	0.7053 (3)	0.0310
O28	-0.0626 (4)	0.5168 (2)	0.64081 (19)	0.0375
N29	-0.0943 (5)	0.6715 (2)	0.6997 (2)	0.0306
C30	-0.0454 (5)	0.7396 (3)	0.7711 (3)	0.0326
C31	0.0618 (5)	0.7156 (3)	0.8524 (3)	0.0305
C32	0.1154 (7)	0.7889 (3)	0.9308 (3)	0.0461
C33	-0.2093 (5)	0.6978 (3)	0.6101 (3)	0.0305
O34	-0.2864 (4)	0.7906 (2)	0.62183 (18)	0.0339
C35	-0.3007 (6)	0.8493 (3)	0.5385 (3)	0.0351
C36	-0.1471 (5)	0.8204 (3)	0.5013 (3)	0.0333
C37	-0.1111 (5)	0.7103 (3)	0.5318 (3)	0.0339
O38	0.0732 (4)	0.7037 (2)	0.5655 (2)	0.0364
C39	0.1460 (6)	0.7998 (3)	0.5517 (3)	0.0354
O40	0.0075 (4)	0.8694 (2)	0.5505 (2)	0.0368
C41	0.2020 (6)	0.8002 (4)	0.4618 (3)	0.0415
C42	0.2914 (6)	0.8224 (4)	0.6342 (3)	0.0473
C43	-0.3599 (6)	0.6221 (3)	0.5830 (3)	0.0347
O44	-0.4208 (4)	0.5954 (2)	0.65563 (19)	0.0351
C45	-0.5767 (6)	0.5327 (3)	0.6364 (3)	0.0430
O46	-0.4246 (4)	0.5982 (2)	0.5041 (2)	0.0448
H71	0.4624	0.1181	0.9722	0.0371*
H111	0.4513	-0.0316	0.7764	0.0408*
H112	0.4950	0.0448	0.7026	0.0412*
H121	0.1912	0.0372	0.6595	0.0412*
H131	0.2163	0.2111	0.6801	0.0380*
H172	-0.1903	0.0060	0.7168	0.0643*
H173	-0.0963	0.0702	0.6514	0.0642*
H171	-0.2232	0.1235	0.7066	0.0639*
H181	-0.0780	0.0161	0.8871	0.0748*
H183	0.0704	0.0951	0.9291	0.0752*
H182	-0.1213	0.1339	0.8809	0.0750*
H222	0.8569	0.3999	0.8455	0.0703*
H221	0.8444	0.3287	0.7598	0.0705*
H223	0.7183	0.4210	0.7528	0.0699*
H232	0.4644	0.0921	1.1292	0.0681*

H233	0.3276	0.1583	1.1672	0.0683*
H231	0.5292	0.1900	1.1863	0.0683*
H301	-0.0922	0.8051	0.7627	0.0356*
H323	0.0689	0.8550	0.9114	0.0654*
H322	0.2425	0.7918	0.9490	0.0654*
H321	0.0697	0.7671	0.9830	0.0652*
H351	-0.4091	0.8324	0.4936	0.0382*
H352	-0.2992	0.9212	0.5538	0.0381*
H361	-0.1674	0.8283	0.4335	0.0381*
H371	-0.1535	0.6632	0.4807	0.0422*
H412	0.2448	0.8668	0.4522	0.0641*
H411	0.2934	0.7499	0.4646	0.0643*
H413	0.1026	0.7836	0.4127	0.0639*
H421	0.3481	0.8846	0.6256	0.0668*
H422	0.3738	0.7673	0.6428	0.0670*
H423	0.2421	0.8276	0.6879	0.0672*
H453	-0.6176	0.5238	0.6929	0.0613*
H452	-0.5496	0.4674	0.6141	0.0609*
H451	-0.6673	0.5654	0.5897	0.0615*
H261	0.1302	0.4919	0.7947	0.0383*
H31	0.2694	0.4353	0.9707	0.0413*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0473 (17)	0.0277 (14)	0.0351 (16)	0.0081 (13)	0.0120 (14)	0.0074 (13)
C2	0.034 (2)	0.0255 (19)	0.033 (2)	0.0003 (17)	0.0092 (18)	0.0010 (18)
N3	0.0444 (19)	0.0259 (15)	0.0338 (19)	0.0032 (16)	0.0095 (16)	-0.0002 (15)
C4	0.034 (2)	0.036 (2)	0.035 (2)	-0.0036 (19)	0.0083 (19)	-0.0014 (19)
O5	0.0476 (18)	0.0436 (17)	0.0347 (16)	-0.0022 (16)	0.0128 (14)	-0.0064 (15)
C6	0.036 (2)	0.035 (2)	0.030 (2)	-0.0005 (19)	0.0074 (18)	0.0005 (18)
C7	0.031 (2)	0.032 (2)	0.032 (2)	0.0015 (17)	0.0062 (18)	0.0057 (17)
N8	0.0345 (17)	0.0266 (16)	0.0267 (17)	0.0057 (14)	0.0090 (14)	0.0010 (13)
C9	0.034 (2)	0.0221 (18)	0.031 (2)	0.0012 (17)	0.0079 (18)	0.0014 (16)
O10	0.0344 (14)	0.0229 (13)	0.0370 (15)	0.0019 (12)	0.0052 (12)	-0.0029 (12)
C11	0.036 (2)	0.029 (2)	0.040 (2)	-0.0002 (19)	0.0102 (19)	-0.0074 (19)
C12	0.035 (2)	0.031 (2)	0.037 (2)	0.0000 (18)	0.0094 (19)	-0.0027 (18)
C13	0.037 (2)	0.028 (2)	0.031 (2)	0.0029 (18)	0.0078 (18)	0.0004 (17)
O14	0.0329 (16)	0.0322 (15)	0.0482 (18)	0.0012 (13)	0.0122 (14)	-0.0035 (13)
C15	0.032 (2)	0.0252 (19)	0.048 (3)	-0.0016 (18)	0.0082 (19)	-0.0045 (19)
O16	0.0344 (16)	0.0272 (14)	0.0482 (18)	0.0020 (13)	0.0122 (14)	0.0032 (13)
C17	0.033 (2)	0.043 (3)	0.059 (3)	-0.002 (2)	0.008 (2)	-0.007 (2)
C18	0.044 (3)	0.043 (2)	0.057 (3)	0.001 (2)	0.018 (2)	0.001 (2)
C19	0.032 (2)	0.0254 (19)	0.041 (2)	0.0035 (18)	0.010 (2)	0.0018 (18)
O20	0.0489 (19)	0.0462 (18)	0.0318 (16)	-0.0022 (15)	0.0131 (15)	0.0039 (14)
O21	0.0370 (16)	0.0353 (15)	0.0405 (16)	-0.0056 (14)	0.0070 (14)	0.0041 (13)
C22	0.034 (2)	0.041 (2)	0.059 (3)	-0.009 (2)	0.013 (2)	0.010 (2)
C23	0.055 (3)	0.049 (3)	0.036 (2)	0.009 (2)	0.011 (2)	0.008 (2)

supplementary materials

O24	0.0423 (17)	0.0327 (15)	0.0333 (15)	0.0038 (14)	0.0012 (14)	0.0006 (13)
C25	0.029 (2)	0.031 (2)	0.033 (2)	-0.0001 (18)	0.0057 (18)	0.0009 (18)
N26	0.0386 (19)	0.0253 (16)	0.0315 (19)	0.0029 (15)	0.0055 (16)	-0.0003 (14)
C27	0.036 (2)	0.0234 (18)	0.034 (2)	-0.0007 (17)	0.0077 (18)	-0.0008 (17)
O28	0.0499 (18)	0.0265 (14)	0.0350 (16)	0.0015 (14)	0.0078 (14)	-0.0022 (13)
N29	0.0365 (19)	0.0266 (16)	0.0277 (17)	0.0020 (15)	0.0054 (15)	-0.0005 (14)
C30	0.037 (2)	0.0256 (19)	0.034 (2)	0.0014 (17)	0.0056 (19)	-0.0034 (17)
C31	0.035 (2)	0.0270 (19)	0.029 (2)	0.0006 (18)	0.0081 (18)	-0.0037 (17)
C32	0.056 (3)	0.033 (2)	0.043 (3)	0.004 (2)	-0.002 (2)	-0.007 (2)
C33	0.038 (2)	0.0236 (18)	0.030 (2)	0.0053 (17)	0.0075 (18)	0.0003 (16)
O34	0.0431 (17)	0.0279 (14)	0.0313 (14)	0.0063 (13)	0.0101 (14)	0.0035 (12)
C35	0.041 (2)	0.029 (2)	0.033 (2)	0.0036 (19)	0.0027 (19)	0.0061 (17)
C36	0.035 (2)	0.030 (2)	0.031 (2)	0.0022 (18)	0.0020 (18)	0.0027 (17)
C37	0.040 (2)	0.0294 (19)	0.032 (2)	-0.0002 (19)	0.0082 (19)	-0.0040 (18)
O38	0.0371 (16)	0.0274 (14)	0.0451 (17)	0.0044 (13)	0.0107 (14)	0.0032 (13)
C39	0.043 (3)	0.0258 (19)	0.039 (2)	0.0027 (19)	0.013 (2)	0.0005 (18)
O40	0.0356 (15)	0.0291 (13)	0.0456 (17)	0.0016 (13)	0.0092 (14)	-0.0033 (13)
C41	0.041 (3)	0.046 (2)	0.040 (2)	-0.002 (2)	0.016 (2)	0.002 (2)
C42	0.041 (3)	0.050 (3)	0.048 (3)	0.004 (2)	0.007 (2)	-0.009 (2)
C43	0.038 (2)	0.030 (2)	0.036 (2)	0.0048 (19)	0.008 (2)	-0.0004 (18)
O44	0.0372 (16)	0.0325 (15)	0.0353 (15)	-0.0068 (13)	0.0081 (13)	-0.0031 (13)
C45	0.045 (3)	0.033 (2)	0.051 (3)	-0.011 (2)	0.012 (2)	-0.001 (2)
O46	0.0479 (18)	0.0505 (18)	0.0323 (15)	-0.0102 (16)	0.0022 (14)	-0.0051 (16)

Geometric parameters (Å, °)

O1—C2	1.233 (5)	O24—C25	1.240 (5)
C2—N3	1.365 (5)	C25—N26	1.384 (5)
C2—N8	1.371 (5)	C25—C31	1.453 (5)
N3—C4	1.401 (5)	N26—C27	1.377 (5)
N3—H31	0.877	N26—H261	0.882
C4—O5	1.230 (5)	C27—O28	1.224 (5)
C4—C6	1.448 (6)	C27—N29	1.380 (5)
C6—C7	1.345 (6)	N29—C30	1.385 (5)
C6—C23	1.512 (6)	N29—C33	1.471 (5)
C7—N8	1.374 (5)	C30—C31	1.346 (5)
C7—H71	0.933	C30—H301	0.946
N8—C9	1.464 (5)	C31—C32	1.507 (6)
C9—O10	1.418 (5)	C32—H323	0.973
C9—C13	1.557 (6)	C32—H322	0.977
C9—C19	1.537 (5)	C32—H321	0.977
O10—C11	1.445 (5)	C33—O34	1.408 (5)
C11—C12	1.497 (6)	C33—C37	1.555 (5)
C11—H111	0.974	C33—C43	1.541 (6)
C11—H112	0.974	O34—C35	1.452 (5)
C12—C13	1.532 (5)	C35—C36	1.497 (6)
C12—O16	1.423 (5)	C35—H351	0.981
C12—H121	0.996	C35—H352	0.986
C13—O14	1.424 (5)	C36—C37	1.544 (6)

C13—H131	0.988	C36—O40	1.427 (5)
O14—C15	1.452 (5)	C36—H361	0.992
C15—O16	1.420 (5)	C37—O38	1.426 (5)
C15—C17	1.512 (6)	C37—H371	0.985
C15—C18	1.496 (6)	O38—C39	1.440 (5)
C17—H172	0.981	C39—O40	1.431 (5)
C17—H173	0.969	C39—C41	1.507 (6)
C17—H171	0.984	C39—C42	1.504 (6)
C18—H181	0.974	C41—H412	0.974
C18—H183	0.973	C41—H411	0.979
C18—H182	0.982	C41—H413	0.965
C19—O20	1.205 (5)	C42—H421	0.966
C19—O21	1.348 (5)	C42—H422	0.971
O21—C22	1.451 (5)	C42—H423	0.972
C22—H222	0.969	C43—O44	1.333 (5)
C22—H221	0.962	C43—O46	1.210 (5)
C22—H223	0.972	O44—C45	1.460 (5)
C23—H232	0.974	C45—H453	0.979
C23—H233	0.979	C45—H452	0.975
C23—H231	0.979	C45—H451	0.977
O1—C2—N3	123.3 (4)	O24—C25—N26	119.7 (4)
O1—C2—N8	120.7 (3)	O24—C25—C31	124.8 (4)
N3—C2—N8	115.9 (3)	N26—C25—C31	115.5 (3)
C2—N3—C4	126.0 (3)	C25—N26—C27	126.7 (3)
C2—N3—H31	116.7	C25—N26—H261	116.2
C4—N3—H31	117.3	C27—N26—H261	117.2
N3—C4—O5	119.6 (4)	N26—C27—O28	123.2 (3)
N3—C4—C6	114.8 (4)	N26—C27—N29	114.8 (3)
O5—C4—C6	125.6 (4)	O28—C27—N29	122.0 (4)
C4—C6—C7	119.0 (4)	C27—N29—C30	121.7 (3)
C4—C6—C23	118.2 (4)	C27—N29—C33	115.2 (3)
C7—C6—C23	122.8 (4)	C30—N29—C33	122.9 (3)
C6—C7—N8	122.6 (4)	N29—C30—C31	122.7 (4)
C6—C7—H71	119.2	N29—C30—H301	118.4
N8—C7—H71	118.2	C31—C30—H301	118.9
C7—N8—C2	121.6 (3)	C25—C31—C30	118.3 (4)
C7—N8—C9	123.1 (3)	C25—C31—C32	118.5 (4)
C2—N8—C9	115.2 (3)	C30—C31—C32	123.2 (4)
N8—C9—O10	107.4 (3)	C31—C32—H323	110.0
N8—C9—C13	113.1 (3)	C31—C32—H322	108.9
O10—C9—C13	107.2 (3)	H323—C32—H322	109.8
N8—C9—C19	110.8 (3)	C31—C32—H321	109.6
O10—C9—C19	105.8 (3)	H323—C32—H321	108.8
C13—C9—C19	112.2 (3)	H322—C32—H321	109.7
C9—O10—C11	108.5 (3)	N29—C33—O34	106.9 (3)
O10—C11—C12	105.3 (3)	N29—C33—C37	113.4 (3)
O10—C11—H111	110.1	O34—C33—C37	107.8 (3)
C12—C11—H111	109.3	N29—C33—C43	110.8 (3)
O10—C11—H112	109.2	O34—C33—C43	106.3 (3)

supplementary materials

C12—C11—H112	112.7	C37—C33—C43	111.2 (3)
H111—C11—H112	110.1	C33—O34—C35	108.5 (3)
C11—C12—C13	104.5 (3)	O34—C35—C36	105.7 (3)
C11—C12—O16	111.1 (3)	O34—C35—H351	109.9
C13—C12—O16	102.2 (3)	C36—C35—H351	109.9
C11—C12—H121	112.9	O34—C35—H352	109.4
C13—C12—H121	114.1	C36—C35—H352	111.7
O16—C12—H121	111.4	H351—C35—H352	110.1
C12—C13—C9	103.5 (3)	C35—C36—C37	104.2 (3)
C12—C13—O14	105.3 (3)	C35—C36—O40	111.1 (3)
C9—C13—O14	112.2 (3)	C37—C36—O40	102.1 (3)
C12—C13—H131	112.6	C35—C36—H361	113.5
C9—C13—H131	112.0	C37—C36—H361	112.1
O14—C13—H131	110.9	O40—C36—H361	112.9
C13—O14—C15	108.0 (3)	C36—C37—C33	103.5 (3)
O14—C15—O16	104.2 (3)	C36—C37—O38	105.3 (3)
O14—C15—C17	109.1 (3)	C33—C37—O38	112.1 (3)
O16—C15—C17	111.0 (3)	C36—C37—H371	111.9
O14—C15—C18	109.0 (3)	C33—C37—H371	112.0
O16—C15—C18	110.2 (4)	O38—C37—H371	111.5
C17—C15—C18	113.0 (4)	C37—O38—C39	107.7 (3)
C12—O16—C15	106.6 (3)	O38—C39—O40	104.5 (3)
C15—C17—H172	108.7	O38—C39—C41	109.9 (3)
C15—C17—H173	109.6	O40—C39—C41	111.7 (3)
H172—C17—H173	110.8	O38—C39—C42	108.4 (4)
C15—C17—H171	109.0	O40—C39—C42	108.3 (3)
H172—C17—H171	108.2	C41—C39—C42	113.7 (4)
H173—C17—H171	110.5	C39—O40—C36	105.5 (3)
C15—C18—H181	108.9	C39—C41—H412	108.6
C15—C18—H183	109.1	C39—C41—H411	109.1
H181—C18—H183	109.7	H412—C41—H411	110.7
C15—C18—H182	108.4	C39—C41—H413	108.6
H181—C18—H182	110.6	H412—C41—H413	110.2
H183—C18—H182	110.0	H411—C41—H413	109.6
C9—C19—O20	123.9 (4)	C39—C42—H421	110.6
C9—C19—O21	110.5 (3)	C39—C42—H422	108.4
O20—C19—O21	125.2 (4)	H421—C42—H422	110.6
C19—O21—C22	115.8 (3)	C39—C42—H423	108.2
O21—C22—H222	108.7	H421—C42—H423	109.6
O21—C22—H221	111.1	H422—C42—H423	109.4
H222—C22—H221	109.6	C33—C43—O44	111.4 (3)
O21—C22—H223	110.3	C33—C43—O46	123.5 (4)
H222—C22—H223	108.4	O44—C43—O46	124.8 (4)
H221—C22—H223	108.7	C43—O44—C45	116.2 (3)
C6—C23—H232	109.2	O44—C45—H453	109.5
C6—C23—H233	110.6	O44—C45—H452	109.7
H232—C23—H233	109.9	H453—C45—H452	109.2
C6—C23—H231	109.0	O44—C45—H451	109.1
H232—C23—H231	107.9	H453—C45—H451	109.6

supplementary materials

H233—C23—H231

110.3

H452—C45—H451

109.8

Hydrogen-bond geometry (Å, °)

D—H···*A*

D—H

H···*A*

D···*A*

D—H···*A*

C11—H111···O5ⁱ

0.97

2.52

3.301 (6)

138

N26—H261···O1

0.88

1.93

2.791 (6)

164

N3—H31···O24

0.88

2.01

2.863 (6)

165

Symmetry codes: (i) $-x+1, y-1/2, -z+2$.

supplementary materials

Fig. 1





Fig. 2



supplementary materials

Fig. 3





Fig. 4



supplementary materials

Fig. 5

