

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

7-[4-(4-Fluorophenyl)-2-methylsulfanyl-1H-imidazol-5-yl]tetrazolo[1,5-a]pyridine

Roland Selig,^a Dieter Schollmeyer,^b Joachim Schlosser,^a Wolfgang Albrecht^c and Stefan Laufer^{a*}^aEberhard-Karls-University Tübingen, Auf der Morgenstelle 8, 72076 Tübingen, Germany, ^bUniversity Mainz, Duesbergweg 10-14, 55099 Mainz, Germany, and ^cc-a-i-r biosciences GmbH, Paul-Ehrlich-Strasse 15, 72076 Tübingen, Germany
Correspondence e-mail: stefan.laufer@uni-tuebingen.de


Received 14 January 2010; accepted 21 January 2010

Key indicators: single-crystal X-ray study; $T = 193$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.038; wR factor = 0.107; data-to-parameter ratio = 13.0.

The crystal structure of the title compound, $\text{C}_{15}\text{H}_{11}\text{FN}_6\text{S}$, forms a three-dimensional network stabilized by π - π interactions between the imidazole core and the tetrazole ring of the tetrazolopyridine unit; the centroid-centroid distance is 3.627 (1) Å. The crystal structure also displays bifurcated $\text{N}-\text{H}\cdots(\text{N},\text{N})$ hydrogen bonding and $\text{C}-\text{H}\cdots\text{F}$ interactions. The former involve the NH H atom of the imidazole core and the tetrazolopyridine N atoms, while the latter involve a methyl H atom, of the methylsulfanyl group, and the 4-fluorophenyl F atom. In the molecule, the imidazole ring makes dihedral angles of 40.45 (9) and 17.09 (8)°, respectively, with the 4-fluorophenyl ring and the tetrazolopyridine ring mean plane.

Related literature

For the biological relevance and the development of p38 MAP kinase inhibitors, see: Peifer *et al.* (2006). For the preparation of 2-fluoro-4-[4-(4-fluorophenyl)-2-(methylthio)-1H-imidazol-5-yl]pyridine, see: Laufer & Liedtke (2006). For the preparation of tetrazolopyridines, see: Capelli *et al.* (2008).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{11}\text{FN}_6\text{S}$
 $M_r = 326.36$
 Monoclinic, $P2_1/c$
 $a = 9.8342$ (7) Å
 $b = 18.1908$ (6) Å
 $c = 8.2374$ (7) Å
 $\beta = 100.292$ (3)°

$V = 1449.89$ (17) Å³
 $Z = 4$
 Cu $K\alpha$ radiation
 $\mu = 2.17$ mm⁻¹
 $T = 193$ K
 $0.30 \times 0.20 \times 0.10$ mm

Data collection

Enraf-Nonius CAD-4
 diffractometer
 Absorption correction: ψ scan
 (CORINC; Dräger & Gattow, 1971)
 $T_{\min} = 0.866$, $T_{\max} = 0.999$

5760 measured reflections
 2733 independent reflections
 2403 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.056$
 3 standard reflections every 60 min
 intensity decay: 3%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.107$
 $S = 1.06$
 2733 reflections

210 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.37$ e Å⁻³
 $\Delta\rho_{\min} = -0.30$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N10}-\text{H10}\cdots\text{N1}^i$	0.92	2.06	2.9703 (19)	174
$\text{N10}-\text{H10}\cdots\text{N2}^i$	0.92	2.60	3.423 (2)	151
$\text{C21}-\text{H21B}\cdots\text{F19}^{ii}$	0.98	2.44	3.286 (3)	144

Symmetry codes: (i) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x + 1, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *CAD-4 Software* (Enraf-Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *CORINC* (Dräger & Gattow, 1971); program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *PLATON*.

The authors would like to thank the Federal Ministry of Education and Research, Germany, Merckle GmbH, Ulm, Germany, and the Fonds der Chemischen Industrie, Germany, for their generous support of this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2158).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Capelli, A., Giuliani, G., Anzini, M., Riitano, D., Giorgi, G. & Vomero, S. (2008). *Bioorg. Med. Chem.* **16**, 6850–6859.
- Dräger, M. & Gattow, G. (1971). *Acta Chem. Scand.* **25**, 761–762.
- Enraf-Nonius (1989). *CAD-4 Software*. Enraf-Nonius, Delft, The Netherlands.
- Laufer, S. & Liedtke, A. (2006). *Tetrahedron Lett.* **47**, 7199–7203.
- Peifer, C., Wagner, G. & Laufer, S. (2006). *Curr. Top. Med. Chem.* **6**, 113–149.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2010). E66, o451 [doi:10.1107/S1600536810002680]

7-[4-(4-Fluorophenyl)-2-methylsulfanyl-1*H*-imidazol-5-yl]tetrazolo[1,5-*a*]pyridine

R. Selig, D. Schollmeyer, J. Schlosser, W. Albrecht and S. Laufer

Comment

Pyridylimidazoles like SB203580 are well known p38 MAP kinase inhibitors (Peifer *et al.*, 2006). The function of the pyridine moiety is to accept a hydrogen bond from the backbone of Met109 in the Hinge region. In the course of our studies we have tried to modify this acceptor system by using the title tetrazolopyridine (Capelli *et al.*, 2008).

The molecular structure of the title compound is given in Fig. 1, and the geometrical parameters are available in the Supplementary information and the archived CIF. The imidazole ring mean plane makes dihedral angles of 40.45 (9)° and 17.09 (8)° with the 4-fluorophenyl ring and the tetrazolopyridine ring mean plane, respectively.

The crystal structure displays asymmetric bifurcated N—H···N hydrogen bonds involving the tetrazolopyridine N-atoms and the NH H-atom of the the imidazole core (Table 1). There is also a C—H···F interaction involving the methylsulfanyl group and the 4-fluorophenyl F-atom (Table 1). The crystal structure of the title compound forms a three dimensional network stabilized by π - π interactions between the imidazole core and the tetrazole moiety of the tetrazolopyridine group; the centroid···centroid distance is 3.627 (1) Å (Table 1).

Experimental

A mixture of 300 mg 2-fluoro-4-[4-(4-fluorophenyl)-2-(methylthio)-1*H*-imidazol-5-yl]pyridine (Laufer & Liedtke, 2006) in anhydrous DMF with 100 mg sodium azide was heated at 353 K for 12 h. The solvent was then removed under reduced pressure and the residue was diluted with ethylacetate. The organic phase was washed with water and concentrated under reduced pressure. The residue was purified by flash chromatography with ethyl acetate/hexane (1/1) to yield 153 mg (47%) of the title compound. Crystals suitable for X-ray analysis were obtained by crystallization from methanol.

Refinement

The H atom attached to N10 was located in a difference Fourier map and refined with a distance restraint of 0.92 (2) Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$. The C-bound H-atoms were placed in calculated positions and refined in the riding-model approximation: C—H = 0.95 - 0.98 Å with $U_{\text{iso}}(\text{H}) = k \times U_{\text{eq}}(\text{C})$, where $k = 1.2$ for H-aromatic and 1.5 for H-methyl.

Figures




Fig. 1. A view of the molecular structure of the title compound. the displacement ellipsoids are drawn at the 50% probability level.

7-[4-(4-Fluorophenyl)-2-methylsulfonyl-1H-imidazol-5-yl]tetrazolo[1,5-a]pyridine

Crystal data

$C_{15}H_{11}FN_6S$

$M_r = 326.36$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 9.8342\ (7)\ \text{\AA}$

$b = 18.1908\ (6)\ \text{\AA}$

$c = 8.2374\ (7)\ \text{\AA}$

$\beta = 100.292\ (3)^\circ$

$V = 1449.89\ (17)\ \text{\AA}^3$

$Z = 4$

$F(000) = 672$

$D_x = 1.495\ \text{Mg m}^{-3}$

Cu $K\alpha$ radiation, $\lambda = 1.54178\ \text{\AA}$

Cell parameters from 25 reflections

$\theta = 65\text{--}70^\circ$

$\mu = 2.17\ \text{mm}^{-1}$

$T = 193\ \text{K}$

Plate, colourless

$0.30 \times 0.20 \times 0.10\ \text{mm}$

Data collection

Enraf–Nonius CAD-4
diffractometer

Radiation source: rotating anode
graphite

$\omega/2\theta$ scans

Absorption correction: ψ scan
(CORINC; Dräger & Gattow, 1971)

$T_{\min} = 0.866$, $T_{\max} = 0.999$

5760 measured reflections

2733 independent reflections

2403 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.056$

$\theta_{\max} = 69.9^\circ$, $\theta_{\min} = 4.6^\circ$

$h = -11 \rightarrow 11$

$k = -22 \rightarrow 22$

$l = -10 \rightarrow 0$

3 standard reflections every 60 min

intensity decay: 3%

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.038$

$wR(F^2) = 0.107$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.051P)^2 + 0.5315P]$

$S = 1.06$	where $P = (F_o^2 + 2F_c^2)/3$
2733 reflections	$(\Delta/\sigma)_{\max} = 0.001$
210 parameters	$\Delta\rho_{\max} = 0.37 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.30 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kFc[1+0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$ Extinction coefficient: 0.0014 (2)

Special details

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell esds are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S20	0.84423 (5)	0.32567 (2)	0.15709 (8)	0.0423 (2)
F19	0.09192 (12)	0.08344 (6)	0.37488 (16)	0.0442 (4)
N1	0.29691 (15)	0.61704 (7)	0.31439 (19)	0.0284 (4)
N2	0.18289 (15)	0.63452 (8)	0.3762 (2)	0.0315 (4)
N3	0.12953 (15)	0.57831 (8)	0.4381 (2)	0.0304 (4)
N3A	0.21213 (13)	0.52053 (7)	0.41594 (17)	0.0232 (4)
N8	0.62050 (14)	0.39105 (8)	0.25381 (18)	0.0265 (4)
N10	0.61319 (14)	0.26910 (7)	0.24463 (18)	0.0271 (4)
C4	0.20076 (17)	0.44913 (9)	0.4666 (2)	0.0267 (5)
C5	0.29443 (17)	0.40040 (9)	0.4318 (2)	0.0264 (5)
C6	0.40054 (16)	0.42075 (8)	0.3415 (2)	0.0221 (4)
C7	0.41195 (16)	0.49369 (9)	0.2991 (2)	0.0231 (4)
C7A	0.31553 (16)	0.54434 (9)	0.3388 (2)	0.0230 (4)
C9	0.68541 (17)	0.33087 (9)	0.2239 (2)	0.0275 (5)
C11	0.49088 (16)	0.29071 (9)	0.2895 (2)	0.0239 (5)
C12	0.49810 (16)	0.36671 (9)	0.2974 (2)	0.0233 (5)
C13	0.38534 (16)	0.23580 (9)	0.3073 (2)	0.0241 (5)
C14	0.24524 (17)	0.24835 (9)	0.2461 (2)	0.0286 (5)
C15	0.14573 (18)	0.19732 (10)	0.2673 (2)	0.0313 (5)
C16	0.18916 (19)	0.13307 (9)	0.3491 (2)	0.0301 (5)
C17	0.32598 (19)	0.11634 (9)	0.4043 (2)	0.0297 (5)
C18	0.42396 (18)	0.16837 (9)	0.3835 (2)	0.0266 (5)
C21	0.8749 (2)	0.42202 (11)	0.1351 (3)	0.0420 (7)
H4	0.12940	0.43470	0.52390	0.0320*
H5	0.29010	0.35100	0.46820	0.0320*

supplementary materials

H7	0.48390	0.50930	0.24390	0.0280*
H10	0.64590	0.22270	0.23220	0.0320*
H14	0.21790	0.29280	0.18890	0.0340*
H15	0.05050	0.20630	0.22690	0.0380*
H17	0.35240	0.07040	0.45520	0.0360*
H18	0.51900	0.15820	0.42160	0.0320*
H21A	0.80250	0.44270	0.05010	0.0630*
H21B	0.96520	0.42910	0.10280	0.0630*
H21C	0.87380	0.44690	0.24030	0.0630*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S20	0.0311 (3)	0.0261 (3)	0.0757 (4)	0.0036 (2)	0.0258 (2)	-0.0011 (2)
F19	0.0400 (6)	0.0307 (6)	0.0661 (8)	-0.0095 (5)	0.0210 (6)	0.0042 (5)
N1	0.0261 (7)	0.0188 (7)	0.0413 (8)	0.0027 (5)	0.0087 (6)	0.0029 (6)
N2	0.0264 (7)	0.0217 (7)	0.0478 (9)	0.0046 (6)	0.0104 (6)	0.0033 (6)
N3	0.0264 (7)	0.0230 (7)	0.0436 (9)	0.0066 (6)	0.0112 (6)	0.0017 (6)
N3A	0.0207 (6)	0.0193 (6)	0.0304 (7)	0.0018 (5)	0.0066 (5)	0.0006 (5)
N8	0.0218 (7)	0.0208 (7)	0.0377 (8)	0.0005 (5)	0.0073 (6)	-0.0022 (6)
N10	0.0239 (7)	0.0182 (6)	0.0395 (8)	0.0026 (5)	0.0067 (6)	-0.0020 (6)
C4	0.0273 (8)	0.0208 (8)	0.0342 (9)	-0.0014 (6)	0.0113 (7)	0.0034 (7)
C5	0.0304 (8)	0.0185 (8)	0.0311 (9)	-0.0003 (6)	0.0076 (7)	0.0027 (7)
C6	0.0209 (7)	0.0192 (7)	0.0255 (8)	0.0000 (6)	0.0019 (6)	-0.0016 (6)
C7	0.0202 (7)	0.0204 (7)	0.0293 (8)	-0.0014 (6)	0.0059 (6)	-0.0012 (6)
C7A	0.0211 (7)	0.0191 (7)	0.0286 (8)	-0.0018 (6)	0.0037 (6)	0.0000 (6)
C9	0.0219 (8)	0.0219 (8)	0.0390 (9)	0.0005 (6)	0.0062 (7)	-0.0014 (7)
C11	0.0228 (8)	0.0194 (8)	0.0287 (8)	0.0020 (6)	0.0027 (6)	-0.0015 (6)
C12	0.0221 (8)	0.0198 (8)	0.0275 (8)	0.0007 (6)	0.0031 (6)	-0.0004 (6)
C13	0.0265 (8)	0.0193 (8)	0.0262 (8)	0.0003 (6)	0.0043 (6)	-0.0028 (6)
C14	0.0283 (9)	0.0198 (8)	0.0364 (10)	0.0018 (7)	0.0022 (7)	0.0005 (7)
C15	0.0246 (8)	0.0256 (8)	0.0432 (10)	0.0009 (7)	0.0044 (7)	-0.0034 (8)
C16	0.0320 (9)	0.0227 (8)	0.0382 (9)	-0.0057 (7)	0.0133 (7)	-0.0041 (7)
C17	0.0389 (10)	0.0212 (8)	0.0299 (9)	0.0016 (7)	0.0088 (7)	0.0025 (7)
C18	0.0280 (8)	0.0222 (8)	0.0288 (8)	0.0027 (6)	0.0026 (7)	-0.0005 (6)
C21	0.0417 (11)	0.0295 (10)	0.0608 (13)	-0.0047 (8)	0.0255 (10)	-0.0009 (9)

Geometric parameters (\AA , $^\circ$)

S20—C9	1.7488 (18)	C11—C12	1.385 (2)
S20—C21	1.793 (2)	C11—C13	1.467 (2)
F19—C16	1.359 (2)	C13—C18	1.399 (2)
N1—N2	1.350 (2)	C13—C14	1.399 (2)
N1—C7A	1.345 (2)	C14—C15	1.382 (2)
N2—N3	1.295 (2)	C15—C16	1.378 (2)
N3—N3A	1.361 (2)	C16—C17	1.375 (3)
N3A—C4	1.375 (2)	C17—C18	1.383 (2)
N3A—C7A	1.362 (2)	C4—H4	0.9500
N8—C9	1.313 (2)	C5—H5	0.9500

N8—C12	1.389 (2)	C7—H7	0.9500
N10—C9	1.356 (2)	C14—H14	0.9500
N10—C11	1.377 (2)	C15—H15	0.9500
N10—H10	0.9200	C17—H17	0.9500
C4—C5	1.346 (2)	C18—H18	0.9500
C5—C6	1.434 (2)	C21—H21A	0.9800
C6—C7	1.382 (2)	C21—H21B	0.9800
C6—C12	1.464 (2)	C21—H21C	0.9800
C7—C7A	1.402 (2)		
C9—S20—C21	98.90 (9)	C11—C13—C14	121.46 (15)
N2—N1—C7A	105.98 (13)	C11—C13—C18	120.11 (15)
N1—N2—N3	112.64 (14)	C13—C14—C15	121.26 (15)
N2—N3—N3A	105.28 (14)	C14—C15—C16	117.82 (16)
N3—N3A—C4	127.32 (14)	F19—C16—C17	118.39 (15)
N3—N3A—C7A	109.28 (13)	F19—C16—C15	118.32 (16)
C4—N3A—C7A	123.36 (14)	C15—C16—C17	123.28 (17)
C9—N8—C12	104.82 (14)	C16—C17—C18	118.02 (15)
C9—N10—C11	107.42 (13)	C13—C18—C17	121.08 (16)
C11—N10—H10	129.00	N3A—C4—H4	121.00
C9—N10—H10	123.00	C5—C4—H4	121.00
N3A—C4—C5	117.50 (15)	C4—C5—H5	119.00
C4—C5—C6	121.98 (15)	C6—C5—H5	119.00
C5—C6—C7	118.59 (14)	C6—C7—H7	121.00
C5—C6—C12	121.68 (14)	C7A—C7—H7	121.00
C7—C6—C12	119.71 (14)	C13—C14—H14	119.00
C6—C7—C7A	118.90 (15)	C15—C14—H14	119.00
N3A—C7A—C7	119.48 (14)	C14—C15—H15	121.00
N1—C7A—N3A	106.81 (14)	C16—C15—H15	121.00
N1—C7A—C7	133.70 (15)	C16—C17—H17	121.00
S20—C9—N8	126.57 (13)	C18—C17—H17	121.00
S20—C9—N10	120.80 (12)	C13—C18—H18	119.00
N8—C9—N10	112.59 (15)	C17—C18—H18	119.00
N10—C11—C12	104.93 (14)	S20—C21—H21A	109.00
C12—C11—C13	134.93 (15)	S20—C21—H21B	109.00
N10—C11—C13	120.00 (14)	S20—C21—H21C	109.00
N8—C12—C11	110.22 (14)	H21A—C21—H21B	109.00
C6—C12—C11	130.65 (15)	H21A—C21—H21C	109.00
N8—C12—C6	119.13 (14)	H21B—C21—H21C	109.00
C14—C13—C18	118.41 (15)		
C21—S20—C9—N8	-3.16 (18)	C5—C6—C12—C11	18.8 (3)
C21—S20—C9—N10	174.43 (15)	C7—C6—C12—N8	16.8 (2)
N2—N1—C7A—C7	178.07 (18)	C7—C6—C12—C11	-162.78 (17)
N2—N1—C7A—N3A	-0.43 (18)	C5—C6—C12—N8	-161.63 (15)
C7A—N1—N2—N3	0.4 (2)	C12—C6—C7—C7A	178.46 (15)
N1—N2—N3—N3A	-0.1 (2)	C6—C7—C7A—N3A	-0.9 (2)
N2—N3—N3A—C7A	-0.15 (18)	C6—C7—C7A—N1	-179.22 (18)
N2—N3—N3A—C4	177.50 (16)	N10—C11—C12—N8	1.62 (18)
N3—N3A—C7A—N1	0.37 (18)	N10—C11—C12—C6	-178.74 (16)

supplementary materials

N3—N3A—C4—C5	-179.88 (15)	C13—C11—C12—N8	-173.91 (17)
C7A—N3A—C4—C5	-2.5 (2)	C13—C11—C12—C6	5.7 (3)
C4—N3A—C7A—C7	3.9 (2)	N10—C11—C13—C14	-136.99 (17)
N3—N3A—C7A—C7	-178.39 (15)	N10—C11—C13—C18	41.5 (2)
C4—N3A—C7A—N1	-177.39 (15)	C12—C11—C13—C14	38.0 (3)
C12—N8—C9—S20	177.95 (13)	C12—C11—C13—C18	-143.46 (19)
C9—N8—C12—C11	-1.15 (18)	C11—C13—C14—C15	-178.13 (15)
C12—N8—C9—N10	0.20 (19)	C18—C13—C14—C15	3.3 (2)
C9—N8—C12—C6	179.17 (15)	C11—C13—C18—C17	178.80 (15)
C9—N10—C11—C12	-1.45 (18)	C14—C13—C18—C17	-2.6 (2)
C9—N10—C11—C13	174.90 (15)	C13—C14—C15—C16	-0.8 (2)
C11—N10—C9—S20	-177.08 (12)	C14—C15—C16—F19	178.12 (15)
C11—N10—C9—N8	0.82 (19)	C14—C15—C16—C17	-2.6 (3)
N3A—C4—C5—C6	-1.7 (2)	F19—C16—C17—C18	-177.45 (15)
C4—C5—C6—C7	4.4 (2)	C15—C16—C17—C18	3.3 (3)
C4—C5—C6—C12	-177.10 (16)	C16—C17—C18—C13	-0.6 (2)
C5—C6—C7—C7A	-3.0 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N10—H10 \cdots N1 ⁱ	0.92	2.06	2.9703 (19)	174
N10—H10 \cdots N2 ⁱ	0.92	2.60	3.423 (2)	151
C7—H7 \cdots N8	0.95	2.53	2.847 (2)	100
C21—H21B \cdots F19 ⁱⁱ	0.98	2.44	3.286 (3)	144

Symmetry codes: (i) $-x+1, y-1/2, -z+1/2$; (ii) $x+1, -y+1/2, z-1/2$.

Fig. 1

