

Dichloridobis[5-nitro-1-trimethylsilylmethyl-1*H*-benzimidazole- κ N³]cobalt(II) *N,N*-dimethylformamide solvate

Mehmet Akkurt,^{a*} Şerife Pınar Yalçın,^b Nihat Şireci,^c Hasan Küçükbay^c and M. Nawaz Tahir^d

^aDepartment of Physics, Faculty of Arts and Sciences, Erciyes University, 38039 Kayseri, Turkey, ^bDepartment of Physics, Faculty of Arts and Sciences, Harran University, 63300 Şanlıurfa, Turkey, ^cDepartment of Chemistry, Faculty of Arts and Sciences, İnönü University, 44280 Malatya, Turkey, and ^dDepartment of Physics, University of Sargodha, Sargodha, Pakistan

Correspondence e-mail: akkurt@erciyes.edu.tr


Received 29 January 2010; accepted 1 February 2010

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.053; wR factor = 0.123; data-to-parameter ratio = 22.9.

The title compound, $[\text{CoCl}_2(\text{C}_{11}\text{H}_{15}\text{N}_3\text{O}_2\text{Si})_2] \cdot \text{C}_3\text{H}_7\text{NO}$, was synthesized from 5-nitro-1-trimethylsilylmethyl-1*H*-benzimidazole and cobalt(II) chloride in dimethylformamide. The Co^{II} atom is coordinated in a distorted tetrahedral environment by two Cl atoms and two N atoms. In the crystal structure, there are a number of C—H...Cl and C—H...O hydrogen-bonding interactions between symmetry-related molecules.

Related literature

For the structures and properties of benzimidazole compounds and their metal complexes, see: Akkurt *et al.* (2005); Castro *et al.* (2002); Küçükbay *et al.* (1996, 2004, 2009); Liu *et al.* (2004); Lukevics *et al.* (2001); Pınar *et al.* (2006); Pan & Xu (2004); Türktekin *et al.* (2004); Tavman *et al.* (2005); Özdemir *et al.* (2005); Çetinkaya *et al.* (1996).



Experimental

Crystal data

$[\text{CoCl}_2(\text{C}_{11}\text{H}_{15}\text{N}_3\text{O}_2\text{Si})_2] \cdot \text{C}_3\text{H}_7\text{NO}$
 $M_r = 701.63$

Triclinic, $P\bar{1}$
 $a = 9.8982$ (4) Å

$b = 11.6936$ (5) Å
 $c = 15.9293$ (6) Å
 $\alpha = 106.041$ (2)°
 $\beta = 107.408$ (2)°
 $\gamma = 99.040$ (3)°
 $V = 1631.97$ (12) Å³

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.81$ mm⁻¹
 $T = 100$ K
 $0.20 \times 0.12 \times 0.08$ mm

Data collection

Bruker APEXII QUAZAR diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2008)
 $T_{\min} = 0.890$, $T_{\max} = 0.937$

30895 measured reflections
8851 independent reflections
5342 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.059$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.123$
 $S = 1.02$
8851 reflections

387 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.90$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.48$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C5—H5...Cl1 ⁱ	0.95	2.79	3.564 (3)	140
C7—H7...O5 ⁱⁱ	0.95	2.32	3.132 (4)	143
C8—H8B...O5 ⁱⁱ	0.99	2.54	3.406 (4)	145
C19—H19A...Cl2 ⁱⁱⁱ	0.99	2.67	3.659 (3)	175
C19—H19B...O4 ⁱ	0.99	2.38	3.182 (4)	137
C22—H22C...Cl1 ^{iv}	0.98	2.82	3.695 (3)	149
C24—H24A...O5	0.98	2.42	2.793 (5)	102

Symmetry codes: (i) $x+1, y, z$; (ii) $-x+1, -y+2, -z+1$; (iii) $-x+1, -y+2, -z+2$; (iv) $-x+1, -y+1, -z+2$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SIR97 (Altomare *et al.*, 1999); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997) and PLATON (Spek, 2009); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON.

We thank Dr Holger Ott (Bruker AXS GmbH, Karlsruhe, Germany) for the data collection. HK & NŞ also thank İnönü University Research Fund (BAPB-2008-60) for financial support of this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5186).

References

- Akkurt, M., Karaca, S., Küçükbay, H., Orhan, E. & Büyükgüngör, O. (2005). *Acta Cryst.* **E61**, m41–m43.
Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
Bruker (2008). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruker (2009). *APEX2* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
Castro, J., Pérez Lourido, P., Sousa-Pedrares, A., Labisbal, E., Piso, J. & García-Vázquez, J. A. (2002). *Acta Cryst.* **C58**, m319–m322.
Çetinkaya, B., Çetinkaya, E., Küçükbay, H. & Durmaz, R. (1996). *Arzneim. Forsch. Drug Res.* **46**, 1154–1158.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.

- Küçükbay, H., Çetinkaya, B., Guesmi, S. & Dixneuf, P. H. (1996). *Organometallics*, **15**, 2434–2439.
- Küçükbay, H., Durmaz, R., Okuyucu, N., Günal, S. & Kazaz, C. (2004). *Arzneim. Forsch. Drug Res.* **54**, 64–68.
- Küçükbay, H., Durmaz, R., Şireci, N. & Günal, S. (2009). *Asian J. Chem.* **21**, 6181–6189.
- Liu, J.-W., Gao, S., Huo, L.-H. & Ng, S. W. (2004). *Acta Cryst.* **E60**, m501–m503.
- Lukevics, E., Arsenyan, P., Shestakova, I., Domracheva, I., Nesterova, A. & Pudova, O. (2001). *Eur. J. Med. Chem.* **36**, 507–515.
- Özdemir, İ., Demir, S. & Çetinkaya, B. (2005). *Tetrahedron*, **61**, 9791–9798.
- Pan, T.-T. & Xu, D.-J. (2004). *Acta Cryst.* **E60**, m56–m58.
- Pınar, Ş., Akkurt, M., Küçükbay, H., Orhan, E. & Büyükgüngör, O. (2006). *Acta Cryst.* **E62**, m1663–m1665.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Tavman, A., Birteksöz, S. & Ötük, G. (2005). *Folia Microbio.* **50**, 467–472.
- Türktekin, S., Akkurt, M., Orhan, E., Küçükbay, F. Z., Küçükbay, H. & Büyükgüngör, O. (2004). *Acta Cryst.* **E60**, m1220–m1222.

supplementary materials

Acta Cryst. (2010). E66, m253-m254 [doi:10.1107/S1600536810003922]

Dichloridobis[5-nitro-1-trimethylsilylmethyl-1*H*-benzimidazole- κN^3]cobalt(II) *N,N*-dimethylformamide solvate

M. Akkurt, S.P. Yalçın, N. Sireci, H. Küçükbay and M. N. Tahir

Comment

Benzimidazole compounds and their metal complexes have been extensively investigated for their versatile properties such as biological activities (Küçükbay *et al.*, 2004; Çetinkaya *et al.*, 1996; Küçükbay *et al.*, 2009; Tavman *et al.*, 2005) and catalytic activities of their metal complexes in many organic syntheses (Küçükbay *et al.*, 1996, Özdemir *et al.*, 2005). Contrary to an extensive chemistry about substituted alkyl derivatives of benzimidazole, there are a limited number of studies of alkylsilyl substituted benzimidazoles (Lukevics *et al.*, 2001). Alkylsilyl substituted benzimidazole derivatives exhibit important *in vitro* cytotoxic activity. The insertion of the silicon atom into the *N*-alkyl chain in benzimidazoles increases the cytotoxic activity. For example, 1-(3-trimethylsilylpropyl)benzimidazole inhibits carcinoma S-180 tumour growth in dose 1 mg kg⁻¹ by 62 % (on ICR mice) (Lukevics *et al.*, 2001). The objective of the present study was to synthesize a trimethylsilylmethyl and NO₂ substituted benzimidazole Co^{II} complex for the first time and compare it to those of related benzimidazole derivatives (Türktekin *et al.*, 2004; Pinar *et al.*, 2006; Akkurt *et al.*, 2005) reported previously.

In the title molecule (Fig. 1), the Co^{II} atom is coordinated in a distorted tetrahedral environment by two Cl atoms and two N atoms. The bond lengths involving the Co atoms are Co1—N1 = 2.013 (3), Co1—N4 = 2.013 (2), Co1—Cl1 = 2.2330 (10) and Co1—Cl2 = 2.2455 (9) Å, while the angles around the Co atom are Cl1—Co1—Cl2 = 112.94 (4), Cl1—Co1—N1 = 111.06 (8), Cl1—Co1—N4 = 110.18 (7), Cl2—Co1—N1 = 106.74 (7), Cl2—Co1—N4 = 111.81 (7) and N1—Co1—N4 = 103.67 (9)°. The average Co—N bond length of 2.013 (3) Å is almost equal to the value of 2.008 (2) Å in dichlorobis[1-(2-ethoxyethyl)-1*H*-benzimidazole- κN^3]cobalt(II) (Türktekin *et al.*, 2004) and 2.032 (2) Å in bis[1-(but-2-enyl)-5-nitro-1*H*-benzimidazole- κN^3]dichlorocobalt(II) (Pinar *et al.*, 2006).

The Co—Cl bond lengths [2.2330 (10) and 2.2455 (9) Å] are comparable to the values of 2.2525 (8) Å in quinolinium trichloro(quinoline-*N*)cobaltate(II) (Pan & Xu, 2004) and 2.236 (1) Å in dichlorobis(1-propylimidazolidine-2-thione- κS)cobalt(II) (Castro *et al.*, 2002) and 2.2680 (8) Å in bis[1-(but-2-enyl)-5-nitro-1*H*-benzimidazole- κN^3]dichlorocobalt(II) (Pinar *et al.*, 2006), but shorter than the value of 2.391 (1) Å observed in aquachlorobis(1,10-phenanthroline)cobalt(II)chloride dimethyl formamide solvate (Liu *et al.*, 2004).

The two benzimidazole ring systems N1/N2/C1—C7 and N4/N5/C12—C18 are almost planar, with maximum deviations of -0.037 (3) for C6 and -0.016 (2) for N4, respectively. The dihedral angle between them is 64.54 (10)°. The angles around the Si atoms with a distorted tetrahedral geometry range from 106.02 (18)° to 113.60 (17)°.

The crystal structure of the title compound is stabilized by C—H \cdots Cl and C—H \cdots O hydrogen-bonding interactions (Fig. 2 and Table 1).

Experimental

1-(Trimethylsilylmethyl)-5-nitrobenzimidazole used in this work as a starting compounds were prepared from reaction of 5(6)-methylbenzimidazole, (chloromethyl)trimethylsilane and KOH under reflux in EtOH.

A solution of 1-(trimethylsilylmethyl)-5-nitrobenzimidazole (2.0 g, 8.02 mmol) and cobalt(II) chloride (0.52 g, 4.01 mmol) in DMF (4 ml) was heated under reflux for 2 h. The mixture was then cooled to room temperature, after which the solvent was then removed from the filtrate in vacuo. The precipitate was then crystallized from EtOH / DMF(2:1). Yield : 2.30 g, 91%; m.p.: 473-474 K. Analysis calculated for $C_{22}H_{30}N_6O_4Si_2CoCl_2 \cdot HCON(CH_3)_2$: C 42.80, H 5.32, N 13.97%. Found: C 42.79, H 5.31, N 13.92%. IR : $\nu_{(C=N)}$: 1523 cm^{-1} . $^1\text{H-NMR}$ (DMSO- d_6): $\delta = 7.97$ ppm (s, 2H, N=CH—N), 6.39 (m, 6H, Ar—H), 4.25 (m, 4H, CH_2Si), -0.27 (s, 18H, Si (CH_3) $_3$).

Refinement

All H atoms were placed at calculated positions and treated as riding atoms, with C—H = 0.95–0.99 Å, and $U_{iso}(H) = 1.2$ or $1.5U_{eq}(C)$.

Figures




Fig. 1. The title molecule showing the atom-labelling scheme. The probability level for the anisotropic displacement parameters is at 50%.




Fig. 2. View of the packing diagram of the title compound in the unit cell. Hydrogen bonds are indicated as dashed lines.

Dichloridobis[5-nitro-1-trimethylsilylmethyl-1H-benzimidazole- κN^3]cobalt(II) *N,N*-dimethylformamide solvate

Crystal data

$[CoCl_2(C_{11}H_{15}N_3O_2Si)_2] \cdot C_3H_7NO$

$M_r = 701.63$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 9.8982(4)\text{ \AA}$

$b = 11.6936(5)\text{ \AA}$

$Z = 2$

$F(000) = 730$

$D_x = 1.428\text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\text{ \AA}$

Cell parameters from 4864 reflections

$\theta = 5.4\text{--}51.0^\circ$

$c = 15.9293 (6) \text{ \AA}$
 $\alpha = 106.041 (2)^\circ$
 $\beta = 107.408 (2)^\circ$
 $\gamma = 99.040 (3)^\circ$
 $V = 1631.97 (12) \text{ \AA}^3$
 $\mu = 0.81 \text{ mm}^{-1}$
 $T = 100 \text{ K}$
 Plate, blue
 $0.20 \times 0.12 \times 0.08 \text{ mm}$

Data collection

Bruker APEXII QUAZAR
 diffractometer
 Radiation source: ImuS
 multilayer
 ω and φ scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 2008)
 $T_{\min} = 0.890, T_{\max} = 0.937$
 30895 measured reflections
 8851 independent reflections
 5342 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.059$
 $\theta_{\max} = 29.4^\circ, \theta_{\min} = 1.4^\circ$
 $h = -13 \rightarrow 13$
 $k = -16 \rightarrow 16$
 $l = -21 \rightarrow 21$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.123$
 $S = 1.02$
 8851 reflections
 387 parameters
 0 restraints
 Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0506P)^2 + 0.0752P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.90 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.48 \text{ e \AA}^{-3}$

Special details

Experimental. Data were collected on an APEX II QUAZAR diffractometer equipped with a brilliant, high intense I μ S (microfocus source) with multilayer mirrors for monochromation and collimation.

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell esds are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating $-R$ -factor-obs etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

supplementary materials

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.38557 (4)	0.69835 (4)	0.77352 (3)	0.0194 (1)
Cl1	0.23442 (8)	0.51641 (7)	0.74255 (6)	0.0299 (2)
Cl2	0.28764 (9)	0.80333 (7)	0.68299 (5)	0.0305 (3)
Si1	0.80522 (9)	0.66034 (8)	0.52977 (6)	0.0220 (3)
Si2	0.76085 (9)	0.79463 (8)	1.19186 (6)	0.0242 (3)
O1	0.6900 (3)	0.3815 (2)	0.93714 (16)	0.0374 (8)
O2	0.9030 (3)	0.3653 (2)	0.93302 (19)	0.0490 (10)
O3	-0.0582 (2)	0.8517 (2)	0.90291 (16)	0.0357 (8)
O4	-0.0395 (2)	0.8620 (2)	1.04422 (17)	0.0393 (8)
N1	0.5723 (3)	0.6770 (2)	0.75277 (16)	0.0202 (8)
N2	0.7576 (3)	0.7092 (2)	0.70128 (16)	0.0210 (8)
N3	0.7937 (3)	0.4046 (2)	0.91233 (19)	0.0310 (9)
N4	0.4540 (2)	0.7981 (2)	0.91071 (16)	0.0183 (7)
N5	0.6010 (2)	0.8884 (2)	1.06061 (15)	0.0172 (7)
N6	0.0135 (3)	0.8609 (2)	0.98339 (19)	0.0264 (8)
C1	0.6695 (3)	0.6074 (3)	0.78068 (19)	0.0206 (9)
C2	0.6672 (3)	0.5322 (3)	0.8338 (2)	0.0239 (9)
C3	0.7870 (3)	0.4827 (3)	0.8532 (2)	0.0240 (10)
C4	0.9025 (3)	0.5029 (3)	0.8219 (2)	0.0262 (10)
C5	0.9033 (3)	0.5751 (3)	0.7674 (2)	0.0249 (10)
C6	0.7850 (3)	0.6276 (3)	0.74777 (19)	0.0209 (9)
C7	0.6315 (3)	0.7364 (3)	0.70664 (19)	0.0211 (9)
C8	0.8477 (3)	0.7615 (3)	0.6555 (2)	0.0233 (9)
C9	0.8366 (4)	0.5082 (3)	0.5292 (2)	0.0316 (11)
C10	0.9278 (3)	0.7459 (3)	0.4868 (2)	0.0286 (10)
C11	0.6093 (3)	0.6419 (3)	0.4624 (2)	0.0301 (11)
C12	0.3698 (3)	0.8356 (3)	0.96419 (19)	0.0173 (9)
C13	0.2206 (3)	0.8261 (3)	0.9372 (2)	0.0190 (9)
C14	0.1713 (3)	0.8717 (3)	1.0084 (2)	0.0215 (9)
C15	0.2602 (3)	0.9252 (3)	1.1030 (2)	0.0221 (9)
C16	0.4089 (3)	0.9373 (3)	1.1298 (2)	0.0206 (9)
C17	0.4614 (3)	0.8913 (3)	1.05910 (19)	0.0167 (8)
C18	0.5901 (3)	0.8324 (3)	0.97189 (19)	0.0176 (8)
C19	0.7347 (3)	0.9258 (3)	1.14482 (19)	0.0204 (9)
C20	0.6391 (3)	0.7803 (3)	1.2593 (2)	0.0302 (11)
C21	0.9569 (4)	0.8365 (4)	1.2669 (3)	0.0517 (16)
C22	0.7128 (4)	0.6516 (3)	1.0898 (2)	0.0424 (14)
O5	0.3510 (3)	1.0259 (2)	0.34832 (19)	0.0470 (10)
N7	0.3118 (3)	0.8861 (2)	0.41855 (17)	0.0256 (8)
C23	0.3349 (4)	0.9199 (4)	0.3519 (3)	0.0478 (16)
C24	0.2906 (4)	0.9746 (4)	0.4965 (3)	0.0484 (14)
C25	0.2834 (4)	0.7605 (3)	0.4171 (3)	0.0472 (14)
H2	0.58920	0.51540	0.85550	0.0290*
H4	0.98130	0.46640	0.83840	0.0310*
H5	0.98000	0.58890	0.74420	0.0300*

H7	0.58970	0.79200	0.68020	0.0250*
H8A	0.95250	0.77410	0.69220	0.0280*
H8B	0.83170	0.84310	0.65610	0.0280*
H9A	0.81830	0.45610	0.46500	0.0470*
H9B	0.76970	0.46860	0.55370	0.0470*
H9C	0.93820	0.51920	0.56880	0.0470*
H10A	1.03020	0.76020	0.52660	0.0430*
H10B	0.90560	0.82520	0.48890	0.0430*
H10C	0.91240	0.69750	0.42220	0.0430*
H11A	0.59310	0.72290	0.46410	0.0450*
H11B	0.54730	0.60370	0.49030	0.0450*
H11C	0.58410	0.58930	0.39730	0.0450*
H13	0.15650	0.79030	0.87350	0.0230*
H15	0.21820	0.95320	1.14870	0.0270*
H16	0.47250	0.97530	1.19340	0.0250*
H18	0.67190	0.81890	0.95490	0.0210*
H19A	0.72860	0.99600	1.19400	0.0250*
H19B	0.82100	0.95400	1.12940	0.0250*
H20A	0.65820	0.71800	1.28850	0.0450*
H20B	0.53650	0.75540	1.21700	0.0450*
H20C	0.65840	0.85980	1.30800	0.0450*
H21A	0.97770	0.76980	1.29050	0.0780*
H21B	0.97900	0.91240	1.31980	0.0780*
H21C	1.01770	0.84960	1.23020	0.0780*
H22A	0.77640	0.66230	1.05430	0.0640*
H22B	0.61020	0.63470	1.04940	0.0640*
H22C	0.72650	0.58240	1.11190	0.0640*
H23	0.33990	0.85760	0.30080	0.0580*
H24A	0.33100	1.05880	0.50050	0.0720*
H24B	0.34120	0.96270	0.55530	0.0720*
H24C	0.18560	0.96110	0.48560	0.0720*
H25A	0.18070	0.73140	0.40970	0.0710*
H25B	0.34760	0.75680	0.47610	0.0710*
H25C	0.30270	0.70800	0.36470	0.0710*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.0190 (2)	0.0224 (2)	0.0177 (2)	0.0079 (2)	0.0067 (2)	0.0069 (2)
Cl1	0.0287 (4)	0.0262 (4)	0.0359 (4)	0.0061 (3)	0.0135 (4)	0.0110 (4)
Cl2	0.0374 (5)	0.0281 (4)	0.0214 (4)	0.0142 (4)	0.0023 (3)	0.0075 (3)
Si1	0.0215 (5)	0.0251 (5)	0.0226 (4)	0.0078 (4)	0.0107 (4)	0.0090 (4)
Si2	0.0231 (5)	0.0326 (5)	0.0214 (4)	0.0123 (4)	0.0080 (4)	0.0135 (4)
O1	0.0466 (15)	0.0345 (14)	0.0415 (14)	0.0148 (12)	0.0206 (12)	0.0211 (12)
O2	0.0418 (15)	0.0509 (17)	0.0700 (19)	0.0243 (13)	0.0154 (14)	0.0429 (15)
O3	0.0227 (12)	0.0408 (15)	0.0360 (14)	0.0129 (11)	0.0039 (11)	0.0066 (12)
O4	0.0273 (13)	0.0520 (16)	0.0509 (15)	0.0176 (12)	0.0260 (12)	0.0193 (13)
N1	0.0220 (13)	0.0228 (14)	0.0194 (13)	0.0090 (11)	0.0087 (11)	0.0094 (11)

supplementary materials

N2	0.0212 (13)	0.0264 (14)	0.0209 (13)	0.0091 (11)	0.0118 (11)	0.0102 (11)
N3	0.0365 (17)	0.0254 (15)	0.0351 (16)	0.0097 (13)	0.0128 (14)	0.0157 (13)
N4	0.0162 (13)	0.0227 (13)	0.0195 (12)	0.0071 (11)	0.0082 (10)	0.0097 (11)
N5	0.0160 (12)	0.0194 (13)	0.0182 (12)	0.0076 (10)	0.0055 (10)	0.0086 (10)
N6	0.0215 (14)	0.0193 (14)	0.0369 (16)	0.0071 (11)	0.0116 (13)	0.0054 (12)
C1	0.0240 (16)	0.0192 (16)	0.0168 (15)	0.0085 (13)	0.0056 (13)	0.0040 (12)
C2	0.0283 (17)	0.0217 (16)	0.0220 (16)	0.0045 (14)	0.0102 (14)	0.0078 (13)
C3	0.0320 (18)	0.0171 (16)	0.0220 (16)	0.0087 (14)	0.0049 (14)	0.0094 (13)
C4	0.0234 (17)	0.0263 (17)	0.0272 (17)	0.0097 (14)	0.0057 (14)	0.0085 (14)
C5	0.0205 (16)	0.0278 (18)	0.0256 (16)	0.0079 (14)	0.0082 (13)	0.0071 (14)
C6	0.0200 (16)	0.0227 (16)	0.0199 (15)	0.0073 (13)	0.0060 (13)	0.0075 (13)
C7	0.0236 (16)	0.0269 (17)	0.0176 (15)	0.0119 (14)	0.0102 (13)	0.0089 (13)
C8	0.0241 (16)	0.0250 (17)	0.0259 (16)	0.0063 (14)	0.0148 (14)	0.0102 (14)
C9	0.043 (2)	0.0286 (19)	0.0313 (18)	0.0171 (16)	0.0195 (16)	0.0115 (15)
C10	0.0207 (16)	0.038 (2)	0.0291 (17)	0.0065 (15)	0.0111 (14)	0.0130 (15)
C11	0.0245 (17)	0.034 (2)	0.0329 (18)	0.0062 (15)	0.0098 (15)	0.0144 (16)
C12	0.0171 (15)	0.0176 (15)	0.0203 (15)	0.0058 (12)	0.0072 (12)	0.0099 (12)
C13	0.0186 (15)	0.0161 (15)	0.0221 (15)	0.0048 (12)	0.0034 (12)	0.0103 (12)
C14	0.0169 (15)	0.0194 (16)	0.0334 (17)	0.0078 (13)	0.0117 (13)	0.0125 (14)
C15	0.0254 (16)	0.0208 (16)	0.0259 (16)	0.0078 (13)	0.0153 (14)	0.0092 (13)
C16	0.0241 (16)	0.0216 (16)	0.0185 (15)	0.0088 (13)	0.0094 (13)	0.0072 (13)
C17	0.0174 (15)	0.0156 (15)	0.0200 (14)	0.0062 (12)	0.0072 (12)	0.0088 (12)
C18	0.0200 (15)	0.0193 (15)	0.0191 (14)	0.0084 (13)	0.0112 (13)	0.0085 (12)
C19	0.0175 (15)	0.0226 (16)	0.0192 (15)	0.0053 (13)	0.0053 (12)	0.0056 (13)
C20	0.0287 (18)	0.038 (2)	0.0277 (17)	0.0101 (16)	0.0114 (14)	0.0151 (16)
C21	0.026 (2)	0.088 (3)	0.058 (3)	0.020 (2)	0.0110 (19)	0.052 (3)
C22	0.074 (3)	0.036 (2)	0.038 (2)	0.030 (2)	0.032 (2)	0.0223 (18)
O5	0.0476 (16)	0.0460 (16)	0.0694 (19)	0.0174 (13)	0.0287 (14)	0.0420 (15)
N7	0.0230 (14)	0.0266 (15)	0.0286 (14)	0.0060 (12)	0.0061 (12)	0.0153 (12)
C23	0.039 (2)	0.062 (3)	0.055 (3)	0.018 (2)	0.020 (2)	0.033 (2)
C24	0.056 (3)	0.044 (2)	0.041 (2)	0.012 (2)	0.018 (2)	0.0085 (19)
C25	0.045 (2)	0.045 (2)	0.065 (3)	0.0190 (19)	0.021 (2)	0.034 (2)

Geometric parameters (Å, °)

Co1—C11	2.2330 (10)	C14—C15	1.396 (4)
Co1—C12	2.2455 (9)	C15—C16	1.376 (4)
Co1—N1	2.013 (3)	C16—C17	1.390 (4)
Co1—N4	2.013 (2)	C2—H2	0.9500
Si1—C8	1.902 (3)	C4—H4	0.9500
Si1—C9	1.852 (4)	C5—H5	0.9500
Si1—C10	1.854 (3)	C7—H7	0.9500
Si1—C11	1.859 (3)	C8—H8B	0.9900
Si2—C19	1.904 (4)	C8—H8A	0.9900
Si2—C20	1.856 (3)	C9—H9A	0.9800
Si2—C21	1.851 (4)	C9—H9B	0.9800
Si2—C22	1.858 (3)	C9—H9C	0.9800
O1—N3	1.224 (4)	C10—H10A	0.9800
O2—N3	1.232 (4)	C10—H10B	0.9800

O3—N6	1.231 (4)	C10—H10C	0.9800
O4—N6	1.230 (4)	C11—H11B	0.9800
O5—C23	1.244 (5)	C11—H11C	0.9800
N1—C1	1.406 (4)	C11—H11A	0.9800
N1—C7	1.333 (4)	C13—H13	0.9500
N2—C7	1.356 (4)	C15—H15	0.9500
N2—C8	1.472 (4)	C16—H16	0.9500
N2—C6	1.373 (4)	C18—H18	0.9500
N3—C3	1.477 (4)	C19—H19B	0.9900
N4—C12	1.395 (4)	C19—H19A	0.9900
N4—C18	1.324 (4)	C20—H20A	0.9800
N5—C18	1.346 (4)	C20—H20B	0.9800
N5—C19	1.474 (4)	C20—H20C	0.9800
N5—C17	1.381 (4)	C21—H21A	0.9800
N6—C14	1.467 (4)	C21—H21B	0.9800
N7—C25	1.444 (5)	C21—H21C	0.9800
N7—C23	1.298 (5)	C22—H22B	0.9800
N7—C24	1.473 (5)	C22—H22C	0.9800
C1—C6	1.407 (4)	C22—H22A	0.9800
C1—C2	1.381 (5)	C23—H23	0.9500
C2—C3	1.390 (5)	C24—H24A	0.9800
C3—C4	1.390 (4)	C24—H24B	0.9800
C4—C5	1.369 (5)	C24—H24C	0.9800
C5—C6	1.400 (5)	C25—H25A	0.9800
C12—C13	1.386 (4)	C25—H25B	0.9800
C12—C17	1.408 (4)	C25—H25C	0.9800
C13—C14	1.374 (4)		
Co1...H2	3.4200	C22...O1	3.338 (4)
Co1...H13	3.2900	C22...C18	3.333 (5)
Cl1...N4	3.484 (3)	C23...C21 ^{xi}	3.446 (6)
Cl1...C5 ⁱ	3.564 (3)	C23...C24 ^{viii}	3.557 (6)
Cl2...N1	3.420 (3)	C24...C23 ^{viii}	3.557 (6)
Cl2...C7	3.546 (3)	C25...C9 ⁱⁱⁱ	3.597 (5)
Cl1...H5 ⁱ	2.7900	C5...H13 ^v	2.9300
Cl1...H22C ⁱⁱ	2.8200	C5...H8A	2.9400
Cl1...H10C ⁱⁱⁱ	2.8600	C5...H9B	3.0600
Cl1...H20A ⁱⁱ	3.0600	C6...H9B	3.0800
Cl2...H7	3.0300	C8...H5	3.0100
Cl2...H10A ⁱ	2.8400	C9...H9C ^{xii}	3.0900
Cl2...H9A ⁱⁱⁱ	3.0700	C9...H5	3.0900
Cl2...H16 ^{iv}	2.9400	C13...H19B ^{iv}	3.0800
Cl2...H19A ^{iv}	2.6700	C16...H20B	3.0800
Si2...O4 ^v	3.657 (2)	C16...H19A	2.9200
O1...C22	3.338 (4)	C18...H22B	2.9100
O1...C12 ⁱⁱ	3.399 (4)	C19...H16	3.0200
O1...C17 ⁱⁱ	3.329 (4)	C23...H24B ^{viii}	3.0000

supplementary materials

O2...C14 ⁱⁱ	3.208 (4)	C23...H21C ^{xi}	2.9900
O2...O4 ⁱⁱ	3.223 (4)	C24...H20C ^{iv}	3.0200
O2...O2 ^{vi}	3.178 (4)	C25...H11A	3.0600
O2...C15 ⁱⁱ	3.335 (4)	C25...H9B ⁱⁱⁱ	2.8400
O3...C5 ⁱ	3.245 (4)	H2...O1	2.4500
O3...C1 ⁱ	3.253 (4)	H2...Co1	3.4200
O3...O4 ^{vii}	3.129 (3)	H4...O2	2.3800
O3...N2 ⁱ	3.000 (3)	H5...H13 ^v	2.5900
O3...C7 ⁱ	3.413 (4)	H5...H9C	2.5700
O3...C6 ⁱ	2.869 (4)	H5...C11 ^v	2.7900
O3...N6 ^{vii}	3.234 (4)	H5...C8	3.0100
O4...O3 ^{vii}	3.129 (3)	H5...C9	3.0900
O4...Si2 ⁱ	3.657 (2)	H5...H8A	2.5500
O4...O2 ⁱⁱ	3.223 (4)	H7...C12	3.0300
O4...C19 ⁱ	3.182 (4)	H7...H8B	2.5400
O5...C8 ^{viii}	3.406 (4)	H7...O5 ^{viii}	2.3200
O5...C7 ^{viii}	3.132 (4)	H8A...C5	2.9400
O1...H2	2.4500	H8A...H5	2.5500
O1...H20B ⁱⁱ	2.6500	H8B...H7	2.5400
O2...H4	2.3800	H8B...O5 ^{viii}	2.5400
O3...H13	2.4800	H9A...C12 ⁱⁱⁱ	3.0700
O4...H18 ⁱ	2.6800	H9B...C5	3.0600
O4...H19B ⁱ	2.3800	H9B...C6	3.0800
O4...H15	2.4600	H9B...C25 ⁱⁱⁱ	2.8400
O4...H22A ⁱ	2.8000	H9B...H25B ⁱⁱⁱ	2.5600
O4...H21C ⁱ	2.8900	H9C...H5	2.5700
O5...H8B ^{viii}	2.5400	H9C...C9 ^{xii}	3.0900
O5...H7 ^{viii}	2.3200	H10A...C12 ^v	2.8400
O5...H24A	2.4200	H10C...C11 ⁱⁱⁱ	2.8600
O5...H15 ^{ix}	2.8700	H11A...H24A ^{viii}	2.4000
N1...N4	3.166 (3)	H11A...C25	3.0600
N1...C12	3.420 (3)	H11A...H25B	2.5800
N1...N2	2.243 (4)	H11B...H11B ⁱⁱⁱ	2.5900
N1...C18	3.397 (4)	H13...C5 ⁱ	2.9300
N2...O3 ^v	3.000 (3)	H13...Co1	3.2900
N2...N1	2.243 (4)	H13...O3	2.4800
N4...N5	2.231 (3)	H13...H5 ⁱ	2.5900
N4...C11	3.484 (3)	H15...O5 ^{xiii}	2.8700
N4...N1	3.166 (3)	H15...O4	2.4600
N5...N4	2.231 (3)	H16...C12 ^{iv}	2.9400
N5...C12 ^{iv}	3.337 (4)	H16...C19	3.0200

N6...O3 ^{vii}	3.234 (4)	H16...H19A	2.5100
N6...N6 ^{vii}	3.216 (4)	H18...O4 ^v	2.6800
N5...H22B	2.9400	H19A...H16	2.5100
C1...O3 ^v	3.253 (4)	H19A...C16	2.9200
C5...C9	3.485 (4)	H19A...C12 ^{iv}	2.6700
C5...C11 ^v	3.564 (3)	H19B...O4 ^v	2.3800
C5...O3 ^v	3.245 (4)	H19B...C13 ^{iv}	3.0800
C6...C9	3.596 (4)	H20A...C11 ⁱⁱ	3.0600
C6...O3 ^v	2.869 (4)	H20B...C16	3.0800
C7...O5 ^{viii}	3.132 (4)	H20B...O1 ⁱⁱ	2.6500
C7...O3 ^v	3.413 (4)	H20C...C24 ^{iv}	3.0200
C8...O5 ^{viii}	3.406 (4)	H20C...H24B ^{iv}	2.5600
C9...C6	3.596 (4)	H21A...H25A ^x	2.5100
C9...C25 ⁱⁱⁱ	3.597 (5)	H21C...O4 ^v	2.8900
C9...C5	3.485 (4)	H21C...C23 ^x	2.9900
C12...O1 ⁱⁱ	3.399 (4)	H22A...O4 ^v	2.8000
C12...N5 ^{iv}	3.337 (4)	H22B...N5	2.9400
C12...C17 ^{iv}	3.532 (5)	H22B...C18	2.9100
C13...C19 ^{iv}	3.515 (5)	H22C...C11 ⁱⁱ	2.8200
C14...O2 ⁱⁱ	3.208 (4)	H23...H25C	2.2900
C15...O2 ⁱⁱ	3.335 (4)	H24A...O5	2.4200
C16...C18 ^{iv}	3.506 (5)	H24A...H11A ^{viii}	2.4000
C17...C18 ^{iv}	3.492 (5)	H24B...H25B	2.4100
C17...O1 ⁱⁱ	3.329 (4)	H24B...C23 ^{viii}	3.0000
C17...C12 ^{iv}	3.532 (5)	H24B...H20C ^{iv}	2.5600
C18...C22	3.333 (5)	H25A...H21A ^{xi}	2.5100
C18...C16 ^{iv}	3.506 (5)	H25B...H11A	2.5800
C18...C17 ^{iv}	3.492 (5)	H25B...H24B	2.4100
C19...C13 ^{iv}	3.515 (5)	H25B...H9B ⁱⁱⁱ	2.5600
C19...O4 ^v	3.182 (4)	H25C...H23	2.2900
C21...C23 ^x	3.446 (6)		
C11—Co1—Cl2	112.94 (4)	C4—C5—H5	122.00
C11—Co1—N1	111.06 (8)	N1—C7—H7	124.00
C11—Co1—N4	110.18 (7)	N2—C7—H7	123.00
Cl2—Co1—N1	106.74 (7)	Si1—C8—H8A	109.00
Cl2—Co1—N4	111.81 (7)	H8A—C8—H8B	108.00
N1—Co1—N4	103.67 (9)	Si1—C8—H8B	109.00
C8—Si1—C9	108.71 (15)	N2—C8—H8A	109.00
C8—Si1—C10	105.46 (15)	N2—C8—H8B	109.00
C8—Si1—C11	107.76 (14)	Si1—C9—H9A	110.00
C9—Si1—C10	113.60 (17)	Si1—C9—H9B	109.00
C9—Si1—C11	109.95 (17)	Si1—C9—H9C	110.00
C10—Si1—C11	111.09 (15)	H9B—C9—H9C	109.00

supplementary materials

C19—Si2—C20	108.88 (15)	H9A—C9—H9B	109.00
C19—Si2—C21	106.02 (18)	H9A—C9—H9C	110.00
C19—Si2—C22	107.56 (14)	Si1—C10—H10B	109.00
C20—Si2—C21	111.80 (17)	H10A—C10—H10C	109.00
C20—Si2—C22	110.98 (17)	Si1—C10—H10C	109.00
C21—Si2—C22	111.36 (19)	H10A—C10—H10B	110.00
Co1—N1—C1	132.6 (2)	Si1—C10—H10A	109.00
Co1—N1—C7	122.7 (2)	H10B—C10—H10C	109.00
C1—N1—C7	104.7 (3)	Si1—C11—H11C	109.00
C6—N2—C7	107.3 (3)	H11A—C11—H11C	110.00
C6—N2—C8	127.5 (3)	H11B—C11—H11C	109.00
C7—N2—C8	125.2 (3)	H11A—C11—H11B	109.00
O1—N3—O2	123.7 (3)	Si1—C11—H11A	109.00
O1—N3—C3	118.3 (3)	Si1—C11—H11B	109.00
O2—N3—C3	118.0 (3)	C14—C13—H13	122.00
Co1—N4—C12	128.49 (18)	C12—C13—H13	122.00
Co1—N4—C18	126.50 (19)	C14—C15—H15	120.00
C12—N4—C18	104.7 (2)	C16—C15—H15	120.00
C17—N5—C18	107.4 (2)	C17—C16—H16	122.00
C17—N5—C19	126.3 (2)	C15—C16—H16	122.00
C18—N5—C19	125.9 (2)	N4—C18—H18	123.00
O3—N6—O4	123.7 (3)	N5—C18—H18	123.00
O3—N6—C14	118.2 (3)	Si2—C19—H19A	109.00
O4—N6—C14	118.1 (3)	H19A—C19—H19B	108.00
C24—N7—C25	114.4 (3)	Si2—C19—H19B	109.00
C23—N7—C24	120.3 (3)	N5—C19—H19A	109.00
C23—N7—C25	124.7 (3)	N5—C19—H19B	109.00
N1—C1—C6	108.8 (3)	Si2—C20—H20A	109.00
N1—C1—C2	130.4 (3)	Si2—C20—H20B	109.00
C2—C1—C6	120.8 (3)	Si2—C20—H20C	109.00
C1—C2—C3	115.2 (3)	H20B—C20—H20C	110.00
N3—C3—C2	118.0 (3)	H20A—C20—H20B	109.00
N3—C3—C4	117.5 (3)	H20A—C20—H20C	109.00
C2—C3—C4	124.6 (3)	Si2—C21—H21B	109.00
C3—C4—C5	120.3 (3)	H21A—C21—H21C	110.00
C4—C5—C6	116.4 (3)	Si2—C21—H21C	109.00
N2—C6—C1	106.2 (3)	H21A—C21—H21B	109.00
N2—C6—C5	131.0 (3)	Si2—C21—H21A	109.00
C1—C6—C5	122.7 (3)	H21B—C21—H21C	109.00
N1—C7—N2	113.0 (3)	Si2—C22—H22C	109.00
Si1—C8—N2	113.4 (2)	H22A—C22—H22C	109.00
C13—C12—C17	120.4 (3)	H22B—C22—H22C	110.00
N4—C12—C17	109.3 (3)	H22A—C22—H22B	109.00
N4—C12—C13	130.3 (3)	Si2—C22—H22A	109.00
C12—C13—C14	115.7 (3)	Si2—C22—H22B	109.00
N6—C14—C13	117.6 (3)	O5—C23—N7	126.4 (4)
N6—C14—C15	117.7 (3)	O5—C23—H23	117.00
C13—C14—C15	124.7 (3)	N7—C23—H23	117.00
C14—C15—C16	119.8 (3)	N7—C24—H24A	109.00

C15—C16—C17	116.7 (3)	N7—C24—H24B	109.00
N5—C17—C12	105.2 (2)	N7—C24—H24C	109.00
N5—C17—C16	132.0 (3)	H24A—C24—H24B	109.00
C12—C17—C16	122.8 (3)	H24A—C24—H24C	110.00
N4—C18—N5	113.3 (3)	H24B—C24—H24C	109.00
Si2—C19—N5	112.1 (2)	N7—C25—H25A	109.00
C1—C2—H2	122.00	N7—C25—H25B	109.00
C3—C2—H2	122.00	N7—C25—H25C	110.00
C3—C4—H4	120.00	H25A—C25—H25B	109.00
C5—C4—H4	120.00	H25A—C25—H25C	109.00
C6—C5—H5	122.00	H25B—C25—H25C	109.00
C11—Co1—N1—C1	-47.5 (3)	Co1—N4—C18—N5	-174.0 (2)
C12—Co1—N1—C1	-171.0 (2)	C17—N5—C19—Si2	86.6 (3)
N4—Co1—N1—C1	70.8 (3)	C18—N5—C17—C12	-1.0 (4)
C11—Co1—N1—C7	134.4 (2)	C19—N5—C18—N4	173.7 (3)
C12—Co1—N1—C7	10.9 (2)	C19—N5—C17—C12	-174.2 (3)
N4—Co1—N1—C7	-107.3 (2)	C17—N5—C18—N4	0.4 (4)
C11—Co1—N4—C12	-58.2 (3)	C18—N5—C19—Si2	-85.4 (3)
C12—Co1—N4—C12	68.3 (3)	C19—N5—C17—C16	6.6 (6)
N1—Co1—N4—C12	-177.1 (3)	C18—N5—C17—C16	179.8 (4)
C11—Co1—N4—C18	114.8 (3)	O3—N6—C14—C15	157.4 (3)
C12—Co1—N4—C18	-118.7 (3)	O4—N6—C14—C15	-22.3 (4)
N1—Co1—N4—C18	-4.1 (3)	O3—N6—C14—C13	-23.9 (4)
C11—Si1—C8—N2	60.1 (3)	O4—N6—C14—C13	156.4 (3)
C10—Si1—C8—N2	178.8 (2)	C24—N7—C23—O5	4.9 (6)
C9—Si1—C8—N2	-59.0 (3)	C25—N7—C23—O5	175.4 (4)
C22—Si2—C19—N5	41.2 (3)	C2—C1—C6—C5	-0.8 (5)
C20—Si2—C19—N5	-79.2 (2)	N1—C1—C6—N2	0.5 (3)
C21—Si2—C19—N5	160.4 (2)	C6—C1—C2—C3	1.6 (4)
C7—N1—C1—C6	-0.9 (3)	C2—C1—C6—N2	-177.7 (3)
Co1—N1—C1—C6	-179.3 (2)	N1—C1—C6—C5	177.3 (3)
C1—N1—C7—N2	1.0 (3)	N1—C1—C2—C3	-176.1 (3)
C7—N1—C1—C2	177.0 (3)	C1—C2—C3—C4	-1.0 (5)
Co1—N1—C7—N2	179.59 (18)	C1—C2—C3—N3	178.1 (3)
Co1—N1—C1—C2	-1.4 (5)	C2—C3—C4—C5	-0.5 (5)
C7—N2—C6—C1	0.2 (3)	N3—C3—C4—C5	-179.6 (3)
C8—N2—C7—N1	-179.4 (3)	C3—C4—C5—C6	1.3 (5)
C6—N2—C7—N1	-0.8 (3)	C4—C5—C6—C1	-0.7 (5)
C7—N2—C8—Si1	-95.2 (3)	C4—C5—C6—N2	175.3 (3)
C8—N2—C6—C5	2.3 (5)	C17—C12—C13—C14	-1.2 (5)
C8—N2—C6—C1	178.8 (3)	C13—C12—C17—C16	1.0 (6)
C7—N2—C6—C5	-176.4 (3)	N4—C12—C17—N5	1.2 (4)
C6—N2—C8—Si1	86.4 (3)	N4—C12—C17—C16	-179.5 (3)
O1—N3—C3—C2	4.0 (4)	N4—C12—C13—C14	179.4 (3)
O2—N3—C3—C2	-176.8 (3)	C13—C12—C17—N5	-178.4 (3)
O1—N3—C3—C4	-176.9 (3)	C12—C13—C14—N6	-178.5 (3)
O2—N3—C3—C4	2.3 (4)	C12—C13—C14—C15	0.0 (6)
C12—N4—C18—N5	0.3 (4)	N6—C14—C15—C16	-180.0 (3)
C18—N4—C12—C17	-1.0 (4)	C13—C14—C15—C16	1.4 (6)

supplementary materials

Co1—N4—C12—C17	173.2 (2)	C14—C15—C16—C17	-1.6 (5)
Co1—N4—C12—C13	-7.3 (5)	C15—C16—C17—N5	179.6 (4)
C18—N4—C12—C13	178.6 (4)	C15—C16—C17—C12	0.5 (5)

Symmetry codes: (i) $x-1, y, z$; (ii) $-x+1, -y+1, -z+2$; (iii) $-x+1, -y+1, -z+1$; (iv) $-x+1, -y+2, -z+2$; (v) $x+1, y, z$; (vi) $-x+2, -y+1, -z+2$; (vii) $-x, -y+2, -z+2$; (viii) $-x+1, -y+2, -z+1$; (ix) $x, y, z-1$; (x) $x+1, y, z+1$; (xi) $x-1, y, z-1$; (xii) $-x+2, -y+1, -z+1$; (xiii) $x, y, z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C5—H5 \cdots C11 ^v	0.95	2.79	3.564 (3)	140
C7—H7 \cdots O5 ^{viii}	0.95	2.32	3.132 (4)	143
C8—H8B \cdots O5 ^{viii}	0.99	2.54	3.406 (4)	145
C19—H19A \cdots C12 ^{iv}	0.99	2.67	3.659 (3)	175
C19—H19B \cdots O4 ^v	0.99	2.38	3.182 (4)	137
C22—H22C \cdots C11 ⁱⁱ	0.98	2.82	3.695 (3)	149
C24—H24A \cdots O5	0.98	2.42	2.793 (5)	102

Symmetry codes: (v) $x+1, y, z$; (viii) $-x+1, -y+2, -z+1$; (iv) $-x+1, -y+2, -z+2$; (ii) $-x+1, -y+1, -z+2$.

Fig. 1




Fig. 2

