

Bis{tris[3-(2-pyridyl)-1*H*-pyrazole]-iron(II)} dodecamolybdo(V,VI)phosphate hexahydrate

Luijiang Hao,* Tongjun Liu, Jiangkui Chen and Xiaofei Zhang

College of Food and Biological Engineering, Shandong Institute of Light Industry, Jinan 250353, People's Republic of China

Correspondence e-mail: luijianghao001@yahoo.com.cn


Received 4 February 2010; accepted 8 February 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.042\text{ \AA}$; H-atom completeness 78%; disorder in main residue; R factor = 0.039; wR factor = 0.142; data-to-parameter ratio = 12.0.

Crystals of the title compound, $[\text{Fe}(\text{C}_8\text{H}_7\text{N}_3)_3]_2[\text{PMo}_{12}\text{O}_{40}] \cdot 6\text{H}_2\text{O}$, prepared under hydrothermal conditions, are isotopic with the Mn^{2+} and Cd^{2+} analogues. The Fe^{2+} cation is in a distorted octahedral coordination by six N atoms from three chelating 3-(2-pyridyl)-1*H*-pyrazole ligands. The heteropolyanion $[\text{PMo}_{12}\text{O}_{40}]^{4-}$ is a one-electron reduced species in which two O atoms of the central PO_4 group ($\bar{1}$ symmetry) are equally disordered about an inversion centre. $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds make a contribution to the crystal packing. The $\text{Fe}-\text{N}$ bond lengths [$2.085(19)$ – $2.15(2)\text{ \AA}$] are somewhat shorter than the $\text{Mn}-\text{N}$ and $\text{Cd}-\text{N}$ bond lengths [$2.224(6)$ – $2.283(5)$ and $2.316(7)$ – $2.334(6)\text{ \AA}$, respectively]. All other bond lengths and angles and the hydrogen-bonding motifs are very similar in the isotopic structures.

Related literature

For the isotopic Mn^{2+} and Cd^{2+} structures, see: Hao, Ma *et al.* (2010); Hao, Wang *et al.* (2010). For general background to polyoxometalates, see: Pope & Müller (1991). For polyoxometalates modified with amines, see: Zhang, Dou *et al.* (2009); Zhang, Wei *et al.* (2009). For other structures containing the one-electron reduced heteropolyanion $[\text{PMo}_{12}\text{O}_{40}]^{4-}$, see: Artero & Proust (2000); Kurmoo *et al.* (1998); Niu *et al.* (1999). For the role of amines in hydrothermal synthesis, see: Yang *et al.* (2003).



Experimental

Crystal data

$[\text{Fe}(\text{C}_8\text{H}_7\text{N}_3)_3]_2[\text{PMo}_{12}\text{O}_{40}] \cdot 6\text{H}_2\text{O}$	$V = 8117.4(8)\text{ \AA}^3$
$M_r = 2913.04$	$Z = 4$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 18.6816(10)\text{ \AA}$	$\mu = 2.26\text{ mm}^{-1}$
$b = 16.2224(10)\text{ \AA}$	$T = 293\text{ K}$
$c = 27.6405(15)\text{ \AA}$	$0.12 \times 0.10 \times 0.08\text{ mm}$
$\beta = 104.295(1)^{\circ}$	

Data collection

Bruker APEXII CCD diffractometer	20565 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2001)	7090 independent reflections
$T_{\min} = 0.773$, $T_{\max} = 0.840$	5550 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.042$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$	592 parameters
$wR(F^2) = 0.142$	H-atom parameters constrained
$S = 1.00$	$\Delta\rho_{\text{max}} = 1.36\text{ e \AA}^{-3}$
7090 reflections	$\Delta\rho_{\text{min}} = -0.80\text{ e \AA}^{-3}$

Table 1
Selected bond lengths (\AA).

$\text{Fe1}-\text{N}8$	$2.085(19)$	$\text{Fe1}-\text{N}7$	$2.15(2)$
$\text{Fe1}-\text{N}5$	$2.08(2)$	$\text{P1}-\text{O}21\text{A}^{\ddagger}$	$1.49(2)$
$\text{Fe1}-\text{N}2$	$2.11(2)$	$\text{P1}-\text{O}21\text{B}^{\ddagger}$	$1.52(2)$
$\text{Fe1}-\text{N}1$	$2.129(17)$	$\text{P1}-\text{O}19\text{B}^{\ddagger}$	$1.54(2)$
$\text{Fe1}-\text{N}4$	$2.13(2)$	$\text{P1}-\text{O}19\text{A}^{\ddagger}$	$1.56(2)$

Symmetry code: (i) $-x + \frac{1}{2}, -y + \frac{1}{2}, -z$.

Table 2
Hydrogen-bond geometry (\AA , $^{\circ}$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}3-\text{H}3\text{A}\cdots\text{O}17^{\ddagger}$	0.86	2.03	2.82 (3)	153
$\text{N}6-\text{H}6\cdots\text{O}2\text{W}$	0.86	1.98	2.82 (5)	167
$\text{N}9-\text{H}9\text{A}\cdots\text{O}1\text{W}$	0.86	1.95	2.77 (3)	157

Symmetry code: (ii) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT-Plus* (Bruker, 2001); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Financial support from the International Cooperation Program for Excellent Lecturers of 2008 of Shandong Provincial Education Department, the Research Award Fund for Outstanding Young and Middle-aged Scientists of Shandong Province (2008BS04022), Shandong Provincial Education Department and Shandong Institute of Education are gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WM2304).

References

- Artero, V. & Proust, A. (2000). *Eur. J. Inorg. Chem.* pp. 2393–2400
- Bruker (2001). *SAINT-Plus* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2004). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Hao, L., Ma, C., Chen, J., Zhang, X. & Zhang, X. (2010). *Acta Cryst. E66*, m231–m232.
- Hao, L., Wang, Y., Zhang, X., Chen, J. & Zhang, X. (2010). *Acta Cryst. E66*, m268–m269.
- Kurmoo, M., Bonamico, M., Bellitto, C., Fares, V., Federici, F., Guionneau, P., Ducasse, L., Kitagawa, H. & Day, P. (1998). *Adv. Mater.* **7**, 545–550.
- Niu, J. Y., Shan, B. Z. & You, X. Z. (1999). *Transition Met. Chem.* **24**, 108–114.
- Pope, M. T. & Müller, A. (1991). *Angew. Chem. Int. Ed.* **30**, 34–38.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Yang, W. B., Lu, C. Z., Wu, C. D. & Zhuang, H. H. (2003). *Chin. J. Struct. Chem.* **22**, 137–142.
- Zhang, X. T., Dou, J. M., Wei, P. H., Li, D. C., Li, B., Shi, C. W. & Hu, B. (2009). *Inorg. Chim. Acta*, **362**, 3325–3332.
- Zhang, X. T., Wei, P. H., Sun, D. F., Ni, Z. H., Dou, J. M., Li, B., Shi, C. W. & Hu, B. (2009). *Cryst. Growth Des.* **9**, 4424–4428.

supplementary materials

Acta Cryst. (2010). E66, m283-m284 [doi:10.1107/S1600536810004861]

Bis{tris[3-(2-pyridyl)-1H-pyrazole]iron(II)} dodecamolybdo(V,VI)phosphate hexahydrate

L. Hao, T. Liu, J. Chen and X. Zhang

Comment

The design and synthesis of polyoxometalates has attracted continuous research interest not only because of their appealing structural and topological novelties, but also due to their interesting optical, electronic, magnetic, and catalytic properties, as well as their potential medical applications (Pope & Müller, 1991). In our research group, organic amines, such as 3-(2-pyridyl)-1H-pyrazole and pyrazine, are used to effectively modify polyoxomolybdates under hydrothermal condicions (Zhang, Dou *et al.*, 2009; Zhang, Wei *et al.*, 2009). Here, we describe the synthesis and structural characterization of the title compound.

As shown in Figure 1, the asymmetric unit of the title compound consists of three subunits, *viz.* of a complex $[\text{Fe}(\text{C}_8\text{H}_7\text{N}_3)_3]^{2+}$ cation, half of a $[\text{PMo}_{12}\text{O}_{40}]^{4-}$ heteropolyanion and of three uncoordinated water molecules. The Fe^{2+} ion is in a distorted octahedral coordination by six N atoms from three chelating 3-(2-pyridyl)-1H-pyrazole ligands.

The heteropolyanion $[\text{PMo}_{12}\text{O}_{40}]^{4-}$ anion is an one-electron reduced derivative of $[\text{PMo}_{12}\text{O}_{40}]^{3-}$, similar to anions with different counter cations as reported by Artero & Proust (2000); Kurmoo *et al.* (1998); Niu *et al.* (1999). The employed organic ligand appears to adjust the pH value, and additionally supplies reducing electrons, which is a commonly observed feature of hydrothermal syntheses when organic amines are used to prepare various hybrid materials, zeolites or metal phosphates (Yang *et al.*, 2003).

In the Keggin-type heteropolyanion, each Mo atom is surrounded by six O atoms and the P atom is located at the center of the anion. There are four kinds of O atoms present in the anion according to their coordination environments: O_a (O atoms in the disordered PO_4 tetrahedron), O_b (bridging O atoms between two triplet groups of MoO_6 octahedra), O_c (bridging O atoms within one triplet group of MoO_6 octahedra) and O_d (terminal O atoms). The P—O bond distances are in the normal range of 1.49 (2)—1.56 (2) Å. The Mo—O bond distances vary widely from 1.636 (15) to 2.51 (2) Å. The shortest Mo—O bonds are in the range of 1.636 (2)—1.659 (15) Å for the terminal oxygen atoms. The longest Mo—O lengths are in the range of 2.43 (2)—2.51 (2) Å for those oxygen atoms connected with both Mo and P atoms.

N—H···O and O—H···O hydrogen bonding between the cationic and anionic moieties and the uncoordinated water molecules leads to a consolidation of the structure (Fig. 2; Table 2).

The crystal structure of $[(\text{Fe}(\text{C}_8\text{H}_7\text{N}_3)_3)_2[\text{PMo}_{12}\text{O}_{40}](\text{H}_2\text{O})_6$ is isotypic with the Mn^{2+} and Cd^{2+} analogues, $[(\text{Mn}(\text{C}_8\text{H}_7\text{N}_3)_3)_2[\text{PMo}_{12}\text{O}_{40}](\text{H}_2\text{O})_6$ (Hao, Ma *et al.*, 2010) and $[(\text{Cd}(\text{C}_8\text{H}_7\text{N}_3)_3)_2[\text{PMo}_{12}\text{O}_{40}](\text{H}_2\text{O})_6$ (Hao, Wang *et al.*, 2010). In comparison with the Mn^{2+} and Cd^{2+} analogues, the Fe—N bond lengths (2.085 (19)—2.15 (2) Å) are somewhat shorter than the Mn —N and Cd —N bond lengths with 2.224 (6)—2.283 (5) Å and 2.316 (7)—2.334 (6) Å, respectively. All other bond lengths and angles and the hydrogen-bonding motifs are very similar in the isotypic structures.

supplementary materials

Experimental

A mixture of 3-(2-pyridyl)-1*H*-pyrazole (0.5 mmol 0.07 g), sodium molybdate (0.4 mmol, 0.10 g), iron(III) chloride hexahydrate (0.25 mmol, 0.07 g), and dipotassium hydrogenphosphate (0.22 mmol, 0.05 g) in 10 ml distilled water was sealed in a 25 ml Teflon-lined stainless steel autoclave and was kept at 433 K for three days. Brown crystals suitable for the X-ray experiment were obtained. IR(cm^{-1}): 3441, 3118, 2674, 2546, 1696, 1605, 1540, 1429, 1356, 1291, 1107, 1014, 931, 876, 802, 719, 562, 507.

TGA curve shows a separation of lattice water molecules and the organic ligands above 352 and 655 K, respectively. The overall thermal decomposition process can be described by the followed equation: $4\text{C}_{48}\text{H}_{54}\text{Fe}_2\text{Mo}_{12}\text{N}_{18}\text{O}_{46}\text{P} + 327\text{O}_2 = 108\text{H}_2\text{O} + 192\text{CO}_2 + 36\text{N}_2\text{O}_5 + 4\text{Fe}_2\text{O}_3 + 2\text{P}_2\text{O}_5 + 48\text{MoO}_3$

Refinement

All hydrogen atoms bound to aromatic carbon atoms were refined in calculated positions using a riding model with a C—H distance of 0.93 Å and $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{C})$. Hydrogen atoms attached to aromatic N atoms were refined with a N—H distance of 0.86 Å and $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{N})$. The hydrogen atoms of the three uncoordinated water molecules could not be located unambiguously from difference Fourier maps, probably due to disorder of the water molecules. Thus the structure was refined without the H atoms of the water molecules (which includes the water O atoms O1W, O2W, O3W). In the PO₄ unit, the two oxygen atoms (O19 and O21) are equally disordered about the inversion centre. In the final difference Fourier map the highest peak is 2.82 Å from atom O2w and the deepest hole is 0.79 Å from atom Mo1. The highest peak is located in the voids of the crystal structure and may be associated with an additional water molecule. However, refinement of this position did not result in a reasonable model. Hence this position was also excluded from the final refinement.

Figures




Fig. 1. The building blocks of the title compound with the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level; H atoms are given as spheres of arbitrary radius.





Fig. 2. The crystal packing of the title compound, displayed with N—H···O and O—H···O hydrogen bonds as dashed lines.

Bis{tris[3-(2-pyridyl)-1*H*-pyrazole]iron(II)} dodecamolybdo(V,VI)phosphate hexahydrate

Crystal data



$$F(000) = 5628$$

$$M_r = 2913.04$$

$$D_x = 2.384 \text{ Mg m}^{-3}$$

Monoclinic, $C2/c$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Hall symbol: -C 2yc	Cell parameters from 9828 reflections
$a = 18.6816(10)$ Å	$\theta = 2.4\text{--}28.2^\circ$
$b = 16.2224(10)$ Å	$\mu = 2.26 \text{ mm}^{-1}$
$c = 27.6405(15)$ Å	$T = 293$ K
$\beta = 104.295(1)^\circ$	Block, brown
$V = 8117.4(8)$ Å ³	$0.12 \times 0.10 \times 0.08$ mm
$Z = 4$	

Data collection

Bruker APEXII CCD diffractometer	7090 independent reflections
Radiation source: fine-focus sealed tube graphite	5550 reflections with $I > 2\sigma(I)$
phi and ω scans	$R_{\text{int}} = 0.042$
Absorption correction: multi-scan (SADABS; Bruker, 2001)	$\theta_{\text{max}} = 25.0^\circ, \theta_{\text{min}} = 1.5^\circ$
$T_{\text{min}} = 0.773, T_{\text{max}} = 0.840$	$h = -22 \rightarrow 17$
20565 measured reflections	$k = -18 \rightarrow 19$
	$l = -32 \rightarrow 31$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.039$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.142$	H-atom parameters constrained
$S = 1.00$	$w = 1/[\sigma^2(F_o^2) + (0.1P)^2 + 20.1545P]$ where $P = (F_o^2 + 2F_c^2)/3$
7090 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
592 parameters	$\Delta\rho_{\text{max}} = 1.36 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.80 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

supplementary materials

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.2037 (13)	0.8989 (13)	0.0951 (8)	0.049 (5)	
H1	0.2470	0.8948	0.0844	0.059*	
C2	0.1427 (14)	0.9335 (14)	0.0638 (8)	0.054 (6)	
H2	0.1446	0.9517	0.0323	0.065*	
C3	0.0793 (13)	0.9410 (15)	0.0791 (8)	0.055 (6)	
H3	0.0379	0.9660	0.0588	0.065*	
C4	0.0780 (13)	0.9107 (15)	0.1252 (9)	0.055 (6)	
H4	0.0345	0.9123	0.1359	0.066*	
C5	0.1406 (12)	0.8781 (12)	0.1553 (7)	0.042 (5)	
C6	0.1448 (12)	0.8483 (13)	0.2053 (8)	0.045 (5)	
C7	0.0917 (16)	0.845 (2)	0.2338 (10)	0.076 (8)	
H7	0.0423	0.8603	0.2244	0.091*	
C8	0.1307 (19)	0.814 (2)	0.2788 (9)	0.079 (9)	
H8	0.1119	0.8047	0.3065	0.095*	
C9	0.3237 (16)	0.9793 (18)	0.2596 (11)	0.071 (7)	
H9	0.2903	0.9589	0.2766	0.085*	
C10	0.3526 (18)	1.0535 (19)	0.2716 (12)	0.076 (8)	
H10	0.3402	1.0833	0.2971	0.091*	
C11	0.400 (2)	1.086 (3)	0.2472 (15)	0.098 (11)	
H11	0.4198	1.1378	0.2557	0.118*	
C12	0.4175 (17)	1.043 (2)	0.2108 (14)	0.090 (10)	
H12	0.4493	1.0655	0.1931	0.108*	
C13	0.3882 (13)	0.9665 (18)	0.2001 (11)	0.066 (7)	
C14	0.4075 (15)	0.915 (2)	0.1621 (11)	0.074 (8)	
C15	0.4577 (19)	0.926 (3)	0.1312 (14)	0.101 (11)	
H15	0.4894	0.9699	0.1306	0.121*	
C16	0.4475 (19)	0.853 (2)	0.1010 (13)	0.091 (10)	
H16	0.4719	0.8392	0.0766	0.109*	
C17	0.4084 (15)	0.758 (2)	0.2901 (12)	0.077 (8)	
H17	0.4135	0.8138	0.2988	0.093*	
C18	0.4475 (19)	0.701 (2)	0.3232 (13)	0.097 (11)	
H18	0.4766	0.7174	0.3540	0.116*	
C19	0.443 (2)	0.622 (2)	0.3106 (14)	0.103 (12)	
H19	0.4709	0.5832	0.3319	0.123*	
C20	0.3977 (17)	0.5978 (18)	0.2657 (11)	0.078 (8)	
H20	0.3943	0.5426	0.2563	0.093*	
C21	0.3579 (13)	0.6562 (15)	0.2356 (9)	0.054 (6)	
C22	0.3055 (13)	0.6380 (14)	0.1877 (9)	0.054 (6)	
C23	0.2857 (17)	0.5644 (17)	0.1626 (10)	0.069 (7)	
H23	0.3049	0.5122	0.1716	0.083*	
C24	0.2312 (19)	0.5861 (18)	0.1213 (11)	0.077 (8)	
H24	0.2051	0.5507	0.0968	0.092*	
Fe1	0.29393 (17)	0.8180 (2)	0.19341 (13)	0.0479 (8)	
Mo1	0.24259 (11)	0.14187 (11)	0.11115 (6)	0.0414 (5)	
Mo2	0.19058 (10)	0.45820 (10)	-0.01680 (7)	0.0420 (5)	

Mo3	0.42269 (9)	0.31452 (11)	-0.01975 (7)	0.0386 (5)	
Mo4	0.35343 (9)	0.40933 (11)	0.07504 (6)	0.0398 (5)	
Mo5	0.41686 (9)	0.20224 (11)	0.09088 (6)	0.0405 (5)	
Mo6	0.17783 (10)	0.34811 (12)	0.09231 (6)	0.0420 (5)	
N1	0.2034 (9)	0.8711 (11)	0.1403 (6)	0.042 (4)	
N2	0.2095 (10)	0.8208 (12)	0.2318 (7)	0.051 (5)	
N3	0.1991 (13)	0.7990 (14)	0.2763 (7)	0.068 (6)	
H3A	0.2327	0.7779	0.3001	0.081*	
N4	0.3407 (12)	0.9323 (13)	0.2238 (8)	0.064 (6)	
N5	0.3703 (13)	0.8449 (17)	0.1521 (8)	0.072 (7)	
N6	0.3958 (16)	0.8079 (19)	0.1153 (10)	0.093 (9)	
H6	0.3805	0.7608	0.1026	0.112*	
N7	0.3639 (11)	0.7377 (13)	0.2466 (8)	0.057 (5)	
N8	0.2677 (11)	0.7008 (11)	0.1633 (7)	0.052 (5)	
N9	0.2227 (12)	0.6693 (13)	0.1233 (8)	0.061 (5)	
H9A	0.1918	0.6975	0.1013	0.073*	
O1	0.4070 (10)	0.3947 (9)	0.0244 (6)	0.062 (5)	
O2	0.1470 (9)	0.3952 (11)	0.1364 (6)	0.059 (4)	
O3	0.2402 (9)	0.0913 (10)	0.1620 (5)	0.056 (4)	
O4	0.3455 (11)	0.1738 (12)	0.1246 (7)	0.077 (6)	
O5	0.4049 (11)	0.3141 (10)	0.1060 (6)	0.065 (5)	
O6	0.2789 (8)	0.3892 (13)	0.1084 (7)	0.071 (5)	
O7	0.4527 (11)	0.2401 (10)	0.0354 (6)	0.065 (5)	
O8	0.4029 (10)	0.4828 (10)	0.1090 (6)	0.062 (4)	
O9	0.4022 (12)	0.2143 (10)	-0.0602 (7)	0.085 (7)	
O10	0.5038 (9)	0.3410 (10)	-0.0293 (7)	0.062 (4)	
O11	0.2853 (9)	0.4702 (15)	0.0264 (7)	0.083 (7)	
O12	0.1036 (11)	0.4003 (12)	-0.0543 (7)	0.083 (7)	
O13	0.1436 (12)	0.1335 (10)	0.0730 (8)	0.088 (7)	
O14	0.1569 (9)	0.4260 (15)	0.0409 (7)	0.084 (6)	
O15	0.1641 (9)	0.5548 (9)	-0.0236 (7)	0.064 (5)	
O16	0.2690 (11)	0.0585 (13)	0.0706 (8)	0.087 (7)	
O17	0.2216 (10)	0.2520 (10)	0.1270 (7)	0.078 (6)	
O18	0.4938 (9)	0.1788 (12)	0.1315 (6)	0.066 (5)	
O19A	0.2032 (13)	0.2368 (14)	0.0391 (8)	0.030 (5)	0.50
O21A	0.3250 (13)	0.2802 (15)	0.0252 (9)	0.030 (5)	0.50
O19B	0.2909 (13)	0.1861 (14)	0.0380 (8)	0.028 (5)	0.50
O21B	0.2480 (14)	0.3317 (15)	0.0266 (9)	0.032 (5)	0.50
O1W	0.1172 (13)	0.7180 (15)	0.0392 (9)	0.096 (7)	
O2W	0.362 (3)	0.657 (2)	0.0638 (17)	0.186 (11)	
O3W	0.534 (2)	0.071 (3)	0.0275 (19)	0.199 (13)	
P1	0.2500	0.2500	0.0000	0.0217 (12)	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.047 (12)	0.048 (12)	0.051 (13)	-0.004 (10)	0.012 (10)	0.016 (10)
C2	0.064 (15)	0.048 (12)	0.046 (12)	-0.015 (12)	0.007 (11)	0.009 (10)

supplementary materials

C3	0.041 (12)	0.062 (14)	0.047 (13)	-0.004 (11)	-0.014 (10)	0.011 (11)
C4	0.039 (12)	0.070 (15)	0.055 (14)	0.001 (11)	0.008 (10)	0.004 (12)
C5	0.043 (11)	0.040 (11)	0.040 (11)	-0.002 (9)	0.005 (9)	0.002 (9)
C6	0.037 (11)	0.053 (12)	0.044 (12)	-0.006 (10)	0.006 (9)	0.003 (10)
C7	0.063 (17)	0.11 (2)	0.067 (17)	0.011 (16)	0.035 (14)	0.013 (16)
C8	0.09 (2)	0.11 (2)	0.037 (13)	-0.009 (19)	0.020 (14)	0.018 (14)
C9	0.070 (18)	0.070 (17)	0.080 (19)	0.018 (15)	0.033 (15)	0.027 (15)
C10	0.08 (2)	0.071 (19)	0.076 (19)	0.009 (16)	0.014 (15)	0.006 (15)
C11	0.08 (2)	0.10 (3)	0.10 (3)	0.01 (2)	0.00 (2)	0.02 (2)
C12	0.051 (17)	0.11 (3)	0.10 (2)	-0.010 (18)	-0.005 (17)	0.02 (2)
C13	0.036 (12)	0.077 (18)	0.076 (17)	0.007 (12)	-0.001 (12)	0.032 (14)
C14	0.049 (15)	0.10 (2)	0.070 (18)	0.022 (16)	0.016 (13)	0.038 (17)
C15	0.068 (17)	0.13 (2)	0.11 (2)	0.018 (17)	0.018 (16)	0.051 (19)
C16	0.082 (18)	0.12 (2)	0.082 (18)	0.038 (17)	0.050 (15)	0.028 (17)
C17	0.053 (16)	0.08 (2)	0.09 (2)	-0.001 (15)	0.006 (15)	-0.021 (17)
C18	0.07 (2)	0.10 (3)	0.09 (2)	-0.006 (19)	-0.027 (18)	0.00 (2)
C19	0.09 (3)	0.09 (2)	0.10 (3)	0.02 (2)	-0.02 (2)	0.01 (2)
C20	0.074 (19)	0.065 (17)	0.08 (2)	0.012 (14)	-0.002 (16)	0.008 (15)
C21	0.044 (12)	0.054 (14)	0.064 (15)	0.003 (11)	0.016 (11)	0.000 (11)
C22	0.054 (14)	0.049 (13)	0.061 (14)	0.010 (11)	0.021 (11)	-0.004 (11)
C23	0.09 (2)	0.053 (14)	0.066 (17)	0.008 (14)	0.018 (15)	-0.002 (13)
C24	0.10 (2)	0.067 (18)	0.063 (17)	0.006 (16)	0.015 (16)	-0.017 (13)
Fe1	0.0393 (17)	0.0462 (18)	0.060 (2)	0.0076 (14)	0.0151 (14)	0.0097 (14)
Mo1	0.0516 (11)	0.0396 (10)	0.0317 (9)	-0.0035 (8)	0.0077 (8)	0.0034 (7)
Mo2	0.0421 (10)	0.0313 (9)	0.0549 (11)	0.0073 (7)	0.0165 (8)	0.0039 (7)
Mo3	0.0257 (9)	0.0415 (10)	0.0474 (11)	-0.0038 (7)	0.0068 (7)	-0.0020 (8)
Mo4	0.0308 (9)	0.0425 (10)	0.0436 (10)	-0.0059 (7)	0.0044 (7)	-0.0096 (7)
Mo5	0.0281 (9)	0.0501 (11)	0.0394 (10)	0.0028 (7)	0.0006 (7)	0.0023 (8)
Mo6	0.0423 (10)	0.0483 (11)	0.0350 (10)	0.0105 (8)	0.0087 (8)	-0.0084 (7)
N1	0.036 (9)	0.047 (10)	0.043 (10)	-0.002 (8)	0.010 (7)	0.005 (8)
N2	0.043 (10)	0.056 (11)	0.044 (10)	-0.007 (9)	-0.005 (8)	0.010 (8)
N3	0.073 (16)	0.082 (15)	0.040 (11)	-0.016 (12)	-0.001 (10)	0.020 (10)
N4	0.058 (13)	0.063 (13)	0.078 (14)	0.019 (11)	0.029 (11)	0.027 (11)
N5	0.062 (14)	0.098 (18)	0.063 (14)	0.034 (14)	0.027 (11)	0.022 (13)
N6	0.09 (2)	0.12 (2)	0.074 (16)	0.054 (18)	0.024 (15)	0.018 (15)
N7	0.040 (10)	0.067 (13)	0.064 (13)	-0.002 (9)	0.010 (9)	-0.007 (10)
N8	0.051 (11)	0.051 (11)	0.050 (11)	0.007 (9)	0.006 (9)	-0.004 (9)
N9	0.071 (14)	0.056 (12)	0.051 (12)	0.003 (10)	0.007 (10)	-0.006 (9)
O1	0.094 (13)	0.045 (9)	0.059 (10)	0.024 (9)	0.042 (9)	0.009 (7)
O2	0.048 (9)	0.081 (12)	0.048 (9)	0.008 (8)	0.013 (7)	-0.022 (8)
O3	0.057 (10)	0.069 (10)	0.042 (8)	-0.010 (8)	0.009 (7)	0.012 (7)
O4	0.071 (12)	0.086 (13)	0.089 (13)	-0.046 (10)	0.049 (11)	-0.049 (11)
O5	0.101 (14)	0.048 (9)	0.062 (10)	0.026 (9)	0.049 (10)	0.012 (8)
O6	0.030 (8)	0.114 (15)	0.067 (11)	0.004 (9)	0.010 (7)	0.040 (10)
O7	0.099 (14)	0.051 (9)	0.055 (10)	0.025 (9)	0.040 (9)	0.011 (8)
O8	0.075 (12)	0.049 (9)	0.058 (10)	-0.020 (9)	0.009 (8)	-0.014 (8)
O9	0.092 (13)	0.039 (8)	0.087 (12)	0.007 (9)	-0.046 (10)	-0.015 (8)
O10	0.047 (9)	0.065 (10)	0.085 (12)	-0.003 (8)	0.038 (9)	-0.001 (9)
O11	0.038 (9)	0.146 (19)	0.068 (11)	0.010 (11)	0.015 (8)	0.052 (12)

O12	0.078 (13)	0.092 (14)	0.096 (14)	-0.049 (11)	0.055 (11)	-0.052 (11)
O13	0.086 (14)	0.043 (9)	0.096 (14)	0.018 (9)	-0.053 (11)	-0.021 (9)
O14	0.037 (9)	0.137 (18)	0.074 (12)	-0.004 (10)	0.007 (8)	0.054 (12)
O15	0.052 (10)	0.036 (8)	0.101 (13)	0.014 (7)	0.012 (9)	0.006 (8)
O16	0.079 (13)	0.097 (14)	0.105 (14)	-0.053 (12)	0.060 (12)	-0.062 (12)
O17	0.069 (12)	0.046 (9)	0.088 (13)	0.007 (8)	-0.038 (10)	-0.013 (9)
O18	0.051 (10)	0.094 (13)	0.043 (9)	0.029 (9)	-0.007 (7)	0.004 (8)
O19A	0.023 (12)	0.037 (13)	0.028 (12)	0.002 (10)	0.004 (10)	-0.005 (10)
O21A	0.022 (12)	0.033 (13)	0.036 (13)	-0.001 (10)	0.008 (10)	-0.001 (10)
O19B	0.032 (13)	0.025 (11)	0.025 (12)	0.001 (10)	0.003 (10)	-0.003 (9)
O21B	0.035 (14)	0.032 (13)	0.029 (13)	0.000 (11)	0.006 (10)	-0.001 (10)
O1W	0.077 (14)	0.102 (16)	0.095 (16)	0.019 (13)	-0.005 (12)	0.004 (13)
O2W	0.231 (11)	0.150 (11)	0.239 (11)	-0.014 (3)	0.179 (4)	-0.011 (3)
O3W	0.166 (13)	0.165 (13)	0.311 (13)	0.000 (3)	0.140 (5)	-0.042 (3)
P1	0.022 (3)	0.020 (3)	0.022 (3)	0.002 (2)	0.004 (2)	-0.001 (2)

Geometric parameters (Å, °)

C1—N1	1.33 (3)		Mo1—O19B	2.51 (2)
C1—C2	1.37 (3)		Mo2—O15	1.641 (14)
C1—H1	0.9300		Mo2—O16 ⁱ	1.847 (17)
C2—C3	1.36 (3)		Mo2—O11	1.884 (18)
C2—H2	0.9300		Mo2—O14	1.925 (17)
C3—C4	1.37 (3)		Mo2—O12	1.941 (18)
C3—H3	0.9300		Mo2—O19B ⁱ	2.46 (2)
C4—C5	1.36 (3)		Mo2—O21B	2.48 (2)
C4—H4	0.9300		Mo3—O10	1.659 (15)
C5—N1	1.34 (3)		Mo3—O1	1.855 (15)
C5—C6	1.45 (3)		Mo3—O13 ⁱ	1.875 (16)
C6—N2	1.33 (3)		Mo3—O7	1.916 (15)
C6—C7	1.41 (3)		Mo3—O9	1.958 (16)
C7—C8	1.37 (4)		Mo3—O19A ⁱ	2.43 (2)
C7—H7	0.9300		Mo3—O21A	2.51 (2)
C8—N3	1.32 (4)		Mo4—O8	1.652 (15)
C8—H8	0.9300		Mo4—O6	1.878 (16)
C9—C10	1.33 (4)		Mo4—O11	1.887 (17)
C9—N4	1.35 (3)		Mo4—O5	1.907 (16)
C9—H9	0.9300		Mo4—O1	1.926 (15)
C10—C11	1.34 (5)		Mo4—O21B	2.44 (2)
C10—H10	0.9300		Mo4—O21A	2.49 (2)
C11—C12	1.33 (5)		Mo5—O18	1.636 (15)
C11—H11	0.9300		Mo5—O4	1.864 (16)
C12—C13	1.36 (4)		Mo5—O5	1.888 (15)
C12—H12	0.9300		Mo5—O7	1.920 (15)
C13—N4	1.35 (3)		Mo5—O12 ⁱ	1.935 (17)
C13—C14	1.45 (4)		Mo5—O19B	2.46 (2)
C14—N5	1.33 (4)		Mo5—O21A	2.51 (2)
C14—C15	1.43 (4)		Mo6—O2	1.657 (14)

supplementary materials

C15—C16	1.43 (5)	Mo6—O9 ⁱ	1.841 (17)
C15—H15	0.9300	Mo6—O14	1.870 (17)
C16—N6	1.34 (4)	Mo6—O17	1.907 (17)
C16—H16	0.9300	Mo6—O6	1.948 (16)
C17—N7	1.33 (3)	Mo6—O19A	2.45 (2)
C17—C18	1.38 (4)	Mo6—O21B	2.50 (2)
C17—H17	0.9300	N2—N3	1.34 (3)
C18—C19	1.32 (5)	N3—H3A	0.8600
C18—H18	0.9300	N5—N6	1.37 (3)
C19—C20	1.38 (4)	N6—H6	0.8600
C19—H19	0.9300	N8—N9	1.32 (3)
C20—C21	1.36 (3)	N9—H9A	0.8600
C20—H20	0.9300	O9—Mo6 ⁱ	1.841 (17)
C21—N7	1.36 (3)	O12—Mo5 ⁱ	1.935 (17)
C21—C22	1.47 (4)	O13—Mo3 ⁱ	1.875 (16)
C22—N8	1.33 (3)	O16—Mo2 ⁱ	1.847 (17)
C22—C23	1.38 (3)	O19A—P1	1.56 (2)
C23—C24	1.37 (4)	O19A—O21A ⁱ	1.75 (3)
C23—H23	0.9300	O19A—Mo3 ⁱ	2.43 (2)
C24—N9	1.36 (3)	O21A—P1	1.49 (2)
C24—H24	0.9300	O21A—O21B	1.67 (3)
Fe1—N8	2.085 (19)	O21A—O19B	1.72 (3)
Fe1—N5	2.08 (2)	O21A—O19A ⁱ	1.75 (3)
Fe1—N2	2.11 (2)	O19B—P1	1.54 (2)
Fe1—N1	2.129 (17)	O19B—O21B ⁱ	1.78 (3)
Fe1—N4	2.13 (2)	O19B—Mo2 ⁱ	2.46 (2)
Fe1—N7	2.15 (2)	O21B—P1	1.52 (2)
Mo1—O3	1.639 (14)	O21B—O19B ⁱ	1.78 (3)
Mo1—O13	1.893 (18)	P1—O21A ⁱ	1.49 (2)
Mo1—O16	1.898 (17)	P1—O21B ⁱ	1.52 (2)
Mo1—O17	1.903 (17)	P1—O19B ⁱ	1.54 (2)
Mo1—O4	1.937 (17)	P1—O19A ⁱ	1.56 (2)
Mo1—O19A	2.48 (2)		
N1—C1—C2	122 (2)	O18—Mo5—O4	102.4 (10)
N1—C1—H1	118.8	O18—Mo5—O5	101.7 (9)
C2—C1—H1	118.8	O4—Mo5—O5	89.4 (8)
C1—C2—C3	120 (2)	O18—Mo5—O7	101.6 (9)
C1—C2—H2	120.2	O4—Mo5—O7	155.9 (9)
C3—C2—H2	120.2	O5—Mo5—O7	87.3 (7)
C2—C3—C4	118 (2)	O18—Mo5—O12 ⁱ	100.7 (10)
C2—C3—H3	120.8	O4—Mo5—O12 ⁱ	88.6 (7)
C4—C3—H3	120.9	O5—Mo5—O12 ⁱ	157.4 (9)
C5—C4—C3	120 (2)	O7—Mo5—O12 ⁱ	85.4 (7)
C5—C4—H4	120.2	O18—Mo5—O19B	159.0 (9)
C3—C4—H4	120.2	O4—Mo5—O19B	64.8 (8)

N1—C5—C4	122 (2)	O5—Mo5—O19B	94.9 (8)
N1—C5—C6	114.9 (18)	O7—Mo5—O19B	91.7 (8)
C4—C5—C6	123 (2)	O12 ⁱ —Mo5—O19B	64.0 (8)
N2—C6—C7	110 (2)	O18—Mo5—O21A	160.3 (9)
N2—C6—C5	117.8 (19)	O4—Mo5—O21A	92.4 (9)
C7—C6—C5	132 (2)	O5—Mo5—O21A	65.1 (8)
C8—C7—C6	103 (3)	O7—Mo5—O21A	64.6 (8)
C8—C7—H7	128.3	O12 ⁱ —Mo5—O21A	92.5 (9)
C6—C7—H7	128.3	O19B—Mo5—O21A	40.6 (7)
N3—C8—C7	109 (2)	O2—Mo6—O9 ⁱ	102.7 (9)
N3—C8—H8	125.6	O2—Mo6—O14	101.7 (9)
C7—C8—H8	125.6	O9 ⁱ —Mo6—O14	90.7 (9)
C10—C9—N4	123 (3)	O2—Mo6—O17	101.0 (9)
C10—C9—H9	118.7	O9 ⁱ —Mo6—O17	89.4 (7)
N4—C9—H9	118.6	O14—Mo6—O17	156.7 (10)
C9—C10—C11	120 (3)	O2—Mo6—O6	99.7 (8)
C9—C10—H10	119.8	O9 ⁱ —Mo6—O6	157.5 (10)
C11—C10—H10	119.8	O14—Mo6—O6	87.2 (7)
C12—C11—C10	120 (4)	O17—Mo6—O6	83.9 (8)
C12—C11—H11	120.0	O2—Mo6—O19A	159.3 (8)
C10—C11—H11	120.1	O9 ⁱ —Mo6—O19A	63.9 (9)
C11—C12—C13	118 (4)	O14—Mo6—O19A	94.4 (9)
C11—C12—H12	120.8	O17—Mo6—O19A	64.9 (8)
C13—C12—H12	120.8	O6—Mo6—O19A	94.0 (8)
N4—C13—C12	123 (3)	O2—Mo6—O21B	157.2 (8)
N4—C13—C14	115 (3)	O9 ⁱ —Mo6—O21B	95.8 (9)
C12—C13—C14	121 (3)	O14—Mo6—O21B	64.5 (9)
N5—C14—C15	111 (3)	O17—Mo6—O21B	92.4 (9)
N5—C14—C13	116 (2)	O6—Mo6—O21B	63.2 (8)
C15—C14—C13	133 (3)	O19A—Mo6—O21B	43.3 (8)
C14—C15—C16	104 (3)	C1—N1—C5	117.9 (18)
C14—C15—H15	128.1	C1—N1—Fe1	126.7 (15)
C16—C15—H15	128.2	C5—N1—Fe1	115.4 (13)
N6—C16—C15	106 (3)	C6—N2—N3	106 (2)
N6—C16—H16	126.9	C6—N2—Fe1	115.0 (14)
C15—C16—H16	126.8	N3—N2—Fe1	138.8 (16)
N7—C17—C18	123 (3)	C8—N3—N2	111 (2)
N7—C17—H17	118.6	C8—N3—H3A	124.4
C18—C17—H17	118.6	N2—N3—H3A	124.4
C19—C18—C17	119 (3)	C9—N4—C13	115 (3)
C19—C18—H18	120.3	C9—N4—Fe1	129.3 (19)
C17—C18—H18	120.3	C13—N4—Fe1	115 (2)
C18—C19—C20	120 (3)	C14—N5—N6	106 (3)
C18—C19—H19	120.0	C14—N5—Fe1	117.3 (19)
C20—C19—H19	120.1	N6—N5—Fe1	137 (2)
C21—C20—C19	118 (3)	N5—N6—C16	113 (3)
C21—C20—H20	120.8	N5—N6—H6	123.6

supplementary materials

C19—C20—H20	120.8	C16—N6—H6	123.5
N7—C21—C20	123 (2)	C17—N7—C21	117 (2)
N7—C21—C22	113 (2)	C17—N7—Fe1	127.5 (19)
C20—C21—C22	124 (2)	C21—N7—Fe1	115.8 (16)
N8—C22—C23	111 (2)	N9—N8—C22	106.2 (19)
N8—C22—C21	117 (2)	N9—N8—Fe1	136.8 (15)
C23—C22—C21	131 (2)	C22—N8—Fe1	117.0 (16)
C24—C23—C22	104 (2)	N8—N9—C24	111 (2)
C24—C23—H23	127.8	N8—N9—H9A	124.4
C22—C23—H23	127.8	C24—N9—H9A	124.4
N9—C24—C23	107 (3)	Mo3—O1—Mo4	139.4 (10)
N9—C24—H24	126.6	Mo5—O4—Mo1	140.2 (11)
C23—C24—H24	126.5	Mo5—O5—Mo4	138.8 (11)
N8—Fe1—N5	95.6 (10)	Mo4—O6—Mo6	137.9 (11)
N8—Fe1—N2	95.1 (8)	Mo3—O7—Mo5	138.2 (10)
N5—Fe1—N2	166.1 (9)	Mo6 ⁱ —O9—Mo3	138.8 (13)
N8—Fe1—N1	91.4 (7)	Mo2—O11—Mo4	139.5 (13)
N5—Fe1—N1	94.2 (7)	Mo5 ⁱ —O12—Mo2	136.6 (12)
N2—Fe1—N1	76.8 (6)	Mo3 ⁱ —O13—Mo1	140.3 (13)
N8—Fe1—N4	169.6 (8)	Mo6—O14—Mo2	139.8 (12)
N5—Fe1—N4	76.3 (10)	Mo2 ⁱ —O16—Mo1	142.9 (12)
N2—Fe1—N4	93.9 (7)	Mo1—O17—Mo6	137.6 (11)
N1—Fe1—N4	95.7 (7)	P1—O19A—O21A ⁱ	53.0 (10)
N8—Fe1—N7	76.4 (8)	P1—O19A—Mo3 ⁱ	124.8 (12)
N5—Fe1—N7	96.8 (8)	O21A ⁱ —O19A—Mo3 ⁱ	71.9 (11)
N2—Fe1—N7	94.2 (7)	P1—O19A—Mo6	122.6 (13)
N1—Fe1—N7	164.3 (7)	O21A ⁱ —O19A—Mo6	132.3 (14)
N4—Fe1—N7	97.8 (8)	Mo3 ⁱ —O19A—Mo6	93.7 (8)
O3—Mo1—O13	102.2 (9)	P1—O19A—Mo1	122.3 (12)
O3—Mo1—O16	102.5 (9)	O21A ⁱ —O19A—Mo1	132.4 (14)
O13—Mo1—O16	88.9 (9)	Mo3 ⁱ —O19A—Mo1	92.5 (8)
O3—Mo1—O17	102.8 (9)	Mo6—O19A—Mo1	92.2 (8)
O13—Mo1—O17	88.2 (7)	P1—O21A—O21B	57.2 (12)
O16—Mo1—O17	154.6 (10)	P1—O21A—O19B	56.7 (11)
O3—Mo1—O4	101.8 (8)	O21B—O21A—O19B	94.2 (16)
O13—Mo1—O4	155.9 (10)	P1—O21A—O19A ⁱ	57.2 (11)
O16—Mo1—O4	85.1 (7)	O21B—O21A—O19A ⁱ	93.2 (16)
O17—Mo1—O4	87.5 (8)	O19B—O21A—O19A ⁱ	92.0 (16)
O3—Mo1—O19A	159.1 (8)	P1—O21A—Mo4	125.7 (13)
O13—Mo1—O19A	62.8 (9)	O21B—O21A—Mo4	68.6 (11)
O16—Mo1—O19A	92.1 (9)	O19B—O21A—Mo4	132.1 (14)
O17—Mo1—O19A	64.2 (8)	O19A ⁱ —O21A—Mo4	131.7 (15)
O4—Mo1—O19A	94.2 (8)	P1—O21A—Mo5	124.8 (13)
O3—Mo1—O19B	157.6 (8)	O21B—O21A—Mo5	132.4 (15)
O13—Mo1—O19B	93.9 (9)	O19B—O21A—Mo5	68.1 (11)
O16—Mo1—O19B	62.0 (8)	O19A ⁱ —O21A—Mo5	129.6 (14)

O17—Mo1—O19B	93.0 (9)	Mo4—O21A—Mo5	90.6 (8)
O4—Mo1—O19B	62.7 (8)	P1—O21A—Mo3	123.9 (13)
O19A—Mo1—O19B	43.3 (7)	O21B—O21A—Mo3	129.4 (15)
O15—Mo2—O16 ⁱ	102.7 (10)	O19B—O21A—Mo3	130.4 (14)
O15—Mo2—O11	100.6 (9)	O19A ⁱ —O21A—Mo3	66.7 (10)
O16 ⁱ —Mo2—O11	91.0 (9)	Mo4—O21A—Mo3	90.3 (8)
O15—Mo2—O14	101.7 (10)	Mo5—O21A—Mo3	91.1 (8)
O16 ⁱ —Mo2—O14	155.4 (10)	P1—O19B—O21A	53.8 (11)
O11—Mo2—O14	87.2 (7)	P1—O19B—O21B ⁱ	54.0 (10)
O15—Mo2—O12	102.3 (9)	O21A—O19B—O21B ⁱ	91.1 (15)
O16 ⁱ —Mo2—O12	87.4 (7)	P1—O19B—Mo2 ⁱ	123.7 (12)
O11—Mo2—O12	156.9 (10)	O21A—O19B—Mo2 ⁱ	135.5 (14)
O14—Mo2—O12	84.8 (8)	O21B ⁱ —O19B—Mo2 ⁱ	69.7 (10)
O15—Mo2—O19B ⁱ	159.9 (8)	P1—O19B—Mo5	125.1 (12)
O16 ⁱ —Mo2—O19B ⁱ	63.8 (8)	O21A—O19B—Mo5	71.3 (11)
O11—Mo2—O19B ⁱ	94.8 (9)	O21B ⁱ —O19B—Mo5	135.2 (15)
O14—Mo2—O19B ⁱ	91.9 (9)	Mo2 ⁱ —O19B—Mo5	94.2 (8)
O12—Mo2—O19B ⁱ	63.9 (8)	P1—O19B—Mo1	121.7 (12)
O15—Mo2—O21B	158.0 (9)	O21A—O19B—Mo1	129.9 (13)
O16 ⁱ —Mo2—O21B	93.0 (10)	O21B ⁱ —O19B—Mo1	128.5 (14)
O11—Mo2—O21B	63.5 (9)	Mo2 ⁱ —O19B—Mo1	91.1 (7)
O14—Mo2—O21B	64.3 (9)	Mo5—O19B—Mo1	91.9 (7)
O12—Mo2—O21B	93.6 (9)	P1—O21B—O21A	55.3 (11)
O19B ⁱ —Mo2—O21B	42.1 (8)	P1—O21B—O19B ⁱ	55.0 (11)
O10—Mo3—O1	102.8 (8)	O21A—O21B—O19B ⁱ	92.2 (16)
O10—Mo3—O13 ⁱ	102.2 (10)	P1—O21B—Mo4	127.1 (14)
O1—Mo3—O13 ⁱ	91.1 (8)	O21A—O21B—Mo4	71.8 (12)
O10—Mo3—O7	101.0 (9)	O19B ⁱ —O21B—Mo4	135.2 (15)
O1—Mo3—O7	89.0 (7)	P1—O21B—Mo2	123.1 (13)
O13 ⁱ —Mo3—O7	156.2 (10)	O21A—O21B—Mo2	132.4 (15)
O10—Mo3—O9	100.5 (10)	O19B ⁱ —O21B—Mo2	68.2 (10)
O1—Mo3—O9	156.6 (9)	Mo4—O21B—Mo2	91.9 (8)
O13 ⁱ —Mo3—O9	85.8 (7)	P1—O21B—Mo6	121.5 (13)
O7—Mo3—O9	84.8 (8)	O21A—O21B—Mo6	132.6 (15)
O10—Mo3—O19A ⁱ	158.1 (8)	O19B ⁱ —O21B—Mo6	126.0 (15)
O1—Mo3—O19A ⁱ	94.8 (8)	Mo4—O21B—Mo6	92.4 (8)
O13 ⁱ —Mo3—O19A ⁱ	64.2 (9)	Mo2—O21B—Mo6	91.2 (8)
O7—Mo3—O19A ⁱ	92.1 (8)	O21A ⁱ —P1—O21A	180 (3)
O9—Mo3—O19A ⁱ	63.0 (9)	O21A ⁱ —P1—O21B ⁱ	67.6 (13)
O10—Mo3—O21A	160.2 (8)	O21A—P1—O21B ⁱ	112.4 (13)
O1—Mo3—O21A	65.0 (8)	O21A ⁱ —P1—O21B	112.4 (13)
O13 ⁱ —Mo3—O21A	93.9 (10)	O21A—P1—O21B	67.6 (13)

supplementary materials

O7—Mo3—O21A	64.7 (8)	O21B ⁱ —P1—O21B	180 (2)
O9—Mo3—O21A	92.0 (9)	O21A ⁱ —P1—O19B	110.5 (12)
O19A ⁱ —Mo3—O21A	41.4 (8)	O21A—P1—O19B	69.5 (12)
O8—Mo4—O6	103.0 (9)	O21B ⁱ —P1—O19B	71.0 (12)
O8—Mo4—O11	102.3 (10)	O21B—P1—O19B	109.0 (12)
O6—Mo4—O11	89.4 (7)	O21A ⁱ —P1—O19B ⁱ	69.5 (12)
O8—Mo4—O5	100.3 (9)	O21A—P1—O19B ⁱ	110.5 (12)
O6—Mo4—O5	89.5 (7)	O21B ⁱ —P1—O19B ⁱ	109.0 (12)
O11—Mo4—O5	157.0 (10)	O21B—P1—O19B ⁱ	71.0 (12)
O8—Mo4—O1	100.6 (8)	O19B—P1—O19B ⁱ	180 (3)
O6—Mo4—O1	156.4 (9)	O21A ⁱ —P1—O19A ⁱ	110.1 (12)
O11—Mo4—O1	85.8 (7)	O21A—P1—O19A ⁱ	69.9 (12)
O5—Mo4—O1	86.1 (6)	O21B ⁱ —P1—O19A ⁱ	72.6 (13)
O8—Mo4—O21B	161.3 (9)	O21B—P1—O19A ⁱ	107.4 (13)
O6—Mo4—O21B	65.4 (8)	O19B—P1—O19A ⁱ	107.2 (12)
O11—Mo4—O21B	64.4 (9)	O19B ⁱ —P1—O19A ⁱ	72.8 (12)
O5—Mo4—O21B	94.5 (8)	O21A ⁱ —P1—O19A	69.9 (12)
O1—Mo4—O21B	91.8 (8)	O21A—P1—O19A	110.1 (12)
O8—Mo4—O21A	159.1 (8)	O21B ⁱ —P1—O19A	107.4 (13)
O6—Mo4—O21A	92.4 (9)	O21B—P1—O19A	72.6 (13)
O11—Mo4—O21A	91.8 (9)	O19B—P1—O19A	72.8 (12)
O5—Mo4—O21A	65.3 (8)	O19B ⁱ —P1—O19A	107.2 (12)
O1—Mo4—O21A	64.7 (7)	O19A ⁱ —P1—O19A	180 (2)
O21B—Mo4—O21A	39.6 (8)		


Symmetry codes: (i) $-x+1/2, -y+1/2, -z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N3—H3A \cdots O17 ⁱⁱ	0.86	2.03	2.82 (3)	153.
N6—H6 \cdots O2W	0.86	1.98	2.82 (5)	167.
N9—H9A \cdots O1W	0.86	1.95	2.77 (3)	157.

Symmetry codes: (ii) $-x+1/2, y+1/2, -z+1/2$.

Fig. 1



supplementary materials

Fig. 2

