

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-Ethylsulfinyl-2-(4-fluorophenyl)-5-iodo-7-methyl-1-benzofuran

Hong Dae Choi,^a Pil Ja Seo,^a Byeng Wha Son^b and Uk Lee^{b*}^aDepartment of Chemistry, Donggeui University, San 24 Kaya-dong Busanjin-gu, Busan 614-714, Republic of Korea, and ^bDepartment of Chemistry, Pukyong National University, 599-1 Daeyeon 3-dong, Nam-gu, Busan 608-737, Republic of Korea

Correspondence e-mail: uklee@pknu.ac.kr


Received 1 February 2010; accepted 10 February 2010

Key indicators: single-crystal X-ray study; $T = 273$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.022; wR factor = 0.059; data-to-parameter ratio = 16.8.

In the title compound, $\text{C}_{17}\text{H}_{14}\text{FIO}_2\text{S}$, the 4-fluorophenyl ring is rotated out of the benzofuran plane, as indicated by the dihedral angle of $15.60(6)^\circ$. The crystal structure exhibits an $\text{I} \cdots \text{O}$ interaction [$3.052(2)$ Å].

Related literature

For the crystal structures of similar 3-ethylsulfinyl-2-(4-fluorophenyl)-5-halo-1-benzofuran derivatives, see: Choi *et al.* (2010*a,b*). For the pharmacological activity of benzofuran compounds, see: Aslam *et al.* (2006); Galal *et al.* (2009); Khan *et al.* (2005). For natural products with benzofuran rings, see: Akgul & Anil (2003); Soekamto *et al.* (2003). For a review of halogen bonding, see: Politzer *et al.* (2007).



Experimental

Crystal data

 $\text{C}_{17}\text{H}_{14}\text{FIO}_2\text{S}$ $M_r = 428.24$ Triclinic, $P\bar{1}$ $a = 7.4681(5)$ Å $b = 10.3023(7)$ Å $c = 10.9761(7)$ Å $\alpha = 76.289(1)^\circ$ $\beta = 89.138(1)^\circ$ $\gamma = 74.802(1)^\circ$ $V = 790.73(9)$ Å³ $Z = 2$ Mo $K\alpha$ radiation $\mu = 2.17$ mm⁻¹ $T = 273$ K $0.50 \times 0.25 \times 0.25$ mm

Data collection

Bruker SMART APEXII CCD diffractometer

Absorption correction: multi-scan (SADABS; Bruker, 2009)

 $T_{\min} = 0.520$, $T_{\max} = 0.582$

6867 measured reflections

3386 independent reflections

3261 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.049$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.022$ $wR(F^2) = 0.059$ $S = 1.06$

3386 reflections

201 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.71$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.81$ e Å⁻³

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and DIAMOND (Brandenburg, 1998); software used to prepare material for publication: SHELXL97.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PB2023).

References

- Akgul, Y. Y. & Anil, H. (2003). *Phytochemistry*, **63**, 939–943.
 Aslam, S. N., Stevenson, P. C., Phythian, S. J., Veitch, N. C. & Hall, D. R. (2006). *Tetrahedron*, **62**, 4214–4226.
 Brandenburg, K. (1998). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
 Bruker (2009). *SADABS. APEX2 and SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2010*a*). *Acta Cryst.* **E66**, o323.
 Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2010*b*). *Acta Cryst.* **E66**, o402.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Galal, S. A., Abd El-All, A. S., Abdallah, M. M. & El-Diwani, H. I. (2009). *Bioorg. Med. Chem. Lett.* **19**, 2420–2428.
 Khan, M. W., Alam, M. J., Rashid, M. A. & Chowdhury, R. (2005). *Bioorg. Med. Chem.* **13**, 4796–4805.
 Politzer, P., Lane, P., Concha, M. C., Ma, Y. & Murray, J. S. (2007). *J. Mol. Model.* **13**, 305–311.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Soekamto, N. H., Achmad, S. A., Ghisalberti, E. L., Hakim, E. H. & Syah, Y. M. (2003). *Phytochemistry*, **64**, 831–834.

supplementary materials

Acta Cryst. (2010). E66, o629 [doi:10.1107/S1600536810005581]

3-Ethylsulfinyl-2-(4-fluorophenyl)-5-iodo-7-methyl-1-benzofuran

H. D. Choi, P. J. Seo, B. W. Son and U. Lee

Comment

Compounds containing benzofuran skeleton show diverse pharmacological activities such as antifungal (Aslam *et al.*, 2006), antitumor and antiviral (Galal *et al.*, 2009), antimicrobial (Khan *et al.*, 2005) properties. These compounds widely occur in nature (Akgul & Anil, 2003; Soekamto *et al.*, 2003). As a part of our ongoing studies of the effect of side chain substituents on the solid state structures of 3-ethylsulfinyl-2-(4-fluorophenyl)-5-halo-1-benzofuran analogues (Choi *et al.*, 2010*a,b*), we report the crystal structure of the title compound (Fig. 1).

The benzofuran unit is essentially planar, with a mean deviation of 0.012 (2) Å from the least-squares plane defined by the nine constituent atoms. The dihedral angle formed by the benzofuran plane and the 4-fluorophenyl ring is 15.60 (6)°. The crystal packing (Fig. 2) is stabilized by an I···O halogen bond between the iodine and the oxygen of the S=O unit [$C4-I\cdots O2^i = 3.052(2)$ Å; $C4-I\cdots O2^i = 168.75(7)^\circ$] (Politzer *et al.*, 2007).

Experimental

77% 3-Chloroperoxybenzoic acid (202 mg, 0.9 mmol) was added in small portions to a stirred solution of 3-ethylsulfinyl-2-(4-fluorophenyl)-5-iodo-7-methyl-1-benzofuran (330 mg, 0.8 mmol) in dichloromethane (25 mL) at 273 K. After being stirred at room temperature for 4h, the mixture was washed with saturated sodium bicarbonate solution and the organic layer was separated, dried over magnesium sulfate, filtered and concentrated at reduced pressure. The residue was purified by column chromatography (hexane–ethyl acetate, 2:1 v/v) to afford the title compound as a colorless solid [yield 82%, m.p. 466–467 K; $R_f = 0.54$ (hexane–ethyl acetate, 2:1 v/v)]. Single crystals suitable for X-ray diffraction were prepared by slow evaporation of a solution of the title compound in acetone at room temperature.

Refinement

All H atoms were positioned geometrically and refined using a riding model, with C—H = 0.93 Å for aryl, 0.97 Å for methylene, and 0.96 Å for methyl H atoms. $U_{iso}(H) = 1.2U_{eq}(C)$ for aryl and methylene, and $1.5U_{eq}(C)$ for methyl H atoms.

Figures




Fig. 1. The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 50 % probability level. H atoms are presented as a small spheres of arbitrary radius.




Fig. 2. C—I...O interactions (dotted lines) in the crystal structure of the title compound. [Symmetry codes: i) $-x + 1, -y + 1, -z + 2.$]

3-Ethylsulfinyl-2-(4-fluorophenyl)-5-iodo-7-methyl-1-benzofuran

Crystal data

$C_{17}H_{14}FIO_2S$	$Z = 2$
$M_r = 428.24$	$F(000) = 420$
Triclinic, $P\bar{1}$	$D_x = 1.799 \text{ Mg m}^{-3}$
Hall symbol: $-P 1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 7.4681 (5) \text{ \AA}$	Cell parameters from 6089 reflections
$b = 10.3023 (7) \text{ \AA}$	$\theta = 2.1\text{--}27.4^\circ$
$c = 10.9761 (7) \text{ \AA}$	$\mu = 2.17 \text{ mm}^{-1}$
$\alpha = 76.289 (1)^\circ$	$T = 273 \text{ K}$
$\beta = 89.138 (1)^\circ$	Block, colourless
$\gamma = 74.802 (1)^\circ$	$0.50 \times 0.25 \times 0.25 \text{ mm}$
$V = 790.73 (9) \text{ \AA}^3$	

Data collection

Bruker SMART APEXII CCD diffractometer	3386 independent reflections
Radiation source: Rotating Anode	3261 reflections with $I > 2\sigma(I)$
Bruker HELIOS graded multilayer optics	$R_{\text{int}} = 0.049$
Detector resolution: $10.0 \text{ pixels mm}^{-1}$	$\theta_{\text{max}} = 27.0^\circ, \theta_{\text{min}} = 2.8^\circ$
φ and ω scans	$h = -9 \rightarrow 9$
Absorption correction: multi-scan (SADABS; Bruker, 2009)	$k = -13 \rightarrow 13$
$T_{\text{min}} = 0.520, T_{\text{max}} = 0.582$	$l = -14 \rightarrow 13$
6867 measured reflections	

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.022$	Hydrogen site location: difference Fourier map
$wR(F^2) = 0.059$	H-atom parameters constrained
$S = 1.06$	$w = 1/[\sigma^2(F_o^2) + (0.0284P)^2 + 0.4222P]$
3386 reflections	where $P = (F_o^2 + 2F_c^2)/3$
201 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.71 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.81 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
I	0.542549 (19)	0.247310 (13)	1.062688 (11)	0.02504 (6)
F	-0.0461 (2)	0.7492 (2)	-0.01911 (13)	0.0441 (4)
O1	0.2937 (2)	0.37876 (15)	0.50304 (13)	0.0214 (3)
O2	0.3809 (2)	0.75140 (17)	0.66534 (15)	0.0284 (3)
C1	0.2629 (3)	0.5691 (2)	0.57785 (18)	0.0204 (4)
C2	0.3398 (3)	0.4484 (2)	0.67903 (19)	0.0201 (4)
C3	0.3934 (3)	0.4255 (2)	0.80613 (18)	0.0215 (4)
H3	0.3824	0.4982	0.8447	0.026*
C4	0.4640 (3)	0.2880 (2)	0.87108 (18)	0.0215 (4)
C5	0.4839 (3)	0.1760 (2)	0.8147 (2)	0.0243 (4)
H5	0.5346	0.0861	0.8620	0.029*
C6	0.4289 (3)	0.1976 (2)	0.68911 (19)	0.0217 (4)
C7	0.3574 (3)	0.3354 (2)	0.62726 (18)	0.0205 (4)
C8	0.2388 (3)	0.5210 (2)	0.47433 (19)	0.0205 (4)
C9	0.1667 (3)	0.5850 (2)	0.34449 (19)	0.0219 (4)
C10	0.0601 (3)	0.7217 (3)	0.3068 (2)	0.0284 (5)
H10	0.0353	0.7757	0.3652	0.034*
C11	-0.0100 (3)	0.7790 (3)	0.1840 (2)	0.0313 (5)
H11	-0.0791	0.8710	0.1587	0.038*
C12	0.0261 (3)	0.6952 (3)	0.1005 (2)	0.0298 (5)
C13	0.1322 (3)	0.5599 (3)	0.1329 (2)	0.0303 (5)
H13	0.1545	0.5065	0.0740	0.036*
C14	0.2052 (3)	0.5048 (2)	0.2553 (2)	0.0259 (4)
H14	0.2799	0.4142	0.2785	0.031*
C15	0.4396 (4)	0.0826 (2)	0.6251 (2)	0.0295 (5)
H15A	0.3163	0.0779	0.6072	0.044*
H15B	0.5075	-0.0038	0.6791	0.044*
H15C	0.5018	0.0999	0.5482	0.044*
C16	0.0218 (3)	0.7605 (2)	0.6731 (2)	0.0279 (4)
H16A	-0.0783	0.7464	0.6269	0.034*
H16B	0.0467	0.6908	0.7520	0.034*
C17	-0.0346 (4)	0.9043 (3)	0.6981 (2)	0.0383 (6)
H17A	0.0664	0.9186	0.7417	0.057*

supplementary materials

H17B	-0.1411	0.9124	0.7486	0.057*
H17C	-0.0646	0.9727	0.6198	0.057*
S1	0.22856 (8)	0.74390 (5)	0.58266 (4)	0.02169 (11)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
I	0.03433 (10)	0.02416 (9)	0.01561 (8)	-0.00848 (6)	-0.00381 (5)	-0.00178 (5)
F	0.0453 (9)	0.0628 (11)	0.0171 (6)	-0.0101 (8)	-0.0098 (6)	-0.0001 (6)
O1	0.0293 (7)	0.0200 (7)	0.0156 (6)	-0.0069 (6)	-0.0013 (5)	-0.0048 (5)
O2	0.0352 (9)	0.0272 (8)	0.0266 (8)	-0.0114 (7)	-0.0022 (6)	-0.0101 (6)
C1	0.0250 (10)	0.0188 (9)	0.0174 (9)	-0.0055 (8)	-0.0002 (7)	-0.0046 (7)
C2	0.0234 (10)	0.0188 (9)	0.0181 (9)	-0.0056 (7)	0.0007 (7)	-0.0046 (7)
C3	0.0294 (10)	0.0200 (9)	0.0166 (9)	-0.0077 (8)	-0.0010 (7)	-0.0057 (7)
C4	0.0274 (10)	0.0222 (10)	0.0152 (8)	-0.0077 (8)	-0.0017 (7)	-0.0035 (7)
C5	0.0302 (11)	0.0197 (10)	0.0219 (10)	-0.0064 (8)	-0.0010 (8)	-0.0029 (8)
C6	0.0276 (10)	0.0193 (10)	0.0200 (9)	-0.0077 (8)	0.0017 (8)	-0.0065 (7)
C7	0.0244 (10)	0.0231 (10)	0.0158 (9)	-0.0089 (8)	0.0006 (7)	-0.0052 (7)
C8	0.0230 (10)	0.0196 (9)	0.0194 (9)	-0.0066 (8)	0.0009 (7)	-0.0042 (7)
C9	0.0245 (10)	0.0268 (10)	0.0161 (9)	-0.0107 (8)	0.0004 (7)	-0.0044 (8)
C10	0.0329 (12)	0.0287 (11)	0.0216 (10)	-0.0056 (9)	-0.0029 (8)	-0.0052 (8)
C11	0.0320 (12)	0.0311 (12)	0.0249 (11)	-0.0049 (10)	-0.0045 (9)	0.0011 (9)
C12	0.0272 (11)	0.0448 (14)	0.0164 (9)	-0.0126 (10)	-0.0034 (8)	-0.0019 (9)
C13	0.0338 (12)	0.0415 (13)	0.0197 (10)	-0.0137 (10)	0.0013 (8)	-0.0111 (9)
C14	0.0276 (11)	0.0303 (11)	0.0211 (10)	-0.0085 (9)	-0.0005 (8)	-0.0078 (8)
C15	0.0445 (13)	0.0200 (10)	0.0263 (10)	-0.0089 (9)	0.0002 (9)	-0.0092 (8)
C16	0.0291 (11)	0.0263 (11)	0.0258 (10)	-0.0035 (9)	0.0021 (8)	-0.0055 (8)
C17	0.0466 (15)	0.0275 (12)	0.0331 (12)	0.0036 (11)	0.0038 (11)	-0.0076 (10)
S1	0.0315 (3)	0.0166 (2)	0.0164 (2)	-0.00631 (19)	-0.00063 (18)	-0.00285 (17)

Geometric parameters (\AA , $^\circ$)

I—C4	2.106 (2)	C9—C14	1.405 (3)
I—O2 ⁱ	3.052 (2)	C10—C11	1.387 (3)
F—C12	1.357 (2)	C10—H10	0.9300
O1—C8	1.373 (2)	C11—C12	1.380 (4)
O1—C7	1.380 (2)	C11—H11	0.9300
O2—S1	1.495 (2)	C12—C13	1.376 (4)
C1—C8	1.374 (3)	C13—C14	1.387 (3)
C1—C2	1.449 (3)	C13—H13	0.9300
C1—S1	1.765 (2)	C14—H14	0.9300
C2—C7	1.389 (3)	C15—H15A	0.9600
C2—C3	1.405 (3)	C15—H15B	0.9600
C3—C4	1.391 (3)	C15—H15C	0.9600
C3—H3	0.9300	C16—C17	1.520 (3)
C4—C5	1.407 (3)	C16—S1	1.815 (2)
C5—C6	1.394 (3)	C16—H16A	0.9700
C5—H5	0.9300	C16—H16B	0.9700

C6—C7	1.385 (3)	C17—H17A	0.9600
C6—C15	1.499 (3)	C17—H17B	0.9600
C8—C9	1.465 (3)	C17—H17C	0.9600
C9—C10	1.391 (3)		
C4—I—O2 ⁱ	168.75 (7)	C12—C11—H11	121.1
C8—O1—C7	106.96 (15)	C10—C11—H11	121.1
C8—C1—C2	106.76 (18)	F—C12—C13	118.6 (2)
C8—C1—S1	126.56 (16)	F—C12—C11	118.3 (2)
C2—C1—S1	126.41 (15)	C13—C12—C11	123.0 (2)
C7—C2—C3	119.10 (19)	C12—C13—C14	118.5 (2)
C7—C2—C1	105.34 (17)	C12—C13—H13	120.7
C3—C2—C1	135.56 (19)	C14—C13—H13	120.7
C4—C3—C2	116.33 (18)	C13—C14—C9	120.4 (2)
C4—C3—H3	121.8	C13—C14—H14	119.8
C2—C3—H3	121.8	C9—C14—H14	119.8
C3—C4—C5	123.00 (18)	C6—C15—H15A	109.5
C3—C4—I	117.95 (15)	C6—C15—H15B	109.5
C5—C4—I	119.06 (15)	H15A—C15—H15B	109.5
C6—C5—C4	121.11 (19)	C6—C15—H15C	109.5
C6—C5—H5	119.4	H15A—C15—H15C	109.5
C4—C5—H5	119.4	H15B—C15—H15C	109.5
C7—C6—C5	114.58 (18)	C17—C16—S1	109.14 (18)
C7—C6—C15	121.77 (18)	C17—C16—H16A	109.9
C5—C6—C15	123.63 (19)	S1—C16—H16A	109.9
O1—C7—C6	123.70 (18)	C17—C16—H16B	109.9
O1—C7—C2	110.45 (18)	S1—C16—H16B	109.9
C6—C7—C2	125.85 (18)	H16A—C16—H16B	108.3
O1—C8—C1	110.48 (17)	C16—C17—H17A	109.5
O1—C8—C9	114.21 (17)	C16—C17—H17B	109.5
C1—C8—C9	135.31 (19)	H17A—C17—H17B	109.5
C10—C9—C14	118.92 (19)	C16—C17—H17C	109.5
C10—C9—C8	122.32 (19)	H17A—C17—H17C	109.5
C14—C9—C8	118.8 (2)	H17B—C17—H17C	109.5
C11—C10—C9	121.3 (2)	O2—S1—C1	107.96 (10)
C11—C10—H10	119.4	O2—S1—C16	106.52 (10)
C9—C10—H10	119.4	C1—S1—C16	97.83 (11)
C12—C11—C10	117.9 (2)		
C8—C1—C2—C7	0.4 (2)	C2—C1—C8—O1	0.4 (2)
S1—C1—C2—C7	-173.92 (16)	S1—C1—C8—O1	174.74 (15)
C8—C1—C2—C3	-179.2 (2)	C2—C1—C8—C9	179.1 (2)
S1—C1—C2—C3	6.5 (4)	S1—C1—C8—C9	-6.6 (4)
C7—C2—C3—C4	0.9 (3)	O1—C8—C9—C10	162.3 (2)
C1—C2—C3—C4	-179.6 (2)	C1—C8—C9—C10	-16.3 (4)
C2—C3—C4—C5	0.6 (3)	O1—C8—C9—C14	-17.0 (3)
C2—C3—C4—I	-179.02 (15)	C1—C8—C9—C14	164.4 (2)
C3—C4—C5—C6	-1.5 (4)	C14—C9—C10—C11	0.7 (4)
I—C4—C5—C6	178.17 (17)	C8—C9—C10—C11	-178.6 (2)
C4—C5—C6—C7	0.7 (3)	C9—C10—C11—C12	1.3 (4)

supplementary materials

C4—C5—C6—C15	-177.5 (2)	C10—C11—C12—F	178.2 (2)
C8—O1—C7—C6	-178.4 (2)	C10—C11—C12—C13	-2.0 (4)
C8—O1—C7—C2	1.4 (2)	F—C12—C13—C14	-179.7 (2)
C5—C6—C7—O1	-179.43 (19)	C11—C12—C13—C14	0.5 (4)
C15—C6—C7—O1	-1.2 (3)	C12—C13—C14—C9	1.7 (4)
C5—C6—C7—C2	0.9 (3)	C10—C9—C14—C13	-2.3 (3)
C15—C6—C7—C2	179.1 (2)	C8—C9—C14—C13	177.1 (2)
C3—C2—C7—O1	178.56 (18)	C8—C1—S1—O2	-139.72 (19)
C1—C2—C7—O1	-1.1 (2)	C2—C1—S1—O2	33.5 (2)
C3—C2—C7—C6	-1.7 (3)	C8—C1—S1—C16	110.0 (2)
C1—C2—C7—C6	178.6 (2)	C2—C1—S1—C16	-76.8 (2)
C7—O1—C8—C1	-1.1 (2)	C17—C16—S1—O2	65.53 (18)
C7—O1—C8—C9	179.96 (18)	C17—C16—S1—C1	176.96 (16)

Symmetry codes: (i) $-x+1, -y+1, -z+2$.

Fig. 1




Fig. 2

