

(E)-N'-(3,5-Dibromo-2-hydroxybenzylidene)nicotinohydrazide

Yong-Qing Su,* Cong Li and Ping Wang

Faculty of Chemistry and Chemical Engineering, Yunnan Normal University, Kunming 650092, People's Republic of China
Correspondence e-mail: yongqing_su@163.com


Received 11 February 2010; accepted 17 February 2010

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.038; wR factor = 0.111; data-to-parameter ratio = 16.3.

In the title Schiff base compound, $\text{C}_{13}\text{H}_9\text{Br}_2\text{N}_3\text{O}_2$, there is an intramolecular $\text{O}-\text{H}\cdots\text{N}$ hydrogen bond involving the hydroxyl substituent and the adjacent hydrazine N atom. The molecule is almost planar, the dihedral angle between the benzene ring and the pyridine ring being $5.7(2)^\circ$. In the crystal structure, symmetry-related molecules are linked via $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, forming chains propagating in [001].

Related literature

For related literature on Schiff bases, see: Archibald *et al.* (1994); Harada *et al.* (1999); Ogawa *et al.* (1998). For similar structures, see: Mohd Lair *et al.* (2009); Li *et al.* (2010); Sun *et al.* (2009); Wang *et al.* (2010); Wen *et al.* (2009).



Experimental

Crystal data

$\text{C}_{13}\text{H}_9\text{Br}_2\text{N}_3\text{O}_2$

$M_r = 399.05$

Monoclinic, $P2_1/c$

$a = 17.013(4)\text{ \AA}$

$b = 8.091(2)\text{ \AA}$

$c = 10.153(3)\text{ \AA}$

$\beta = 92.194(13)^\circ$

$V = 1396.6(6)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 5.81\text{ mm}^{-1}$

$T = 298\text{ K}$

$0.23 \times 0.21 \times 0.20\text{ mm}$

Data collection

Bruker APEXII CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 2004)
 $T_{\min} = 0.349$, $T_{\max} = 0.390$

8024 measured reflections
3023 independent reflections
1933 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.111$
 $S = 1.00$
3023 reflections
185 parameters
1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.64\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.69\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1 \cdots N1	0.82	1.89	2.609 (4)	146
N2—H2 \cdots O2 ⁱ	0.91 (2)	2.14 (2)	3.017 (4)	162 (3)

Symmetry code: (i) $x, -y + \frac{3}{2}, z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors would like to acknowledge the support provided by the Yunnan Province Science and Technology Department, the Program of Yunnan Province's Young-middle aged Reserve Scientific and Technological Principal Culture Plan (grant No. 2006PY01-50) and the fund of Yunnan Province's Applied Research Plan (grant No. 2006E0032M).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2163).

References

- Archibald, S. J., Blake, A. J., Schroder, M. & Winpenny, R. E. P. (1994). *Chem. Commun.* pp. 1669–1670.
Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Harada, J., Uekusa, H. & Ohashi, Y. (1999). *J. Am. Chem. Soc.* **121**, 5809–5810.
Li, C., Wang, P. & Su, Y.-Q. (2010). *Acta Cryst. E* **66**, o520.
Mohd Lair, N., Mohd Ali, H. & Ng, S. W. (2009). *Acta Cryst. E* **65**, o189.
Ogawa, K., Kasahara, Y., Ohtani, Y. & Harada, J. (1998). *J. Am. Chem. Soc.* **120**, 7107–7108.
Sheldrick, G. M. (2004). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Sun, Y., Li, H.-G., Wang, X., Fu, S. & Wang, D. (2009). *Acta Cryst. E* **65**, o262.
Wang, P., Li, C. & Su, Y.-Q. (2010). *Acta Cryst. E* **66**, o542.
Wen, L., Yin, H., Li, W. & Li, K. (2009). *Acta Cryst. E* **65**, o2623.

supplementary materials

Acta Cryst. (2010). E66, o670 [doi:10.1107/S1600536810006276]

(E)-N¹-(3,5-Dibromo-2-hydroxybenzylidene)nicotinohydrazide

Y.-Q. Su, C. Li and P. Wang

Comment

Schiff bases have received much attention in recent years (Ogawa *et al.*, 1998; Archibald *et al.*, 1994; Harada *et al.*, 1999). Recently, we reported on the crystal structures of two new Schiff bases (Li *et al.*, 2010; Wang *et al.*, 2010). As a further investigation of the structures of Schiff base compounds the title compound was prepared by the reaction of 3,5-dibromo-2-hydroxybenzaldehyde with nicotinic acid hydrazide in methanol; its crystal structure is reported on here.

In the title compound (Fig. 1), there is an intramolecular O-H···N hydrogen bond involving the hydroxyl substituent and the adjacent hydrazine N-atom, N2 (Table 1). The benzene ring is inclined to the pyridine ring by 5.7 (2) $^{\circ}$. All the bond lengths are comparable with those observed in similar Schiff bases reported on previously (Wen *et al.*, 2009; Mohd Lair *et al.*, 2009; Sun *et al.*, 2009).

In the crystal structure, symmetry related molecules are linked via intermolecular N—H···O hydrogen bonds so forming chains running along the *c* axis (Table 1, Fig. 2).

Experimental

3,5-Dibromo-2-hydroxybenzaldehyde (1.0 mmol, 280 mg) and nicotinic acid hydrazide (1.0 mmol, 137 mg) were dissolved in methanol (30 mL). The mixture was stirred at room temperature for about 1 h to give a colorless solution. After allowing the solution to evaporate slowly in air for 8 days, colorless block-like crystals, suitable for X-ray analysis, were formed.

Refinement

Atom H2 was located in a difference Fourier map and refined with a N—H distance restraint of 0.90 (1) Å and $U_{\text{iso}}(\text{H}) = 0.08$. The other H atoms were placed in idealized positions and constrained to ride on their parent atoms: O—H = 0.82 Å, C—H = 0.93 Å, with $U_{\text{iso}}(\text{H}) = k \times U_{\text{eq}}(\text{parent atom})$, where $k = 1.2$ for C-bound H-atoms and $= 1.5$ for the hydroxyl H-atom.

Figures




Fig. 1. The molecular structure of the title compound, showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. The intramolecular O-H···N hydrogen bond is shown as a dashed line.

supplementary materials




Fig. 2. The crystal packing of the title compound viewed along the b -axis. The intra- and intermolecular hydrogen bonds are shown as dashed lines (see Table 1 for details).

(E)-N¹-(3,5-Dibromo-2-hydroxybenzylidene)nicotinohydrazide

Crystal data

C ₁₃ H ₉ Br ₂ N ₃ O ₂	$F(000) = 776$
$M_r = 399.05$	$D_x = 1.898 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 2084 reflections
$a = 17.013 (4) \text{ \AA}$	$\theta = 2.4\text{--}25.2^\circ$
$b = 8.091 (2) \text{ \AA}$	$\mu = 5.81 \text{ mm}^{-1}$
$c = 10.153 (3) \text{ \AA}$	$T = 298 \text{ K}$
$\beta = 92.194 (13)^\circ$	Block, colourless
$V = 1396.6 (6) \text{ \AA}^3$	$0.23 \times 0.21 \times 0.20 \text{ mm}$
$Z = 4$	

Data collection

Bruker APEXII CCD area-detector diffractometer	3023 independent reflections
Radiation source: fine-focus sealed tube graphite	1933 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.036$
Absorption correction: multi-scan (SADABS; Sheldrick, 2004)	$\theta_{\text{max}} = 27.0^\circ, \theta_{\text{min}} = 1.2^\circ$
$T_{\text{min}} = 0.349, T_{\text{max}} = 0.390$	$h = -19 \rightarrow 21$
8024 measured reflections	$k = -10 \rightarrow 10$
	$l = -12 \rightarrow 8$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.038$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.111$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.00$	$w = 1/[\sigma^2(F_o^2) + (0.0589P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$

3023 reflections	$(\Delta/\sigma)_{\max} = 0.001$
185 parameters	$\Delta\rho_{\max} = 0.64 \text{ e \AA}^{-3}$
1 restraint	$\Delta\rho_{\min} = -0.69 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.38611 (2)	1.15620 (7)	0.38681 (5)	0.0731 (2)
Br2	0.44353 (2)	0.85756 (6)	0.88313 (4)	0.06341 (18)
N1	0.11733 (17)	0.8462 (3)	0.5380 (3)	0.0393 (7)
N2	0.04257 (16)	0.7852 (4)	0.5483 (3)	0.0402 (7)
N3	-0.16645 (18)	0.5160 (4)	0.5780 (3)	0.0501 (8)
O1	0.22622 (14)	1.0061 (3)	0.4163 (2)	0.0514 (7)
H1	0.1820	0.9696	0.4270	0.077*
O2	0.01730 (14)	0.8169 (3)	0.3293 (2)	0.0508 (7)
C1	0.2459 (2)	0.8814 (4)	0.6310 (3)	0.0383 (8)
C2	0.27257 (19)	0.9725 (4)	0.5231 (3)	0.0377 (8)
C3	0.3494 (2)	1.0292 (4)	0.5283 (3)	0.0437 (9)
C4	0.4001 (2)	0.9964 (4)	0.6347 (3)	0.0455 (9)
H4	0.4512	1.0374	0.6373	0.055*
C5	0.3737 (2)	0.9026 (4)	0.7363 (3)	0.0403 (8)
C6	0.2980 (2)	0.8450 (4)	0.7371 (3)	0.0430 (9)
H6	0.2813	0.7823	0.8076	0.052*
C7	0.1655 (2)	0.8224 (4)	0.6360 (3)	0.0396 (8)
H7	0.1490	0.7673	0.7104	0.048*
C8	-0.00388 (19)	0.7712 (4)	0.4372 (3)	0.0366 (8)
C9	-0.08166 (19)	0.6936 (4)	0.4557 (3)	0.0358 (8)
C10	-0.1424 (2)	0.7215 (5)	0.3642 (4)	0.0492 (9)
H10	-0.1347	0.7889	0.2917	0.059*
C11	-0.2145 (2)	0.6488 (5)	0.3811 (4)	0.0541 (11)
H11	-0.2561	0.6666	0.3207	0.065*
C12	-0.2236 (2)	0.5493 (5)	0.4894 (4)	0.0510 (10)
H12	-0.2727	0.5025	0.5011	0.061*
C13	-0.0972 (2)	0.5869 (4)	0.5595 (3)	0.0417 (9)
H13	-0.0563	0.5637	0.6200	0.050*
H2	0.024 (2)	0.764 (6)	0.629 (2)	0.080*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0414 (3)	0.1045 (4)	0.0730 (3)	-0.0128 (2)	-0.0014 (2)	0.0354 (3)
Br2	0.0494 (3)	0.0779 (3)	0.0609 (3)	0.0041 (2)	-0.0244 (2)	0.0057 (2)
N1	0.0304 (16)	0.0528 (18)	0.0347 (16)	-0.0058 (13)	0.0011 (12)	-0.0056 (13)
N2	0.0285 (16)	0.0596 (19)	0.0324 (16)	-0.0048 (14)	-0.0016 (12)	-0.0034 (14)
N3	0.0425 (19)	0.057 (2)	0.0510 (19)	-0.0091 (15)	0.0009 (15)	0.0027 (15)
O1	0.0355 (15)	0.079 (2)	0.0393 (14)	-0.0054 (13)	-0.0059 (11)	0.0100 (13)
O2	0.0404 (15)	0.082 (2)	0.0301 (14)	-0.0050 (13)	-0.0007 (11)	0.0079 (13)
C1	0.0308 (19)	0.044 (2)	0.040 (2)	-0.0013 (15)	-0.0025 (15)	-0.0067 (16)
C2	0.0319 (19)	0.047 (2)	0.0344 (18)	0.0030 (15)	-0.0022 (14)	-0.0012 (16)
C3	0.035 (2)	0.051 (2)	0.045 (2)	-0.0012 (16)	-0.0013 (16)	0.0044 (17)
C4	0.030 (2)	0.051 (2)	0.055 (2)	-0.0002 (17)	-0.0051 (17)	-0.0015 (19)
C5	0.036 (2)	0.044 (2)	0.041 (2)	0.0035 (16)	-0.0078 (15)	-0.0012 (17)
C6	0.043 (2)	0.047 (2)	0.039 (2)	0.0002 (17)	-0.0033 (16)	-0.0008 (16)
C7	0.037 (2)	0.044 (2)	0.037 (2)	-0.0012 (15)	-0.0033 (16)	-0.0010 (15)
C8	0.0315 (19)	0.043 (2)	0.0347 (19)	0.0033 (15)	-0.0029 (15)	-0.0034 (16)
C9	0.0322 (19)	0.043 (2)	0.0318 (18)	0.0028 (15)	-0.0015 (14)	-0.0087 (15)
C10	0.042 (2)	0.066 (3)	0.038 (2)	-0.0023 (19)	-0.0067 (17)	0.0071 (19)
C11	0.034 (2)	0.077 (3)	0.050 (2)	0.0003 (19)	-0.0100 (17)	-0.011 (2)
C12	0.037 (2)	0.056 (2)	0.061 (3)	-0.0079 (18)	0.0017 (18)	-0.012 (2)
C13	0.038 (2)	0.047 (2)	0.039 (2)	-0.0013 (17)	-0.0042 (16)	0.0019 (16)

Geometric parameters (\AA , $^\circ$)

Br1—C3	1.892 (4)	C3—C4	1.382 (4)
Br2—C5	1.905 (3)	C4—C5	1.370 (5)
N1—C7	1.278 (4)	C4—H4	0.9300
N1—N2	1.372 (4)	C5—C6	1.371 (5)
N2—C8	1.357 (4)	C6—H6	0.9300
N2—H2	0.901 (10)	C7—H7	0.9300
N3—C12	1.326 (4)	C8—C9	1.483 (5)
N3—C13	1.330 (4)	C9—C10	1.381 (5)
O1—C2	1.344 (4)	C9—C13	1.396 (5)
O1—H1	0.8200	C10—C11	1.377 (5)
O2—C8	1.223 (4)	C10—H10	0.9300
C1—C6	1.401 (5)	C11—C12	1.376 (5)
C1—C2	1.409 (5)	C11—H11	0.9300
C1—C7	1.452 (5)	C12—H12	0.9300
C2—C3	1.384 (5)	C13—H13	0.9300
C7—N1—N2	117.1 (3)	C1—C6—H6	120.3
C8—N2—N1	118.6 (3)	N1—C7—C1	120.0 (3)
C8—N2—H2	122 (3)	N1—C7—H7	120.0
N1—N2—H2	119 (3)	C1—C7—H7	120.0
C12—N3—C13	116.5 (3)	O2—C8—N2	122.5 (3)
C2—O1—H1	109.5	O2—C8—C9	122.4 (3)

C6—C1—C2	119.6 (3)	N2—C8—C9	115.1 (3)
C6—C1—C7	118.3 (3)	C10—C9—C13	116.7 (3)
C2—C1—C7	122.1 (3)	C10—C9—C8	119.6 (3)
O1—C2—C3	119.2 (3)	C13—C9—C8	123.6 (3)
O1—C2—C1	122.4 (3)	C11—C10—C9	119.6 (4)
C3—C2—C1	118.4 (3)	C11—C10—H10	120.2
C4—C3—C2	121.7 (3)	C9—C10—H10	120.2
C4—C3—Br1	118.9 (3)	C12—C11—C10	118.5 (4)
C2—C3—Br1	119.4 (3)	C12—C11—H11	120.8
C5—C4—C3	118.9 (3)	C10—C11—H11	120.8
C5—C4—H4	120.6	N3—C12—C11	124.0 (4)
C3—C4—H4	120.6	N3—C12—H12	118.0
C4—C5—C6	121.9 (3)	C11—C12—H12	118.0
C4—C5—Br2	118.9 (3)	N3—C13—C9	124.6 (3)
C6—C5—Br2	119.2 (3)	N3—C13—H13	117.7
C5—C6—C1	119.4 (3)	C9—C13—H13	117.7
C5—C6—H6	120.3		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O1—H1···N1	0.82	1.89	2.609 (4)	146
N2—H2···O2 ⁱ	0.91 (2)	2.14 (2)	3.017 (4)	162 (3)

Symmetry codes: (i) $x, -y+3/2, z+1/2$.

supplementary materials

Fig. 1




Fig. 2

