

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-Chloroquinoline-3-carboxylic acid

Souheila Ladraa,^a Abdelmalek Bouraiou,^{a*} Sofiane Bouacida,^b Thierry Roisnel^c and Ali Belfaitah^a

^aLaboratoire des Produits Naturels d'Origine Végétale et de Synthèse Organique, PHYSYNOR, Université Mentouri-Constantine, 25000 Constantine, Algeria, ^bUnité de Recherche de Chimie de l'Environnement et Moléculaire Structurale, CHEMS, Université Mentouri-Constantine, 25000 Algeria, and ^cCentre de diffractométrie X, UMR 6226 CNRS Unité Sciences Chimiques de Rennes, Université de Rennes I, 263 Avenue du Général Leclerc, 35042 Rennes, France
Correspondence e-mail: bouraiou.abdelmalek@yahoo.fr


Received 17 February 2010; accepted 19 February 2010

Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.038; wR factor = 0.094; data-to-parameter ratio = 15.0.

The crystal structure of the title compound, $\text{C}_{10}\text{H}_6\text{ClNO}_2$, can be described by two types of crossed layers which are parallel to (110) and $(\bar{1}10)$. The crystal packing is stabilized by intermolecular $\text{C}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{N}$ hydrogen bonds, resulting in the formation of a two-dimensional network and reinforcing the cohesion of the structure.

Related literature

For our previous work on the preparation of α -aminonitriles, see: Ladraa *et al.* (2009); Belfaitah *et al.* (2006). For the removal of chiral auxiliaries using ceric ammonium nitrate, see: Bhanu Prasad *et al.* (2004).



Experimental

Crystal data

 $\text{C}_{10}\text{H}_6\text{ClNO}_2$ $M_r = 207.61$ Orthorhombic, $P2_1nb$ $a = 5.8193$ (2) Å $b = 8.0689$ (3) Å $c = 18.1780$ (5) Å $V = 853.55$ (5) Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.41$ mm⁻¹ $T = 120$ K $0.19 \times 0.12 \times 0.08$ mm

Data collection

Nonius KappaCCD diffractometer

Absorption correction: multi-scan

(SADABS; Sheldrick, 2002)

 $T_{\min} = 0.915$, $T_{\max} = 0.967$

13714 measured reflections

1938 independent reflections

1746 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.046$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.038$ $wR(F^2) = 0.094$ $S = 1.14$

1938 reflections

129 parameters

1 restraint

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.32$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.35$ e Å⁻³

Absolute structure: Flack (1983),

862 Friedel pairs

Flack parameter: 0.28 (9)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O2}-\text{H2}\cdots\text{N1}^{\text{i}}$	0.84	1.95	2.768 (3)	164
$\text{C8}-\text{H8}\cdots\text{O1}^{\text{ii}}$	0.95	2.37	3.290 (4)	163

Symmetry codes: (i) $x - 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *COLLECT* (Nonius, 1998); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* (Otwinowski & Minor 1997) and *SCALEPACK*; program(s) used to solve structure: *SIR2002* (Burla *et al.*, 2003); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *DIAMOND* (Brandenburg & Berndt, 2001); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

We are grateful to all personnel of the laboratory PHYSYNOR, Université Mentouri-Constantine, Algérie for their assistance. Thanks are due to MESRS (Ministère de l'Enseignement Supérieur et de la Recherche Scientifique – Algérie) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2197).

References

- Belfaitah, A., Ladraa, S., Bouraiou, A., Benali-Cherif, N., Debache, A. & Rhouati, S. (2006). *Acta Cryst.* **E62**, o1355–o1357.
 Bhanu Prasad, B. A., Bisai, A. & Singh, V. K. (2004). *Tetrahedron Lett.* **45**, 9565–9567.
 Brandenburg, K. & Berndt, M. (2001). *DIAMOND*. Crystal Impact, Bonn, Germany.
 Burla, M. C., Camalli, M., Carrozzini, B., Cascarano, G. L., Giacovazzo, C., Polidori, G. & Spagna, R. (2003). *J. Appl. Cryst.* **36**, 1103.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Ladraa, S., Bouraiou, A., Bouacida, S., Roisnel, T. & Belfaitah, A. (2009). *Acta Cryst.* **C65**, o475–o478.
 Nonius (1998). *COLLECT*. Nonius BV, Delft The Netherlands.
 Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
 Sheldrick, G. M. (2002). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2010). E66, o693 [doi:10.1107/S1600536810006501]

2-Chloroquinoline-3-carboxylic acid

S. Ladraa, A. Bouraiou, S. Bouacida, T. Roisnel and A. Belfaitah

Comment

In a continuation of our previous work related to the preparation of α -aminonitrile (Ladraa *et al.*, 2009; Belfaitah *et al.*, 2006) and in order to prepare chiral N-protected α -aminonitrile, we have explored the oxidative debenylation of N-protected α -aminonitrile. The removal of the chiral auxiliaries has already been investigated using ceric ammonium nitrate (CAN) (Bhanu Prasad *et al.*, 2004). Surprisingly, our attempts to remove the chiral auxiliary using CAN were failed to undergo the desired adduct and led to the 2-chloroquinoline-3-carboxylic acid (I). This unexpected cleavage of 2-[(*S*)-2-chloro-3-quinoly]-2-[(*R*)-1-(4-methoxyphenyl) ethylamino]acetonitrile may result from the decomposition of the α -aminonitrile into cyanide and imine which in turn undergo hydrolysis/oxidative sequence. In this paper, we report the structure determination of compound (I), resulting from unwanted decomposition of chiral N-protected α -aminonitrile.

The molecular geometry and the atom-numbering scheme of (I) are shown in Fig. 1. The two rings of quinolyl moiety are fused in an axial fashion and form a dihedral angle of $0.42(9)^\circ$. The crystal packing can be described by two types of crossed layers which quinolyl ring is parallel to (110) and (-110) planes respectively (Fig. 2). The crystal packing is stabilized by inter and intramolecular hydrogen bonds (O—H \cdots N and C—H \cdots O) linked molecules in the same layer, resulting in the formation of a two dimensional network and reinforcing a cohesion of structure. Hydrogen-bonding parameters are listed in table 1.

Experimental

A solution of 327 mg (3 eq., 0.59 mmol.) of ceric ammonium nitrate (CAN) in 1 ml of water was added to precooled stirred solution of 2-[(*S*)-2-chloro-3-quinoly]-2-[(*R*)-1-(4-methoxyphenyl) ethylamino] acetonitrile (70 mg, 0.19 mmol) in 9 ml of CH₃CN. After completion of the reaction (checked by TLC), the reaction mixture was poured into cold water and the residue obtained was filtered off. Crystals of (I) suitable for X-ray analysis were obtained by slow evaporation of the filtrate.

Refinement

All non-H atoms were refined with anisotropic atomic displacement parameters. All H atoms were localized on Fourier maps but introduced in calculated positions and treated as riding on their parent C and O atoms. (with C—H = 0.95 Å - O—H = 0.84 Å and $U_{\text{iso}}(\text{H}) = 1.2$ or 1.5 (carrier atom)).

Figures




Fig. 1. (Farrugia, 1997) the structure of the title compound with the atomic labeling scheme. Displacements are drawn at the 50% probability level.




Fig. 2. (Brandenburg & Berndt, 2001) A diagram of the layered crystal packing of (I) viewed down the c axis.




Fig. 3. (Brandenburg & Berndt, 2001) Unit cell of (I) showing hydrogen bond [O—H...N and C—H...O] as dashed line.

2-Chloroquinoline-3-carboxylic acid

Crystal data

$C_{10}H_6ClNO_2$

$M_r = 207.61$

Orthorhombic, $P2_1nb$

Hall symbol: P -2bc 2a

$a = 5.8193 (2) \text{ \AA}$

$b = 8.0689 (3) \text{ \AA}$

$c = 18.1780 (5) \text{ \AA}$

$V = 853.55 (5) \text{ \AA}^3$

$Z = 4$

$F(000) = 424$

$D_x = 1.616 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 21289 reflections

$\theta = 2.9\text{--}27.5^\circ$

$\mu = 0.41 \text{ mm}^{-1}$

$T = 120 \text{ K}$

Prism, yellow

$0.19 \times 0.12 \times 0.08 \text{ mm}$

Data collection

Nonius KappaCCD
diffractometer

Radiation source: Enraf–Nonius FR590
graphite

Detector resolution: 9 pixels mm^{-1}

CCD rotation images, thin slices scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 2002)

$T_{\min} = 0.915$, $T_{\max} = 0.967$

13714 measured reflections

1938 independent reflections

1746 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.046$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.4^\circ$

$h = -7 \rightarrow 7$

$k = -10 \rightarrow 10$

$l = -23 \rightarrow 23$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.038$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$wR(F^2) = 0.094$	$w = 1/[\sigma^2(F_o^2) + (0.0388P)^2 + 0.7398P]$
$S = 1.14$	where $P = (F_o^2 + 2F_c^2)/3$
1938 reflections	$(\Delta/\sigma)_{\max} = 0.001$
129 parameters	$\Delta\rho_{\max} = 0.32 \text{ e } \text{\AA}^{-3}$
1 restraint	$\Delta\rho_{\min} = -0.35 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 862 Friedel pairs Flack parameter: 0.28 (9)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.1466 (5)	0.1851 (3)	0.23620 (14)	0.0167 (5)
C2	-0.0521 (4)	0.0966 (3)	0.21287 (14)	0.0161 (5)
C3	-0.0939 (5)	0.0943 (3)	0.13814 (14)	0.0182 (5)
H3	-0.2255	0.0377	0.1201	0.022*
C4	0.0535 (5)	0.1735 (3)	0.08831 (15)	0.0175 (5)
C5	0.0162 (5)	0.1732 (3)	0.01110 (15)	0.0204 (6)
H5	-0.113	0.1173	-0.009	0.024*
C6	0.1668 (5)	0.2538 (3)	-0.03428 (14)	0.0196 (6)
H6	0.1416	0.254	-0.0859	0.024*
C7	0.3592 (5)	0.3363 (3)	-0.00485 (15)	0.0217 (6)
H7	0.4622	0.3918	-0.0371	0.026*
C8	0.4009 (5)	0.3382 (3)	0.06923 (15)	0.0200 (6)
H8	0.5315	0.3945	0.0882	0.024*
C9	0.2487 (5)	0.2563 (3)	0.11715 (13)	0.0164 (5)
C10	-0.2112 (4)	0.0090 (3)	0.26469 (13)	0.0168 (5)
N1	0.2901 (4)	0.2608 (3)	0.19237 (12)	0.0173 (5)
O1	-0.1908 (4)	0.0077 (3)	0.33080 (10)	0.0268 (5)
O2	-0.3754 (4)	-0.0713 (3)	0.22913 (11)	0.0256 (5)
H2	-0.4571	-0.1233	0.2594	0.038*
Cl1	0.22006 (13)	0.19998 (8)	0.32850 (3)	0.02526 (17)

Atomic displacement parameters (\AA^2)

U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
----------	----------	----------	----------	----------	----------

supplementary materials

C1	0.0190 (13)	0.0187 (13)	0.0124 (11)	0.0014 (10)	-0.0030 (9)	-0.0007 (10)
C2	0.0152 (13)	0.0183 (12)	0.0148 (13)	0.0012 (10)	0.0014 (9)	0.0008 (10)
C3	0.0137 (13)	0.0211 (13)	0.0198 (13)	-0.0030 (10)	0.0003 (11)	-0.0030 (10)
C4	0.0168 (13)	0.0180 (12)	0.0176 (12)	0.0023 (10)	0.0013 (10)	-0.0001 (10)
C5	0.0183 (14)	0.0233 (13)	0.0195 (13)	-0.0025 (11)	-0.0025 (11)	-0.0032 (11)
C6	0.0235 (17)	0.0227 (13)	0.0127 (12)	0.0034 (10)	-0.0010 (10)	-0.0001 (9)
C7	0.0221 (15)	0.0239 (14)	0.0191 (13)	-0.0008 (11)	0.0041 (11)	0.0029 (11)
C8	0.0174 (14)	0.0220 (13)	0.0208 (13)	0.0003 (11)	-0.0002 (10)	-0.0002 (11)
C9	0.0170 (14)	0.0169 (11)	0.0151 (11)	0.0006 (11)	-0.0019 (11)	-0.0008 (8)
C10	0.0142 (13)	0.0168 (12)	0.0195 (13)	-0.0022 (10)	0.0035 (10)	-0.0008 (10)
N1	0.0171 (11)	0.0176 (10)	0.0173 (10)	-0.0023 (8)	-0.0008 (9)	-0.0002 (9)
O1	0.0259 (10)	0.0380 (12)	0.0165 (9)	-0.0116 (9)	0.0015 (8)	-0.0001 (8)
O2	0.0242 (11)	0.0347 (11)	0.0180 (10)	-0.0156 (9)	0.0012 (8)	0.0041 (9)
Cl1	0.0242 (3)	0.0352 (3)	0.0164 (3)	-0.0106 (3)	-0.0016 (3)	0.0029 (2)

Geometric parameters (\AA , $^\circ$)

C1—N1	1.306 (3)	C6—C7	1.408 (4)
C1—C2	1.424 (4)	C6—H6	0.95
C1—Cl1	1.736 (3)	C7—C8	1.368 (4)
C2—C3	1.380 (4)	C7—H7	0.95
C2—C10	1.498 (3)	C8—C9	1.407 (4)
C3—C4	1.402 (4)	C8—H8	0.95
C3—H3	0.95	C9—N1	1.389 (3)
C4—C9	1.418 (4)	C10—O1	1.208 (3)
C4—C5	1.420 (4)	C10—O2	1.323 (3)
C5—C6	1.368 (4)	O2—H2	0.84
C5—H5	0.95		
N1—C1—C2	124.9 (2)	C5—C6—H6	119.8
N1—C1—Cl1	113.6 (2)	C7—C6—H6	119.8
C2—C1—Cl1	121.5 (2)	C8—C7—C6	121.3 (3)
C3—C2—C1	116.3 (2)	C8—C7—H7	119.3
C3—C2—C10	120.2 (2)	C6—C7—H7	119.3
C1—C2—C10	123.5 (2)	C7—C8—C9	119.5 (3)
C2—C3—C4	121.5 (3)	C7—C8—H8	120.2
C2—C3—H3	119.3	C9—C8—H8	120.2
C4—C3—H3	119.3	N1—C9—C8	119.2 (2)
C3—C4—C9	117.8 (2)	N1—C9—C4	121.0 (2)
C3—C4—C5	123.0 (3)	C8—C9—C4	119.8 (2)
C9—C4—C5	119.2 (2)	O1—C10—O2	123.6 (2)
C6—C5—C4	119.8 (3)	O1—C10—C2	124.7 (2)
C6—C5—H5	120.1	O2—C10—C2	111.7 (2)
C4—C5—H5	120.1	C1—N1—C9	118.5 (2)
C5—C6—C7	120.3 (2)	C10—O2—H2	109.5
N1—C1—C2—C3	-0.8 (4)	C7—C8—C9—C4	-0.3 (4)
Cl1—C1—C2—C3	179.6 (2)	C3—C4—C9—N1	-1.0 (4)
N1—C1—C2—C10	179.0 (2)	C5—C4—C9—N1	179.4 (2)
Cl1—C1—C2—C10	-0.6 (4)	C3—C4—C9—C8	-179.7 (2)
C1—C2—C3—C4	0.7 (4)	C5—C4—C9—C8	0.7 (4)

C10—C2—C3—C4	-179.2 (2)	C3—C2—C10—O1	-178.2 (3)
C2—C3—C4—C9	0.2 (4)	C1—C2—C10—O1	1.9 (4)
C2—C3—C4—C5	179.8 (3)	C3—C2—C10—O2	3.3 (4)
C3—C4—C5—C6	179.7 (3)	C1—C2—C10—O2	-176.6 (2)
C9—C4—C5—C6	-0.6 (4)	C2—C1—N1—C9	0.1 (4)
C4—C5—C6—C7	0.3 (4)	C11—C1—N1—C9	179.68 (18)
C5—C6—C7—C8	0.1 (4)	C8—C9—N1—C1	179.6 (2)
C6—C7—C8—C9	0.0 (4)	C4—C9—N1—C1	0.9 (4)
C7—C8—C9—N1	-179.0 (3)		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H2 \cdots N1 ⁱ	0.84	1.95	2.768 (3)	164.
C3—H3 \cdots O2	0.95	2.34	2.685 (4)	101.
C8—H8 \cdots O1 ⁱⁱ	0.95	2.37	3.290 (4)	163.

Symmetry codes: (i) $x-1, y-1/2, -z+1/2$; (ii) $x+1, y+1/2, -z+1/2$.

Fig. 1




Fig. 2




Fig. 3

