

(2*R*)-2-Methylpiperazinedium tetrachloridocuprate(II)

Li Zhaung Chen* and Sheng Wan

School of Material Science and Engineering, Jiangsu University of Science and Technology, Zhenjiang 212003, People's Republic of China

Correspondence e-mail: clz1977@sina.com


Received 21 January 2010; accepted 2 March 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.009\text{ \AA}$; R factor = 0.025; wR factor = 0.060; data-to-parameter ratio = 23.7.

In the title compound, $(\text{C}_5\text{H}_{14}\text{N}_2)[\text{CuCl}_4]$, the copper(II) ion has a slightly tetrahedrally distorted square-planar coordination geometry and the diprotonated piperazine ring adopts a chair conformation. In the crystal structure, cations and anions are linked by intermolecular $\text{N}-\text{H}\cdots\text{Cl}$ hydrogen bonds, forming a three-dimensional network.

Related literature

For the ferroelectric and non-linear optical properties of chiral organic ligands, see: Fu *et al.* (2007); Qu *et al.* (2003). For transition metal complexes of 2-methylpiperazine, see: Ye *et al.* (2009). For puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data

$(\text{C}_5\text{H}_{14}\text{N}_2)[\text{CuCl}_4]$

$M_r = 307.53$

Orthorhombic, $P2_12_12_1$

$a = 6.0169 (12)\text{ \AA}$

$b = 12.985 (3)\text{ \AA}$

$c = 14.644 (3)\text{ \AA}$

$V = 1144.1 (4)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 2.80\text{ mm}^{-1}$

$T = 293\text{ K}$

$0.30 \times 0.25 \times 0.22\text{ mm}$

Data collection

Rigaku SCXmini diffractometer

Absorption correction: multi-scan
(*CrystalClear*; Rigaku, 2005)

$T_{\min} = 0.87$, $T_{\max} = 0.90$

11992 measured reflections

2627 independent reflections

2469 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.051$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.025$

$wR(F^2) = 0.060$

$S = 1.10$

2627 reflections

111 parameters

1 restraint

H-atom parameters constrained

$\Delta\rho_{\max} = 0.30\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.53\text{ e \AA}^{-3}$

Absolute structure: Flack (1983),
1091 Friedel pairs

Flack parameter: 0.00 (3)

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1A \cdots Cl1 ⁱ	0.90	2.36	3.149 (6)	147
N1—H1B \cdots Cl1 ⁱⁱ	0.90	2.31	3.182 (6)	163
N2—H2A \cdots Cl4 ⁱⁱⁱ	0.90	2.33	3.218 (6)	168
N2—H2B \cdots Cl4 ^{iv}	0.90	2.39	3.192 (6)	148

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{5}{2}, -z$; (ii) $x - \frac{1}{2}, -y + \frac{5}{2}, -z$; (iii) $-x + 2, y - \frac{1}{2}, -z + \frac{1}{2}$;
(iv) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

This work was supported by a start-up grant from Jiangsu University of Science and Technology

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2412).

References

- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Fu, D.-W., Song, Y.-M., Wang, G.-X., Ye, Q., Xiong, R.-G., Akutagawa, T., Nakamura, T., Chan, P. W. H. & Huang, S. D. (2007). *J. Am. Chem. Soc.* **129**, 5346–5347.
- Qu, Z.-R., Zhao, H., Wang, X.-S., Li, Y.-H., Song, Y.-M., Lui, Y.-J., Ye, Q., Xiong, R.-G., Abrahams, B. F., Xue, Z.-L. & You, X.-Z. (2003). *Inorg. Chem.* **42**, 7710–7712.
- Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Ye, H.-Y., Fu, D.-W., Zhang, Y., Zhang, W., Xiong, R.-G. & Huang, S. D. (2009). *J. Am. Chem. Soc.* **131**, 42–43.

supplementary materials

Acta Cryst. (2010). E66, m376 [doi:10.1107/S1600536810007877]

(2*R*)-2-Methylpiperazinium tetrachloridocuprate(II)

L. Z. Chen and S. Wan

Comment

The existence of a chiral centre in an organic ligand is very important for the construction noncentrosymmetric or chiral coordination polymers that exhibit desirable physical properties such as ferroelectricity (Fu *et al.*, 2007) and nonlinear optical second harmonic generation (Qu *et al.*, 2003). Chiral (*R*)-2-methylpiperazine has a chiral centre which have shown tremendous scope in the synthesis of transition metal complexes (Ye *et al.*, 2009). The construction of new members of this family of ligands is an important direction in the development of modern coordination chemistry. We report here the crystal structure of the title compound

The asymmetric unit of the title compound consists of a diprotonated (*R*)-2-methylpiperazine cation and a tetrachlorocuprate anion (Fig. 1). The copper(II) metal centre is in a slightly tetrahedrally distorted square-planar coordination geometry (maximum displacement 0.0252 (18) Å for atom Cl1). The 6-membered piperazine ring adopts a chair conformation, with puckering parameters (Cremer & Pople, 1975) $Q = 0.570$ (6) Å, $\theta = 178.6$ (6)° and $\varphi = -127.3$ (3)°. The crystal structure is stabilized by inter-ion N—H···Cl hydrogen interactions (Table 1) forming a three-dimensional network (Fig. 2).

Experimental

A mixture of (*R*)-2-methylpiperazine (1 mmol, 0.1 g), CuCl₂ (1 mmol, 0.136 g) and 10% aqueous HCl (6 ml) were mixed and dissolved in 30 ml water by heating to 353 K (10 minute) forming a clear solution. The reaction mixture was then cooled slowly to room temperature. Single crystals of the title compound suitable for X-ray analysis were formed after 12 days on slow evaporation of the solvent.

Refinement

All H atoms were placed in calculated positions with C—H = 0.93–0.98 Å, N—H = 0.90 Å, and refined using a riding model, with U_{iso}(H) = 1.2U_{eq}(C, N) or 1.5 U_{eq}(C) for methyl H atoms.

Figures




Fig. 1. The asymmetric unit of the title compound with atom labels. Displacement ellipsoids were drawn at the 30% probability level.

supplementary materials




Fig. 2. Packing diagram viewed approximately along the a axis. Hydrogen bonds are drawn as dashed lines.

(2*R*)-2-Methylpiperazinium tetrachloridocuprate(II)

Crystal data

$(C_5H_{14}N_2)[CuCl_4]$	$F(000) = 620$
$M_r = 307.53$	$D_x = 1.785 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: P 2ac 2ab	Cell parameters from 2469 reflections
$a = 6.0169 (12) \text{ \AA}$	$\theta = 3.1\text{--}27.5^\circ$
$b = 12.985 (3) \text{ \AA}$	$\mu = 2.80 \text{ mm}^{-1}$
$c = 14.644 (3) \text{ \AA}$	$T = 293 \text{ K}$
$V = 1144.1 (4) \text{ \AA}^3$	Block, yellow
$Z = 4$	$0.30 \times 0.25 \times 0.22 \text{ mm}$

Data collection

Rigaku SCXmini diffractometer	2627 independent reflections
Radiation source: fine-focus sealed tube graphite	2469 reflections with $I > 2\sigma(I)$
Detector resolution: 13.6612 pixels mm^{-1}	$R_{\text{int}} = 0.051$
ω scans	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 3.1^\circ$
Absorption correction: multi-scan (CrystalClear; Rigaku, 2005)	$h = -7 \rightarrow 7$
$T_{\text{min}} = 0.87, T_{\text{max}} = 0.90$	$k = -16 \rightarrow 16$
11992 measured reflections	$l = -18 \rightarrow 18$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.025$	$w = 1/[\sigma^2(F_o^2) + (0.0239P)^2 + 0.0175P]$
$wR(F^2) = 0.060$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.10$	$(\Delta/\sigma)_{\text{max}} < 0.001$
2627 reflections	$\Delta\rho_{\text{max}} = 0.30 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.53 \text{ e \AA}^{-3}$

111 parameters	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = k F_c [1 + 0.001 x F_c^2 \lambda^3 / \sin(2\theta)]^{1/4}$
1 restraint	Extinction coefficient: 0.193 (9)
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 1091 Friedel pairs
Secondary atom site location: difference Fourier map	Flack parameter: 0.00 (3)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.82482 (13)	1.34903 (5)	0.01022 (5)	0.0237 (3)
Cl4	0.8203 (3)	1.32659 (11)	0.17172 (9)	0.0251 (4)
Cl3	0.8247 (3)	1.17476 (11)	-0.00811 (9)	0.0256 (4)
Cl2	0.8317 (3)	1.52141 (11)	0.02454 (12)	0.0365 (4)
Cl1	0.8196 (3)	1.36387 (11)	-0.14937 (10)	0.0250 (4)
N1	0.8224 (9)	1.0679 (4)	0.1788 (3)	0.0289 (11)
H1A	0.9338	1.1063	0.1559	0.035*
H1B	0.6932	1.0976	0.1626	0.035*
N2	0.6773 (9)	0.8919 (4)	0.2799 (3)	0.0263 (11)
H2A	0.8089	0.8638	0.2951	0.032*
H2B	0.5692	0.8523	0.3038	0.032*
C5	0.6696 (17)	0.7855 (5)	0.1398 (5)	0.055 (2)
H5C	0.5702	0.7410	0.1725	0.083*
H5B	0.8190	0.7603	0.1458	0.083*
H5A	0.6288	0.7867	0.0764	0.083*
C2	0.6553 (12)	0.8933 (5)	0.1786 (4)	0.0300 (14)
H2C	0.5097	0.9221	0.1630	0.036*
C4	0.8395 (11)	1.0661 (5)	0.2797 (4)	0.0309 (14)
H4B	0.8227	1.1354	0.3035	0.037*
H4A	0.9850	1.0408	0.2975	0.037*
C3	0.6623 (12)	0.9975 (5)	0.3195 (4)	0.0304 (13)
H3B	0.6801	0.9939	0.3853	0.036*
H3A	0.5170	1.0264	0.3067	0.036*
C1	0.8335 (12)	0.9624 (5)	0.1390 (4)	0.0323 (14)
H1D	0.9786	0.9329	0.1512	0.039*
H1C	0.8150	0.9666	0.0733	0.039*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0339 (4)	0.0198 (4)	0.0174 (4)	-0.0001 (3)	-0.0003 (3)	0.0000 (3)
Cl4	0.0236 (7)	0.0309 (8)	0.0209 (7)	-0.0006 (7)	-0.0001 (7)	-0.0035 (6)
Cl3	0.0392 (8)	0.0197 (7)	0.0178 (7)	-0.0004 (7)	0.0001 (7)	0.0001 (5)
Cl2	0.0540 (10)	0.0222 (8)	0.0333 (9)	-0.0006 (8)	-0.0017 (10)	-0.0016 (6)
Cl1	0.0238 (7)	0.0308 (8)	0.0205 (7)	-0.0010 (7)	-0.0001 (7)	0.0047 (6)
N1	0.025 (2)	0.034 (3)	0.028 (3)	-0.004 (3)	0.000 (3)	0.013 (2)
N2	0.026 (2)	0.028 (3)	0.024 (3)	-0.004 (3)	0.002 (3)	0.011 (2)
C5	0.090 (6)	0.035 (4)	0.041 (5)	-0.005 (5)	-0.005 (6)	0.000 (3)
C2	0.033 (3)	0.031 (3)	0.026 (3)	-0.001 (3)	-0.003 (3)	0.007 (3)
C4	0.037 (3)	0.031 (3)	0.025 (3)	-0.004 (3)	-0.003 (3)	0.009 (3)
C3	0.037 (3)	0.032 (3)	0.023 (3)	0.001 (3)	0.005 (3)	0.006 (3)
C1	0.037 (3)	0.039 (4)	0.022 (3)	0.000 (3)	0.005 (3)	0.007 (3)

Geometric parameters (\AA , $^\circ$)

Cu1—Cl2	2.2485 (16)	C5—H5C	0.9600
Cu1—Cl3	2.2788 (16)	C5—H5B	0.9600
Cu1—Cl1	2.3451 (15)	C5—H5A	0.9600
Cu1—Cl4	2.3831 (15)	C2—C1	1.514 (9)
N1—C4	1.482 (8)	C2—H2C	0.9800
N1—C1	1.490 (8)	C4—C3	1.506 (9)
N1—H1A	0.9000	C4—H4B	0.9700
N1—H1B	0.9000	C4—H4A	0.9700
N2—C2	1.489 (8)	C3—H3B	0.9700
N2—C3	1.492 (7)	C3—H3A	0.9700
N2—H2A	0.9000	C1—H1D	0.9700
N2—H2B	0.9000	C1—H1C	0.9700
C5—C2	1.513 (9)		
Cl2—Cu1—Cl3	178.25 (7)	N2—C2—C5	111.0 (5)
Cl2—Cu1—Cl1	90.66 (6)	N2—C2—C1	109.0 (5)
Cl3—Cu1—Cl1	87.95 (5)	C5—C2—C1	111.4 (6)
Cl2—Cu1—Cl4	91.68 (6)	N2—C2—H2C	108.5
Cl3—Cu1—Cl4	89.74 (5)	C5—C2—H2C	108.5
Cl1—Cu1—Cl4	177.28 (6)	C1—C2—H2C	108.5
C4—N1—C1	111.9 (4)	N1—C4—C3	110.3 (6)
C4—N1—H1A	109.2	N1—C4—H4B	109.6
C1—N1—H1A	109.2	C3—C4—H4B	109.6
C4—N1—H1B	109.2	N1—C4—H4A	109.6
C1—N1—H1B	109.2	C3—C4—H4A	109.6
H1A—N1—H1B	107.9	H4B—C4—H4A	108.1
C2—N2—C3	111.8 (4)	N2—C3—C4	110.5 (5)
C2—N2—H2A	109.3	N2—C3—H3B	109.6
C3—N2—H2A	109.3	C4—C3—H3B	109.6
C2—N2—H2B	109.3	N2—C3—H3A	109.6

C3—N2—H2B	109.3	C4—C3—H3A	109.6
H2A—N2—H2B	107.9	H3B—C3—H3A	108.1
C2—C5—H5C	109.5	N1—C1—C2	111.3 (5)
C2—C5—H5B	109.5	N1—C1—H1D	109.4
H5C—C5—H5B	109.5	C2—C1—H1D	109.4
C2—C5—H5A	109.5	N1—C1—H1C	109.4
H5C—C5—H5A	109.5	C2—C1—H1C	109.4
H5B—C5—H5A	109.5	H1D—C1—H1C	108.0
C3—N2—C2—C5	−179.7 (7)	N1—C4—C3—N2	56.1 (7)
C3—N2—C2—C1	57.3 (7)	C4—N1—C1—C2	56.4 (7)
C1—N1—C4—C3	−55.8 (7)	N2—C2—C1—N1	−56.0 (7)
C2—N2—C3—C4	−58.1 (7)	C5—C2—C1—N1	−178.8 (6)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1A···Cl1 ⁱ	0.90	2.36	3.149 (6)	147
N1—H1B···Cl1 ⁱⁱ	0.90	2.31	3.182 (6)	163
N2—H2A···Cl4 ⁱⁱⁱ	0.90	2.33	3.218 (6)	168
N2—H2B···Cl4 ^{iv}	0.90	2.39	3.192 (6)	148

Symmetry codes: (i) $x+1/2, -y+5/2, -z$; (ii) $x-1/2, -y+5/2, -z$; (iii) $-x+2, y-1/2, -z+1/2$; (iv) $-x+1, y-1/2, -z+1/2$.

supplementary materials

Fig. 1




Fig. 2

