

8-Hydroxy-2-methylquinolinium tetrachlorido(quinolin-8-olato- κ^2N,O)stannate(IV) acetonitrile monosolvate

Marzieh Vafae, ^a Gholamhossein Mohammadnezhad, ^a Mostafa M. Amini ^a and Seik Weng Ng ^{b*}

^aDepartment of Chemistry, General Campus, Shahid Beheshti University, Tehran 1983963113, Iran, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 1 March 2010; accepted 2 March 2010

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.004$ Å; R factor = 0.023; wR factor = 0.064; data-to-parameter ratio = 19.3.

In the title solvated salt, $(C_{10}H_{10}NO)[SnCl_4(C_9H_6NO)] \cdot CH_3CN$, the Sn^{IV} atom is chelated by the N,O -bidentate 8-hydroxyquinolinato ligand and four chloride ions, generating a distorted $SnONCl_4$ octahedral coordination geometry for the metal. In the crystal, the cations are linked to the anions and the solvent molecules by $O-H \cdots O$ and $N-H \cdots N$ hydrogen bonds, respectively.

Related literature

For the spectroscopic characterization of the tetrachlorido(quinolinato)stannate(IV) anion in other salts, see: Cunningham *et al.* (1977); Douek *et al.* (1967); Frazer & Goffer (1996); Frazer & Rimmer (1968); Greenwood & Ruddick (1967). For the structures of dichloridobis(quinolin-8-olato)tin and dichloridobis(2-methylquinolin-8-olato)tin, see: Archer *et al.* (1987); Lo & Ng (2009). For a related structure, see: Mohammadnezhad *et al.* (2010).



Experimental

Crystal data

$(C_{10}H_{10}NO)[SnCl_4(C_9H_6NO)] \cdot CH_3CN$

$M_r = 605.88$
Monoclinic, $C2/c$

$a = 29.1672$ (14) Å
 $b = 10.9415$ (5) Å
 $c = 15.4907$ (8) Å
 $\beta = 99.9411$ (6)°
 $V = 4869.4$ (4) Å³

$Z = 8$
Mo $K\alpha$ radiation
 $\mu = 1.51$ mm⁻¹
 $T = 296$ K
 $0.40 \times 0.30 \times 0.20$ mm

Data collection

Bruker SMART APEX diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{min} = 0.583$, $T_{max} = 0.752$

22804 measured reflections
5594 independent reflections
4663 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.027$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.023$
 $wR(F^2) = 0.064$
 $S = 1.02$
5594 reflections
290 parameters
2 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{max} = 0.28$ e Å⁻³
 $\Delta\rho_{min} = -0.54$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Sn1—N1	2.203 (2)	Sn1—Cl2	2.4055 (6)
Sn1—O1	2.077 (1)	Sn1—Cl3	2.4130 (6)
Sn1—Cl1	2.3741 (6)	Sn1—Cl4	2.4176 (6)
O1—Sn1—N1	78.08 (6)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O2—H2o \cdots O1	0.85 (1)	1.91 (2)	2.715 (2)	158 (3)
N2—H2n \cdots N3	0.85 (1)	2.15 (1)	2.972 (3)	162 (2)

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2010).

We thank Shahid Beheshti University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5348).

References

- Archer, S. J., Koch, K. R. & Schmidt, S. (1987). *Inorg. Chim. Acta*, **126**, 209–218.
Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Bruker (2009). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Cunningham, D., Finnegan, J., Donaldson, J. D. & Frazer, M. J. (1977). *J. Chem. Soc. Dalton Trans.* pp. 162–164.
Douek, I., Frazer, M. J., Goffer, Z., Goldstein, M., Rimmer, B. & Willis, H. A. (1967). *Spectrochim. Acta*, **23A**, 373–381.
Frazer, M. J. & Goffer, Z. (1996). *J. Chem. Soc. A*, pp. 544–549.
Frazer, M. J. & Rimmer, B. (1968). *J. Chem. Soc. A*, pp. 69–74.

Greenwood, N. N. & Ruddick, J. N. R. (1967). *J. Chem. Soc. A*, pp. 1679–1683.
Lo, K. M. & Ng, S. W. (2009). *Acta Cryst. E***65**, m719.
Mohammadnezhad, G., Amini, M. M. & Langer, V. (2010). *Acta Cryst. C***66**, m44–m47.

Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A***64**, 112–122.
Westrip, S. P. (2010). *publCIF*. In preparation.

supplementary materials

Acta Cryst. (2010). E66, m381-m382 [doi:10.1107/S160053681000810X]

8-Hydroxy-2-methylquinolinium tetrachlorido(quinolin-8-olato- κ^2N,O)stannate(IV) acetonitrile monosolvate

M. Vafae, G. Mohammadnezhad, M. M. Amini and S. W. Ng

Comment

The 8-hydroxyquinoline anion furnishes a number of compounds with both inorganic tin(IV) and organotin(IV) systems; for example, the crystal structure of dichlorobis(quinolin-8-olato)tin has been known some time back (Archer *et al.*, 1987). The presence of a methyl substituent in the 2-position introduces steric problems (Mohammadnezhad *et al.*, 2010); this may affect the ability of the complexes to crystallize well so that fewer such complexes have been reported. The crystal structure of dichlorobis(2-methylquinolin-8-olato)tin has only recently been reported (Lo & Ng, 2009). On the other hand, mixed chelate complexes are difficult to synthesize as the compounds disproportionate into the symmetrical derivatives.

The reaction of tin(IV) chloride with 8-hydroxyquinoline and 2-methyl-8-hydroxyquinoline in acetonitrile yields instead the salt, 8-hydroxy-2-methylquinolinium tetrachloro(quinolin-8-olato)stannate as the acetonitrile solvate (Scheme I, Fig. 1). Owing to the steric bulk of the methyl group, the 2-methyl-8-hydroxyquinoline component does not engage in binding to the tin atom but merely functions as a proton abstractor. In the salt, the tin atom is chelated by the 8-hydroxyquinolinato unit and it exists in an octahedral coordination geometry. The tin-chlorine bonds *trans* to the chelating atoms are significantly shorter than the other tin-chlorine bonds (Table 1). The cation is linked to the anion by an O–H \cdots O hydrogen bond; the cation is linked to solvent molecule by an N–H \cdots N hydrogen bond (Table 2). The stannate has been spectroscopically characterized in other salts (Cunningham *et al.*, 1977; Douek *et al.*, 1967; Frazer & Goffer, 1996; Frazer & Rimmer, 1968; Greenwood & Ruddick, 1967).

Experimental

Stannic chloride pentahydrate (1 mmol, 0.35 g), 8-hydroxyquinoline (1 mmol, 0.15 g) and 2-methyl-8-hydroxyquinoline (1 mmol, 0.16 g) were placed in a convection tube; the tube was filled with acetonitrile and kept at 333 K. Yellow prisms of (I) were collected after a week.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C–H 0.93 to 0.96 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to 1.2 to 1.5 $U(C)$. The nitrogen- and oxygen-bound ones were located in a difference Fourier map, and were refined with distance restraints of N–H = O–H 0.86 \pm 0.01 Å; their temperature factors were refined.

Figures




Fig. 1. The molecular structure of (I): displacement ellipsoids are drawn at the 50% probability level and H atoms are of arbitrary radius.

8-Hydroxy-2-methylquinolinium tetrachlorido(quinolin-8-olato- κ^2N,O)stannate(IV) acetonitrile monosolvate

Crystal data

(C₁₀H₁₀NO)[SnCl₄(C₉H₆NO)]·C₂H₃N

$M_r = 605.88$

Monoclinic, $C2/c$

Hall symbol: $-C\ 2yc$

$a = 29.1672$ (14) Å

$b = 10.9415$ (5) Å

$c = 15.4907$ (8) Å

$\beta = 99.9411$ (6)°

$V = 4869.4$ (4) Å³

$Z = 8$

$F(000) = 2400$

$D_x = 1.653$ Mg m⁻³

Melting point: 408 K

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9937 reflections

$\theta = 2.3$ – 28.3 °

$\mu = 1.51$ mm⁻¹

$T = 296$ K

Prism, yellow

$0.40 \times 0.30 \times 0.20$ mm

Data collection

Bruker SMART APEX
diffractometer

Radiation source: fine-focus sealed tube
graphite

ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)

$T_{\min} = 0.583$, $T_{\max} = 0.752$

22804 measured reflections

5594 independent reflections

4663 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.027$

$\theta_{\max} = 27.5$ °, $\theta_{\min} = 1.4$ °

$h = -37$ → 37

$k = -14$ → 14

$l = -20$ → 20

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.023$

$wR(F^2) = 0.064$

$S = 1.02$

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0319P)^2 + 3.2971P]$

where $P = (F_o^2 + 2F_c^2)/3$

5594 reflections

$$(\Delta/\sigma)_{\max} = 0.001$$

290 parameters

$$\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$$

2 restraints

$$\Delta\rho_{\min} = -0.54 \text{ e } \text{\AA}^{-3}$$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn1	0.395979 (5)	0.698425 (12)	0.408891 (9)	0.03485 (6)
C11	0.37060 (2)	0.76784 (6)	0.53800 (4)	0.05477 (15)
C12	0.40349 (2)	0.89500 (5)	0.34322 (5)	0.06272 (18)
C13	0.476603 (19)	0.70536 (6)	0.47849 (4)	0.04984 (14)
C14	0.31714 (2)	0.68113 (6)	0.33007 (4)	0.05257 (15)
O1	0.39266 (5)	0.51704 (12)	0.44682 (9)	0.0390 (3)
O2	0.34376 (6)	0.38174 (17)	0.54651 (13)	0.0573 (5)
H2O	0.3527 (11)	0.436 (2)	0.5137 (19)	0.092 (12)*
N1	0.41915 (6)	0.60155 (16)	0.29942 (11)	0.0395 (4)
N2	0.29662 (7)	0.20670 (18)	0.61441 (12)	0.0443 (4)
H2N	0.3234 (5)	0.197 (2)	0.6006 (16)	0.044 (7)*
N3	0.38826 (10)	0.1151 (2)	0.58312 (19)	0.0753 (7)
C1	0.43162 (9)	0.6467 (3)	0.22790 (15)	0.0533 (6)
H1	0.4306	0.7308	0.2191	0.064*
C2	0.44621 (9)	0.5712 (3)	0.16521 (16)	0.0621 (7)
H2	0.4549	0.6050	0.1154	0.075*
C3	0.44759 (9)	0.4489 (3)	0.17715 (16)	0.0614 (7)
H3	0.4571	0.3987	0.1351	0.074*
C4	0.43490 (8)	0.3968 (2)	0.25239 (15)	0.0488 (6)
C5	0.43577 (10)	0.2710 (3)	0.2723 (2)	0.0648 (8)
H5	0.4447	0.2149	0.2332	0.078*
C6	0.42365 (10)	0.2317 (2)	0.3480 (2)	0.0618 (7)
H6	0.4248	0.1484	0.3603	0.074*
C7	0.40938 (8)	0.31228 (19)	0.40858 (17)	0.0453 (5)
H7	0.4015	0.2821	0.4602	0.054*
C8	0.40704 (7)	0.43536 (18)	0.39199 (14)	0.0358 (4)
C9	0.42052 (7)	0.47834 (19)	0.31345 (13)	0.0364 (4)
C10	0.30128 (11)	-0.0040 (3)	0.6638 (2)	0.0714 (8)
H10A	0.3128	-0.0270	0.6117	0.107*
H10B	0.2803	-0.0657	0.6777	0.107*
H10C	0.3269	0.0042	0.7116	0.107*
C11	0.27629 (9)	0.1142 (2)	0.64902 (15)	0.0547 (6)
C12	0.23234 (10)	0.1348 (3)	0.67019 (19)	0.0693 (8)
H12	0.2173	0.0719	0.6946	0.083*
C13	0.21137 (9)	0.2439 (3)	0.65597 (19)	0.0687 (8)
H13	0.1821	0.2549	0.6707	0.082*
C14	0.23274 (8)	0.3424 (3)	0.61921 (15)	0.0530 (6)
C15	0.21283 (10)	0.4587 (3)	0.6017 (2)	0.0702 (8)
H15	0.1831	0.4746	0.6129	0.084*
C16	0.23708 (11)	0.5477 (3)	0.5685 (2)	0.0718 (8)
H16	0.2240	0.6251	0.5588	0.086*

supplementary materials

C17	0.28137 (10)	0.5257 (3)	0.54830 (18)	0.0617 (7)
H17	0.2971	0.5879	0.5247	0.074*
C18	0.30146 (8)	0.4131 (2)	0.56320 (15)	0.0456 (5)
C19	0.27715 (8)	0.3204 (2)	0.59882 (14)	0.0431 (5)
C20	0.42250 (11)	0.0664 (2)	0.58977 (18)	0.0582 (7)
C21	0.46668 (12)	0.0043 (3)	0.5990 (3)	0.0996 (13)
H21A	0.4639	-0.0652	0.5606	0.149*
H21B	0.4758	-0.0224	0.6585	0.149*
H21C	0.4898	0.0592	0.5840	0.149*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn1	0.03691 (9)	0.02715 (8)	0.04003 (8)	0.00082 (5)	0.00535 (6)	0.00091 (5)
Cl1	0.0529 (3)	0.0571 (4)	0.0555 (3)	0.0079 (3)	0.0127 (3)	-0.0166 (3)
Cl2	0.0668 (4)	0.0322 (3)	0.0873 (5)	-0.0024 (3)	0.0084 (3)	0.0155 (3)
Cl3	0.0356 (3)	0.0614 (4)	0.0513 (3)	0.0017 (2)	0.0039 (2)	-0.0024 (3)
Cl4	0.0404 (3)	0.0569 (4)	0.0562 (3)	-0.0032 (2)	-0.0035 (2)	0.0031 (3)
O1	0.0508 (9)	0.0289 (7)	0.0405 (8)	0.0029 (6)	0.0172 (6)	0.0023 (6)
O2	0.0440 (9)	0.0543 (11)	0.0803 (13)	0.0083 (8)	0.0295 (9)	0.0185 (9)
N1	0.0414 (10)	0.0411 (10)	0.0362 (9)	-0.0020 (8)	0.0074 (7)	0.0025 (7)
N2	0.0389 (10)	0.0557 (12)	0.0395 (9)	-0.0072 (9)	0.0099 (8)	-0.0046 (8)
N3	0.0781 (18)	0.0534 (14)	0.099 (2)	0.0056 (13)	0.0267 (15)	0.0007 (13)
C1	0.0574 (15)	0.0613 (16)	0.0421 (12)	-0.0056 (12)	0.0111 (11)	0.0116 (11)
C2	0.0541 (15)	0.099 (2)	0.0353 (12)	-0.0032 (15)	0.0130 (11)	0.0036 (13)
C3	0.0488 (14)	0.095 (2)	0.0413 (13)	0.0014 (14)	0.0098 (11)	-0.0218 (14)
C4	0.0411 (12)	0.0569 (15)	0.0481 (13)	0.0005 (11)	0.0069 (10)	-0.0171 (11)
C5	0.0618 (17)	0.0542 (16)	0.0785 (19)	0.0066 (13)	0.0119 (14)	-0.0315 (14)
C6	0.0592 (16)	0.0318 (12)	0.093 (2)	0.0027 (11)	0.0107 (15)	-0.0111 (13)
C7	0.0413 (12)	0.0335 (11)	0.0608 (14)	0.0005 (9)	0.0079 (10)	0.0024 (10)
C8	0.0314 (10)	0.0326 (10)	0.0429 (11)	0.0017 (8)	0.0049 (8)	-0.0023 (8)
C9	0.0305 (10)	0.0399 (11)	0.0382 (10)	-0.0002 (8)	0.0038 (8)	-0.0053 (8)
C10	0.084 (2)	0.0607 (18)	0.0682 (18)	-0.0182 (16)	0.0103 (15)	0.0090 (14)
C11	0.0583 (15)	0.0628 (16)	0.0432 (12)	-0.0222 (13)	0.0089 (11)	-0.0087 (11)
C12	0.0610 (17)	0.087 (2)	0.0643 (17)	-0.0347 (17)	0.0225 (14)	-0.0148 (16)
C13	0.0393 (14)	0.105 (2)	0.0656 (17)	-0.0219 (16)	0.0196 (12)	-0.0243 (17)
C14	0.0364 (12)	0.0810 (18)	0.0420 (12)	-0.0035 (12)	0.0073 (10)	-0.0158 (12)
C15	0.0425 (14)	0.103 (2)	0.0658 (17)	0.0164 (16)	0.0119 (13)	-0.0178 (17)
C16	0.0644 (18)	0.079 (2)	0.0728 (19)	0.0302 (16)	0.0144 (15)	-0.0023 (16)
C17	0.0585 (16)	0.0626 (17)	0.0656 (17)	0.0112 (13)	0.0152 (13)	0.0048 (13)
C18	0.0381 (11)	0.0555 (14)	0.0443 (12)	0.0036 (10)	0.0102 (9)	-0.0019 (10)
C19	0.0350 (11)	0.0592 (14)	0.0345 (10)	-0.0042 (10)	0.0044 (9)	-0.0085 (9)
C20	0.0721 (19)	0.0400 (13)	0.0665 (17)	-0.0085 (13)	0.0229 (14)	-0.0031 (12)
C21	0.068 (2)	0.070 (2)	0.167 (4)	-0.0003 (17)	0.036 (2)	-0.001 (2)

Geometric parameters (\AA , $^\circ$)

Sn1—N1	2.203 (2)	C6—H6	0.9300
Sn1—O1	2.077 (1)	C7—C8	1.371 (3)

Sn1—C11	2.3741 (6)	C7—H7	0.9300
Sn1—C12	2.4055 (6)	C8—C9	1.422 (3)
Sn1—C13	2.4130 (6)	C10—C11	1.482 (4)
Sn1—C14	2.4176 (6)	C10—H10A	0.9600
O1—C8	1.349 (2)	C10—H10B	0.9600
O2—C18	1.348 (3)	C10—H10C	0.9600
O2—H2O	0.849 (10)	C11—C12	1.395 (4)
N1—C1	1.320 (3)	C12—C13	1.341 (5)
N1—C9	1.365 (3)	C12—H12	0.9300
N2—C11	1.332 (3)	C13—C14	1.413 (4)
N2—C19	1.371 (3)	C13—H13	0.9300
N2—H2N	0.852 (10)	C14—C19	1.406 (3)
N3—C20	1.121 (4)	C14—C15	1.406 (4)
C1—C2	1.396 (4)	C15—C16	1.356 (4)
C1—H1	0.9300	C15—H15	0.9300
C2—C3	1.351 (4)	C16—C17	1.401 (4)
C2—H2	0.9300	C16—H16	0.9300
C3—C4	1.403 (4)	C17—C18	1.367 (3)
C3—H3	0.9300	C17—H17	0.9300
C4—C5	1.410 (4)	C18—C19	1.404 (3)
C4—C9	1.415 (3)	C20—C21	1.442 (4)
C5—C6	1.352 (4)	C21—H21A	0.9600
C5—H5	0.9300	C21—H21B	0.9600
C6—C7	1.402 (4)	C21—H21C	0.9600
O1—Sn1—N1	78.08 (6)	O1—C8—C9	118.80 (18)
O1—Sn1—C11	91.77 (4)	C7—C8—C9	118.2 (2)
N1—Sn1—C11	169.83 (5)	N1—C9—C4	121.3 (2)
O1—Sn1—C12	170.48 (4)	N1—C9—C8	117.27 (18)
N1—Sn1—C12	92.42 (5)	C4—C9—C8	121.4 (2)
C11—Sn1—C12	97.74 (3)	C11—C10—H10A	109.5
O1—Sn1—C13	89.60 (4)	C11—C10—H10B	109.5
N1—Sn1—C13	87.37 (5)	H10A—C10—H10B	109.5
C11—Sn1—C13	92.01 (2)	C11—C10—H10C	109.5
C12—Sn1—C13	90.45 (2)	H10A—C10—H10C	109.5
O1—Sn1—C14	88.93 (4)	H10B—C10—H10C	109.5
N1—Sn1—C14	88.00 (5)	N2—C11—C12	117.6 (3)
C11—Sn1—C14	92.45 (2)	N2—C11—C10	118.9 (2)
C12—Sn1—C14	90.26 (2)	C12—C11—C10	123.5 (3)
C13—Sn1—C14	175.34 (2)	C13—C12—C11	121.1 (3)
C8—O1—Sn1	114.85 (12)	C13—C12—H12	119.4
C18—O2—H2O	109 (2)	C11—C12—H12	119.4
C1—N1—C9	119.9 (2)	C12—C13—C14	121.6 (3)
C1—N1—Sn1	129.17 (17)	C12—C13—H13	119.2
C9—N1—Sn1	110.92 (13)	C14—C13—H13	119.2
C11—N2—C19	124.1 (2)	C19—C14—C15	118.5 (3)
C11—N2—H2N	119.6 (16)	C19—C14—C13	116.6 (3)
C19—N2—H2N	116.3 (16)	C15—C14—C13	124.9 (3)
N1—C1—C2	121.6 (3)	C16—C15—C14	119.9 (3)
N1—C1—H1	119.2	C16—C15—H15	120.1

supplementary materials

C2—C1—H1	119.2	C14—C15—H15	120.1
C3—C2—C1	119.8 (2)	C15—C16—C17	121.5 (3)
C3—C2—H2	120.1	C15—C16—H16	119.2
C1—C2—H2	120.1	C17—C16—H16	119.2
C2—C3—C4	120.7 (2)	C18—C17—C16	120.1 (3)
C2—C3—H3	119.7	C18—C17—H17	119.9
C4—C3—H3	119.7	C16—C17—H17	119.9
C3—C4—C5	125.5 (2)	O2—C18—C17	125.2 (2)
C3—C4—C9	116.8 (2)	O2—C18—C19	115.6 (2)
C5—C4—C9	117.7 (2)	C17—C18—C19	119.1 (2)
C6—C5—C4	120.2 (2)	N2—C19—C18	120.2 (2)
C6—C5—H5	119.9	N2—C19—C14	119.0 (2)
C4—C5—H5	119.9	C18—C19—C14	120.8 (2)
C5—C6—C7	122.2 (2)	N3—C20—C21	179.5 (4)
C5—C6—H6	118.9	C20—C21—H21A	109.5
C7—C6—H6	118.9	C20—C21—H21B	109.5
C8—C7—C6	120.2 (2)	H21A—C21—H21B	109.5
C8—C7—H7	119.9	C20—C21—H21C	109.5
C6—C7—H7	119.9	H21A—C21—H21C	109.5
O1—C8—C7	122.9 (2)	H21B—C21—H21C	109.5
N1—Sn1—O1—C8	2.45 (13)	C3—C4—C9—N1	0.0 (3)
Cl1—Sn1—O1—C8	-176.95 (13)	C5—C4—C9—N1	179.4 (2)
Cl3—Sn1—O1—C8	-84.95 (13)	C3—C4—C9—C8	-179.8 (2)
Cl4—Sn1—O1—C8	90.63 (13)	C5—C4—C9—C8	-0.5 (3)
O1—Sn1—N1—C1	179.6 (2)	O1—C8—C9—N1	1.6 (3)
Cl1—Sn1—N1—C1	-177.0 (2)	C7—C8—C9—N1	-178.21 (19)
Cl2—Sn1—N1—C1	0.1 (2)	O1—C8—C9—C4	-178.54 (18)
Cl3—Sn1—N1—C1	-90.3 (2)	C7—C8—C9—C4	1.6 (3)
Cl4—Sn1—N1—C1	90.2 (2)	C19—N2—C11—C12	0.3 (3)
O1—Sn1—N1—C9	-1.57 (13)	C19—N2—C11—C10	-179.2 (2)
Cl1—Sn1—N1—C9	1.8 (4)	N2—C11—C12—C13	0.1 (4)
Cl2—Sn1—N1—C9	178.93 (13)	C10—C11—C12—C13	179.6 (3)
Cl3—Sn1—N1—C9	88.58 (13)	C11—C12—C13—C14	-0.1 (4)
Cl4—Sn1—N1—C9	-90.90 (13)	C12—C13—C14—C19	-0.3 (4)
C9—N1—C1—C2	0.2 (3)	C12—C13—C14—C15	179.4 (3)
Sn1—N1—C1—C2	178.98 (17)	C19—C14—C15—C16	-2.0 (4)
N1—C1—C2—C3	0.2 (4)	C13—C14—C15—C16	178.2 (3)
C1—C2—C3—C4	-0.6 (4)	C14—C15—C16—C17	2.0 (5)
C2—C3—C4—C5	-178.9 (3)	C15—C16—C17—C18	-1.0 (5)
C2—C3—C4—C9	0.5 (4)	C16—C17—C18—O2	-179.9 (3)
C3—C4—C5—C6	178.6 (3)	C16—C17—C18—C19	0.0 (4)
C9—C4—C5—C6	-0.7 (4)	C11—N2—C19—C18	179.1 (2)
C4—C5—C6—C7	0.8 (4)	C11—N2—C19—C14	-0.8 (3)
C5—C6—C7—C8	0.4 (4)	O2—C18—C19—N2	0.0 (3)
Sn1—O1—C8—C7	176.79 (17)	C17—C18—C19—N2	-179.9 (2)
Sn1—O1—C8—C9	-3.1 (2)	O2—C18—C19—C14	179.8 (2)
C6—C7—C8—O1	178.6 (2)	C17—C18—C19—C14	-0.1 (3)
C6—C7—C8—C9	-1.6 (3)	C15—C14—C19—N2	-179.1 (2)
C1—N1—C9—C4	-0.3 (3)	C13—C14—C19—N2	0.7 (3)

Sn1—N1—C9—C4	-179.30 (16)	C15—C14—C19—C18	1.1 (3)
C1—N1—C9—C8	179.5 (2)	C13—C14—C19—C18	-179.1 (2)
Sn1—N1—C9—C8	0.5 (2)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H2o \cdots O1	0.85 (1)	1.91 (2)	2.715 (2)	158 (3)
N2—H2n \cdots N3	0.85 (1)	2.15 (1)	2.972 (3)	162 (2)

Fig. 1

