

Bis(acetonitrile- κN)(1,10-phenanthroline- $\kappa^2 N,N'$)platinum(II) bis(perchlorate)

Kwang Ha

School of Applied Chemical Engineering, The Research Institute of Catalysis, Chonnam National University, Gwangju 500-757, Republic of Korea
Correspondence e-mail: hakwang@chonnam.ac.kr


Received 22 February 2010; accepted 4 March 2010

Key indicators: single-crystal X-ray study; $T = 200\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.012\text{ \AA}$; disorder in solvent or counterion; R factor = 0.042; wR factor = 0.100; data-to-parameter ratio = 11.6.

The asymmetric unit of the title compound, $[\text{Pt}(\text{CH}_3\text{CN})_2(\text{C}_{12}\text{H}_8\text{N}_2)](\text{ClO}_4)_2$, contains one half of a cationic Pt^{II} complex and pair of half perchlorate anions, one of which is disordered over two sites in a 0.53 (3):0.47 (3) ratio. The complex and anions are disposed about a crystallographic mirror plane parallel to the ac plane passing through the Pt and Cl atoms. In the complex, the Pt^{II} ion lies in a distorted square-planar environment defined by four N atoms of the chelating 1,10-phenanthroline ligand and two distinct acetonitrile molecules. The component ions interact by means of intermolecular C—H···O hydrogen bonds.

Related literature

For the synthesis of $[\text{PtCl}_2(\text{phen})]$ ($\text{phen} = 1,10\text{-phenanthroline}$), see: Hodges & Rund (1975). For the crystal structure of $[\text{Pd}(\text{phen})(\text{CH}_3\text{CN})_2](\text{O}_3\text{SCF}_3)_2$, see: Adrian *et al.* (2008).



Experimental

Crystal data

$[\text{Pt}(\text{C}_2\text{H}_3\text{N})_2(\text{C}_{12}\text{H}_8\text{N}_2)](\text{ClO}_4)_2$

$M_r = 656.30$

Orthorhombic, $Pnma$

$a = 9.1407(5)\text{ \AA}$

$b = 11.7822(7)\text{ \AA}$

$c = 18.3215(11)\text{ \AA}$

$V = 1973.2(2)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 7.44\text{ mm}^{-1}$

$T = 200\text{ K}$

$0.28 \times 0.12 \times 0.04\text{ mm}$

Data collection

Bruker SMART 1000 CCD

diffractometer

Absorption correction: multi-scan
(SADABS; Bruker, 2000)

$T_{\min} = 0.763$, $T_{\max} = 1.000$

11860 measured reflections

2043 independent reflections

1540 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.087$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$

$wR(F^2) = 0.100$

$S = 1.02$

2043 reflections

176 parameters

18 restraints

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 2.23\text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -1.82\text{ e \AA}^{-3}$

Table 1

Selected bond angles (°).

N2 ⁱ —Pt1—N2	87.9 (3)	N1—Pt1—N1 ⁱ	81.9 (3)
Symmetry code: (i) $x, -y + \frac{1}{2}, z$.			

Table 2

Hydrogen-bond geometry (Å, °).

D—H···A	D—H	H···A	D···A	D—H···A
C1—H1···O6 ⁱⁱ	0.95	2.60	3.48 (2)	155
C3—H3···O2 ⁱⁱⁱ	0.95	2.54	3.210 (8)	127
C3—H3···O3 ^{iv}	0.95	2.54	3.486 (11)	178
C7—H7A···O6 ⁱⁱ	0.98	2.39	3.066 (18)	126
C7—H7B···O3 ^v	0.98	2.41	3.143 (11)	131

Symmetry codes: (ii) $x + \frac{1}{2}, -y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 1, y - \frac{1}{2}, -z$; (iv) $-x + 1, -y, -z$; (v) $x + \frac{1}{2}, y, -z + \frac{1}{2}$.

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and PLATON (Spek, 2009); software used to prepare material for publication: SHELXL97.

This work was supported by the Priority Research Centers Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2009-0094056).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2230).

References

- Adrian, R. A., Broker, G. A., Tiekink, E. R. T. & Walmsley, J. A. (2008). *Inorg. Chim. Acta*, **361**, 1261–1266.
- Bruker (2000). SADABS, SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Hodges, K. D. & Rund, J. V. (1975). *Inorg. Chem.* **14**, 525–528.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2010). E66, m389 [doi:10.1107/S1600536810008299]

Bis(acetonitrile- κN)(1,10-phenanthroline- $\kappa^2 N,N'$)platinum(II) bis(perchlorate)

K. Ha

Comment

The asymmetric unit of the title compound, $[\text{Pt}(\text{phen})(\text{CH}_3\text{CN})_2](\text{ClO}_4)_2$ (where phen is 1,10-phenanthroline, $C_{12}\text{H}_8\text{N}_2$), contains one half of a cationic Pt^{II} complex and half a perchlorate anion (Fig. 1). The complex and anions are disposed about a crystallographic mirror plane parallel to the ac plane passing through the Pt and Cl atoms (Fig. 2). In the complex, the Pt^{II} ion lies in a distorted square-planar environment defined by four N atoms of the chelating 1,10-phenanthroline ligand and two distinct acetonitrile molecules. The main contribution to the distortion is the tight $\text{N}1-\text{Pt}1-\text{N}1'$ [symmetry code: (i) $x,-y+1/2,z$] chelate angle [$81.9(3)^\circ$], which results in non-linear trans arrangement [$\angle \text{N}1-\text{Pt}1-\text{N}2' = 177.0(2)^\circ$]. The Pt—N bond lengths are almost equal [$\text{Pt}1-\text{N}(\text{phen})$: 2.001 (6) Å; $\text{Pt}1-\text{N}(\text{CH}_3\text{CN})$: 1.994 (7) Å] (Table 1). The component ions interact by means of intermolecular C—H···O hydrogen bonds (Fig. 2 and Table 2).

Experimental

To a solution of $\text{AgClO}_4 \cdot \text{H}_2\text{O}$ (0.1006 g, 0.446 mmol) in CH_3CN (70 ml) was added $[\text{PtCl}_2(\text{phen})]$ (0.0996 g, 0.223 mmol) and refluxed for 7 h. The mixture was filtered to remove AgCl and then the resulting solution was dried under vacuum. The residue was washed with CH_2Cl_2 and dried at 50 °C, to give a pale gray powder (0.1401 g). Crystals suitable for X-ray analysis were obtained by slow evaporation from a CH_3CN solution.

Refinement

H atoms were positioned geometrically and allowed to ride on their respective parent atoms [C—H = 0.95 (aromatic) or 0.98 Å (CH_3) and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{methyl C})$]. The highest peak ($2.23 \text{ e } \text{\AA}^{-3}$) and the deepest hole ($-1.82 \text{ e } \text{\AA}^{-3}$) in the difference Fourier map are located 0.86 and 0.96 Å, respectively, from the atom Pt1. The O atoms (O4, O5 and O6) of the ClO_4 anion displayed relatively large displacement factors so that the anion appears to be partially disordered. The anion was modelled as disordered over two sites with a major site occupancy factor of 0.53 (3). A total of 18 restraints were used in the refinement using the following SHELXL97 (Sheldrick, 2008) commands: SAME 0.020 $\text{Cl}2' > \text{O}6'$ and DELU 0.010 $\text{Cl}2 > \text{O}6'$. In addition, the displacement parameters of the major and minor component atoms $\text{Cl}2$ $\text{Cl}2'$ were constrained using the EADP command.

supplementary materials

Figures




Fig. 1. The structure of the title compound with displacement ellipsoids drawn at the 50% probability level for non-H atoms. Unlabelled atoms are related to labelled atoms by the symmetry operation and the bonds of the minor components of the disordered ClO_4^- anion are shown with dashed lines.




Fig. 2. View of the unit-cell contents of the title compound. Hydrogen-bond interactions are drawn with dashed lines.

Bis(acetonitrile- κN)(1,10-phenanthroline- $\kappa^2\text{N},\text{N}'$)platinum(II) bis(perchlorate)

Crystal data

$[\text{Pt}(\text{C}_2\text{H}_3\text{N})_2(\text{C}_{12}\text{H}_8\text{N}_2)](\text{ClO}_4)_2$	$F(000) = 1256$
$M_r = 656.30$	$D_x = 2.209 \text{ Mg m}^{-3}$
Orthorhombic, $Pnma$	$\text{Mo K}\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ac 2n	Cell parameters from 3403 reflections
$a = 9.1407 (5) \text{ \AA}$	$\theta = 2.2\text{--}25.0^\circ$
$b = 11.7822 (7) \text{ \AA}$	$\mu = 7.44 \text{ mm}^{-1}$
$c = 18.3215 (11) \text{ \AA}$	$T = 200 \text{ K}$
$V = 1973.2 (2) \text{ \AA}^3$	Rod, colorless
$Z = 4$	$0.28 \times 0.12 \times 0.04 \text{ mm}$

Data collection

Bruker SMART 1000 CCD diffractometer	2043 independent reflections
Radiation source: fine-focus sealed tube graphite	1540 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.087$
Absorption correction: multi-scan (SADABS; Bruker, 2000)	$\theta_{\text{max}} = 26.0^\circ, \theta_{\text{min}} = 2.1^\circ$
$T_{\text{min}} = 0.763, T_{\text{max}} = 1.000$	$h = -11 \rightarrow 7$
11860 measured reflections	$k = -14 \rightarrow 14$
	$l = -22 \rightarrow 21$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.042$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.100$	H-atom parameters constrained
$S = 1.02$	$w = 1/[\sigma^2(F_o^2) + (0.0514P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
2043 reflections	$(\Delta/\sigma)_{\max} < 0.001$
176 parameters	$\Delta\rho_{\max} = 2.23 \text{ e \AA}^{-3}$
18 restraints	$\Delta\rho_{\min} = -1.82 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Pt1	0.96551 (5)	0.2500	0.10603 (2)	0.02158 (17)	
N1	0.8473 (6)	0.1387 (5)	0.0483 (3)	0.0219 (14)	
N2	1.0774 (7)	0.1326 (5)	0.1611 (4)	0.0278 (16)	
C1	0.8490 (8)	0.0253 (6)	0.0516 (4)	0.0264 (18)	
H1	0.9141	-0.0111	0.0845	0.032*	
C2	0.7580 (9)	-0.0405 (7)	0.0081 (5)	0.034 (2)	
H2	0.7628	-0.1209	0.0110	0.041*	
C3	0.6622 (9)	0.0097 (7)	-0.0387 (5)	0.031 (2)	
H3	0.5999	-0.0355	-0.0682	0.038*	
C4	0.6557 (8)	0.1302 (7)	-0.0431 (5)	0.0304 (19)	
C5	0.7512 (7)	0.1902 (6)	0.0018 (4)	0.0229 (17)	
C6	0.5587 (9)	0.1932 (8)	-0.0901 (4)	0.034 (2)	
H6	0.4937	0.1535	-0.1215	0.041*	
C7	1.2397 (10)	0.0000 (8)	0.2375 (5)	0.040 (2)	
H7A	1.2170	-0.0791	0.2254	0.061*	
H7B	1.2217	0.0129	0.2896	0.061*	
H7C	1.3427	0.0154	0.2264	0.061*	
C8	1.1474 (8)	0.0750 (6)	0.1947 (4)	0.0255 (18)	
Cl1	0.4808 (3)	0.2500	0.14861 (16)	0.0273 (6)	
O1	0.3890 (9)	0.2500	0.2122 (5)	0.040 (2)	
O2	0.3939 (10)	0.2500	0.0837 (4)	0.040 (2)	
O3	0.5699 (7)	0.1506 (6)	0.1490 (4)	0.0515 (19)	
Cl2	0.430 (2)	0.7500	0.341 (4)	0.033 (2)	0.53 (3)
O4	0.272 (2)	0.7500	0.340 (2)	0.080 (10)	0.53 (3)

supplementary materials

O5	0.468 (3)	0.7500	0.4176 (11)	0.088 (10)	0.53 (3)
O6	0.4825 (19)	0.6495 (12)	0.3106 (12)	0.077 (8)	0.53 (3)
Cl2'	0.461 (3)	0.7500	0.340 (4)	0.033 (2)	0.47 (3)
O4'	0.312 (3)	0.7500	0.365 (2)	0.068 (10)	0.47 (3)
O5'	0.453 (3)	0.7500	0.2625 (11)	0.111 (15)	0.47 (3)
O6'	0.5367 (19)	0.6564 (15)	0.3651 (15)	0.073 (8)	0.47 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pt1	0.0226 (3)	0.0172 (2)	0.0250 (3)	0.000	-0.00007 (19)	0.000
N1	0.027 (4)	0.014 (3)	0.025 (4)	-0.002 (2)	0.003 (3)	-0.003 (3)
N2	0.032 (4)	0.020 (3)	0.031 (4)	-0.005 (3)	-0.005 (3)	-0.003 (3)
C1	0.026 (4)	0.022 (4)	0.031 (5)	0.004 (3)	0.001 (3)	-0.005 (3)
C2	0.038 (5)	0.020 (4)	0.044 (6)	-0.006 (4)	0.006 (4)	-0.006 (4)
C3	0.031 (5)	0.030 (5)	0.034 (5)	-0.012 (4)	0.003 (4)	-0.010 (4)
C4	0.028 (5)	0.033 (5)	0.030 (5)	0.001 (3)	-0.004 (4)	-0.006 (4)
C5	0.019 (4)	0.027 (4)	0.022 (4)	0.000 (3)	0.003 (3)	0.002 (3)
C6	0.037 (5)	0.057 (6)	0.008 (4)	-0.011 (4)	0.000 (3)	-0.005 (3)
C7	0.047 (6)	0.038 (5)	0.036 (6)	0.012 (4)	-0.012 (4)	0.002 (4)
C8	0.025 (4)	0.020 (4)	0.032 (5)	-0.001 (3)	-0.002 (4)	0.000 (4)
Cl1	0.0272 (14)	0.0256 (14)	0.0292 (17)	0.000	-0.0033 (12)	0.000
O1	0.038 (5)	0.039 (5)	0.042 (6)	0.000	0.004 (4)	0.000
O2	0.055 (6)	0.031 (5)	0.035 (5)	0.000	-0.014 (4)	0.000
O3	0.058 (4)	0.055 (4)	0.041 (4)	0.033 (3)	-0.003 (3)	-0.003 (3)
Cl2	0.030 (7)	0.0226 (15)	0.047 (3)	0.000	-0.003 (11)	0.000
O4	0.031 (9)	0.077 (18)	0.13 (3)	0.000	-0.017 (11)	0.000
O5	0.11 (2)	0.09 (2)	0.063 (10)	0.000	-0.040 (13)	0.000
O6	0.072 (11)	0.034 (8)	0.124 (19)	0.002 (8)	0.025 (12)	-0.031 (10)
Cl2'	0.030 (7)	0.0226 (15)	0.047 (3)	0.000	-0.003 (11)	0.000
O4'	0.023 (11)	0.09 (2)	0.09 (2)	0.000	0.005 (13)	0.000
O5'	0.09 (2)	0.21 (4)	0.034 (9)	0.000	-0.016 (11)	0.000
O6'	0.057 (11)	0.037 (10)	0.12 (2)	0.013 (8)	0.007 (11)	0.028 (12)

Geometric parameters (\AA , $^\circ$)

Pt1—N2 ⁱ	1.994 (7)	C6—H6	0.9500
Pt1—N2	1.994 (7)	C7—C8	1.452 (11)
Pt1—N1	2.001 (6)	C7—H7A	0.9800
Pt1—N1 ⁱ	2.001 (6)	C7—H7B	0.9800
N1—C1	1.337 (9)	C7—H7C	0.9800
N1—C5	1.366 (9)	Cl1—O3 ⁱ	1.427 (6)
N2—C8	1.118 (9)	Cl1—O3	1.427 (6)
C1—C2	1.388 (11)	Cl1—O2	1.430 (9)
C1—H1	0.9500	Cl1—O1	1.436 (9)
C2—C3	1.360 (12)	Cl2—O6 ⁱⁱ	1.40 (3)
C2—H2	0.9500	Cl2—O6	1.40 (3)
C3—C4	1.424 (11)	Cl2—O5	1.44 (7)

C3—H3	0.9500	Cl2—O4	1.44 (2)
C4—C5	1.392 (10)	Cl2'—O6 ⁱⁱ	1.38 (3)
C4—C6	1.442 (11)	Cl2'—O6'	1.38 (3)
C5—C5 ⁱ	1.410 (14)	Cl2'—O5'	1.42 (8)
C6—C6 ⁱ	1.338 (19)	Cl2'—O4'	1.44 (2)
N2 ⁱ —Pt1—N2	87.9 (3)	C4—C6—H6	119.5
N2 ⁱ —Pt1—N1	177.0 (2)	C8—C7—H7A	109.5
N2—Pt1—N1	95.1 (2)	C8—C7—H7B	109.5
N2 ⁱ —Pt1—N1 ⁱ	95.1 (2)	H7A—C7—H7B	109.5
N2—Pt1—N1 ⁱ	177.0 (2)	C8—C7—H7C	109.5
N1—Pt1—N1 ⁱ	81.9 (3)	H7A—C7—H7C	109.5
C1—N1—C5	118.6 (6)	H7B—C7—H7C	109.5
C1—N1—Pt1	128.7 (5)	N2—C8—C7	179.2 (9)
C5—N1—Pt1	112.7 (5)	O3 ⁱ —Cl1—O3	110.4 (6)
C8—N2—Pt1	173.4 (6)	O3 ⁱ —Cl1—O2	108.7 (3)
N1—C1—C2	121.7 (7)	O3—Cl1—O2	108.7 (3)
N1—C1—H1	119.1	O3 ⁱ —Cl1—O1	109.3 (3)
C2—C1—H1	119.1	O3—Cl1—O1	109.3 (3)
C3—C2—C1	120.3 (8)	O2—Cl1—O1	110.5 (6)
C3—C2—H2	119.9	O6 ⁱⁱ —Cl2—O6	116 (4)
C1—C2—H2	119.9	O6 ⁱⁱ —Cl2—O5	108 (3)
C2—C3—C4	119.7 (7)	O6—Cl2—O5	108 (3)
C2—C3—H3	120.1	O6 ⁱⁱ —Cl2—O4	110 (2)
C4—C3—H3	120.1	O6—Cl2—O4	110 (2)
C5—C4—C3	116.5 (7)	O5—Cl2—O4	105 (3)
C5—C4—C6	118.5 (7)	O6 ⁱⁱ —Cl2'—O6'	106 (4)
C3—C4—C6	125.0 (8)	O6 ⁱⁱ —Cl2'—O5'	111 (3)
N1—C5—C4	123.1 (7)	O6'—Cl2'—O5'	111 (3)
N1—C5—C5 ⁱ	116.3 (4)	O6 ⁱⁱ —Cl2'—O4'	111 (3)
C4—C5—C5 ⁱ	120.5 (5)	O6'—Cl2'—O4'	111 (3)
C6 ⁱ —C6—C4	121.0 (5)	O5'—Cl2'—O4'	106 (4)
C6 ⁱ —C6—H6	119.5		
N2—Pt1—N1—C1	1.5 (7)	C1—N1—C5—C4	-0.7 (11)
N1 ⁱ —Pt1—N1—C1	-178.3 (5)	Pt1—N1—C5—C4	-179.0 (6)
N2—Pt1—N1—C5	179.6 (5)	C1—N1—C5—C5 ⁱ	178.5 (5)
N1 ⁱ —Pt1—N1—C5	-0.2 (6)	Pt1—N1—C5—C5 ⁱ	0.2 (5)
C5—N1—C1—C2	1.2 (11)	C3—C4—C5—N1	-0.1 (11)
Pt1—N1—C1—C2	179.2 (6)	C6—C4—C5—N1	179.9 (7)
N1—C1—C2—C3	-1.0 (12)	C3—C4—C5—C5 ⁱ	-179.2 (5)
C1—C2—C3—C4	0.2 (12)	C6—C4—C5—C5 ⁱ	0.8 (9)
C2—C3—C4—C5	0.3 (12)	C5—C4—C6—C6 ⁱ	-0.8 (9)
C2—C3—C4—C6	-179.7 (8)	C3—C4—C6—C6 ⁱ	179.2 (6)

Symmetry codes: (i) $x, -y+1/2, z$; (ii) $x, -y+3/2, z$.


supplementary materials

Hydrogen-bond geometry (\AA , $^\circ$)

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C1—H1 \cdots O6 ⁱⁱⁱ	0.95	2.60	3.48 (2)	155.
C3—H3 \cdots O2 ^{iv}	0.95	2.54	3.210 (8)	127.
C3—H3 \cdots O3 ^v	0.95	2.54	3.486 (11)	178.
C7—H7A \cdots O6 ⁱⁱⁱ	0.98	2.39	3.066 (18)	126.
C7—H7B \cdots O3 ^{vi}	0.98	2.41	3.143 (11)	131.

Symmetry codes: (iii) $x+1/2, -y+1/2, -z+1/2$; (iv) $-x+1, y-1/2, -z$; (v) $-x+1, -y, -z$; (vi) $x+1/2, y, -z+1/2$.

Fig. 1



supplementary materials

Fig. 2

