

Bis[4-(2-carbamoylhydrazin-1-ylidene- κ^2N^1,O)-5-hydroxymethyl-2-methylpyridinium-3-olato- κO^3]cobalt(II) dinitrate dihydrate

Dragoslav Vidovic^{a*} and Violeta Jevtovic^b

^aChemistry Research Laboratory, Chemistry Department, University of Oxford, Mansfield Road, Oxford OX1 3TA, England, and ^bDepartment of Chemistry, Faculty of Sciences, University of Novi Sad, 21000 Novi Sad, Trg D. Obradovica 3, Serbia
Correspondence e-mail: drasko.vidovic@gmail.com


Received 11 January 2010; accepted 28 January 2010

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(C-C) = 0.003$ Å; R factor = 0.033; wR factor = 0.041; data-to-parameter ratio = 13.4.

The asymmetric unit of the title compound, $[Co(C_9H_{12}N_4O_3)_2](NO_3)_2 \cdot 2H_2O$, consists of a discrete cationic $[Co(PLSC)_2]^{2+}$ complex unit [PLSC is 4-(2-carbamoylhydrazin-1-ylidene)-5-hydroxymethyl-2-methylpyridinium-3-olato], two NO_3^- and two water molecules. The two tridentate PLSC ligands of the cation are zwitterions related to each other by a non-crystallographic C_2 axis. The Co^{II} ion is in a disorted octahedral coordination environment. The crystal structure is composed of alternating NO_3^-/H_2O and complex layers supported by extensive $C-H \cdots O$, $N-H \cdots O$ and $N-H \cdots N$ hydrogen bonding.

Related literature

For the preparation and structure of other complexes incorporating PLSC ligands, see for example: Poleti *et al.* (2003); Leovac *et al.* (2007a); Jacimovic *et al.* (2007); Knezevic *et al.* (2003). For the preparation and structures of similar complexes incorporating thiosemicarbazone (TSC) ligands, see: Belicchi Ferrari *et al.* (1998); Leovac *et al.* (2007b). For background to the biological activity of semicarbazones and thiosemicarbazones, see: West *et al.* (1991). For puckering parameters, see: Cremer & Pople (1975). For the Chebychev weighting scheme, see: Prince (1982); Watkin (1994).



Experimental

Crystal data

$[Co(C_9H_{12}N_4O_3)_2](NO_3)_2 \cdot 2H_2O$
 $M_r = 667.41$
 Monoclinic, $P2_1/c$
 $a = 11.0358$ (1) Å
 $b = 18.4859$ (2) Å
 $c = 13.8380$ (1) Å
 $\beta = 106.5705$ (6)°

$V = 2705.80$ (4) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.72$ mm⁻¹
 $T = 150$ K
 $0.38 \times 0.08 \times 0.03$ mm

Data collection

Nonius KappaCCD diffractometer
 Absorption correction: multi-scan
DENZO/SCALEPACK
 (Otwinowski & Minor, 1997)
 $T_{min} = 0.92$, $T_{max} = 0.98$

38405 measured reflections
 5212 independent reflections
 4121 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.056$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.041$
 $S = 1.14$
 5212 reflections

389 parameters
 H-atom parameters constrained
 $\Delta\rho_{max} = 0.56$ e Å⁻³
 $\Delta\rho_{min} = -0.42$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
C7—H71 \cdots O11	0.94	2.33	2.992 (3)	127
C17—H172 \cdots O11 ⁱ	0.97	2.58	3.331 (3)	135
C22—H221 \cdots O2 ⁱⁱ	0.93	2.35	3.153 (3)	144
N27—H271 \cdots O35 ⁱⁱⁱ	0.86	2.18	2.950 (3)	149
N27—H271 \cdots N36 ⁱⁱⁱ	0.86	2.57	3.400 (3)	163
N27—H271 \cdots O37 ⁱⁱⁱ	0.86	2.27	2.995 (3)	142
C30—H301 \cdots O37 ^{iv}	0.97	2.49	3.369 (3)	151
O11—H325 \cdots O16 ⁱ	0.82	2.00	2.783 (3)	159
O11—H325 \cdots O18 ⁱ	0.82	2.59	3.132 (3)	125
O39—H11 \cdots O40	0.84	2.01	2.851 (3)	175
O31—H17 \cdots O42 ^v	0.81	1.95	2.718 (3)	158
C17—H23 \cdots O40	0.96	2.59	3.391 (3)	141
O39—H45 \cdots O31 ^{vi}	0.85	2.05	2.892 (3)	171
O34—H7 \cdots O11 ^{vii}	0.83	1.96	2.767 (3)	164
O34—H19 \cdots O25	0.83	2.06	2.852 (3)	158
N13—H131 \cdots O40	0.75	2.22	2.963 (3)	169
N13—H131 \cdots O42	0.75	2.38	2.971 (3)	136
N5—H2 \cdots O38 ⁱ	0.86	1.93	2.789 (3)	174
N4—H3 \cdots O37 ^{viii}	0.86	2.09	2.948 (3)	174
N33—H4 \cdots O39 ^{ix}	0.86	2.08	2.893 (3)	158
N33—H5 \cdots O34 ⁱⁱ	0.88	2.06	2.874 (3)	153
N4—H6 \cdots O35 ⁱ	0.88	2.09	2.967 (3)	174
N20—H9 \cdots O34 ⁱⁱ	0.86	2.21	2.946 (3)	143

Symmetry codes: (i) $-x + 2, -y + 1, -z + 1$; (ii) $-x + 2, -y + 1, -z$; (iii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (iv) $x, -y + \frac{3}{2}, z - \frac{1}{2}$; (v) $x, y, z - 1$; (vi) $-x + 1, -y + 1, -z$; (vii) $x, -y + \frac{1}{2}, z - \frac{1}{2}$; (viii) $-x + 2, y - \frac{1}{2}, -z + \frac{1}{2}$; (ix) $x + 1, y, z$.

Data collection: *COLLECT* (Nonius, 2001); cell refinement: *DENZO/SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO/SCALEPACK*; program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *CRYSTALS* (Betteridge *et al.*, 2003); molecular graphics: *CAMERON* (Watkin *et al.*, 1996) and *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *CRYSTALS*.

The authors acknowledge the Oxford Chemical Crystallography Service for the use of the instrumentation and the Chemical Crystallography Research Group for helpful discussions.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2980).

References

- Altomare, A., Cascarano, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
- Belicchi Ferrari, M., Gasparri Fava, G., Pelizzi, C., Pelosi, G. & Tarasconi, P. (1998). *Inorg. Chim. Acta*, **269**, 297–301.
- Betteridge, P. W., Carruthers, J. R., Cooper, R. I., Prout, K. & Watkin, D. J. (2003). *J. Appl. Cryst.* **36**, 1487.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Jacimovic, Z., Leovac, V., Giester, G., Tomic, Z. & Szecsenyi, K. (2007). *J. Therm. Anal. Calorim.* **90**, 549–555.
- Knezevic, N., Leovac, M., Jevtovic, V., Grguric-Sipka, S. & Sabo, T. (2003). *Inorg. Chem. Commun.* **6**, 561–563.
- Leovac, V., Jovanovic, Lj., Divjakovic, V., Pevec, A., Leban, I. & Armstrong, T. (2007a). *Polyhedron*, **26**, 49–58.
- Leovac, V., Jovanovic, Lj., Jevtovic, V., Pelosi, G. & Bisceglie, F. (2007b). *Polyhedron*, **26**, 2971–2978.
- Nonius (2001). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Poleti, D., Karanović, L., Leovac, V. M. & Jevtović, V. S. (2003). *Acta Cryst. C* **59**, m73–m75.
- Prince, E. (1982). *Mathematical Techniques in Crystallography and Materials Science*. New York: Springer-Verlag.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Watkin, D. (1994). *Acta Cryst. A* **50**, 411–437.
- Watkin, D. J., Prout, C. K. & Pearce, L. J. (1996). *CAMERON*. Chemical Crystallography Laboratory, Oxford, England.
- West, D. X., Padhye, S. B. & Sonawane, P. B. (1991). *Struct. Bonding (Berlin)*, **76**, 1–50.

supplementary materials

Acta Cryst. (2010). E66, m408-m409 [doi:10.1107/S1600536810003570]

Bis[4-(2-carbamoylhydrazin-1-ylidene- κ^2N^1,O)-5-hydroxymethyl-2-methylpyridinium-3-olato- κO^3]cobalt(II) dinitrate dihydrate

D. Vidovic and V. Jevtovic

Comment

Semicarbazones (SC) and thiosemicarbazones (TSC) are excellent chelating ligands of different denticity that possess a broad range of biological activity as antifungal, anti-viral, anti-malarian and anti-tumour agents (West *et al.*, 1991). The synthetic, structural as well as the biological activity of TSC-based ligands have been explored to a greater extent than their SC-based analogues. In fact, only a handful of reports have been published revealing the syntheses and structures of complexes incorporating 3-hydroxy-5-hydroxymethyl-2-methyl-pyridine-4-carbaldehyde semicarbazone (PLSC) ligand (see for example Knezevic *et al.*, 2003) and all the reports describe complexes incorporating one PLSC ligand except for a report by Leovac *et al.*, (2007b), which includes the synthetic and structural descriptions of a complex with two PTSC ligands in its coordination sphere. It is worth noting that PLSC ligand can adopt three different forms in the coordination sphere of a transition metal namely neutral (but zwitterionic) H_2L , monoanionic HL^- (pyridinium deprotonation) and dianionic L^{2-} (both pyridinium and hydrazine deprotonation) forms (see Fig. 1).

Herein, we report the second PLSC-based complex ($[Co(H_2L)_2] \cdot 2NO_3 \cdot 2H_2O$, **1** (Figure 2), which contains two PLSC ligands in its coordination sphere. However, complex **1** includes both PLSC ligands in their neutral zwitterionic forms while in the corresponding complex reported by Leovac *et al.*, (2007b) both ligands are in their monodeprotonated forms HL^- . Thus, the title complex **1** is the first bis-PLSC-based complex that contains neutral PLSC ligands.

The molecular structure for **1** is shown in Fig. 2 and it contains a discrete dicationic unit $[Co(H_2L)_2]^{2+}$, two NO_3^- anions and two H_2O molecules. The crystal structure is connected by an extensive hydrogen-bonding network. The structure is best described as layered (Fig. 3) in which one layer consists of $[Co(H_2L)_2]^{2+}$ units connected by alternating $O16-H \cdots O11$ and $C22-H \cdots O2$ hydrogen bonds additionally stabilized by a water molecule. The other layer consists of NO_3^- anions and water molecules that also form several hydrogen bonds with the first layer.

The values for certain bond lengths and angles of the ligand backbone (O(phenolic)-C-C-C-N (hydrazine)-N-C-O(carbonyl)) are crystallographic evidence used to determine which form (H_2L , HL^- or L^{2-}) the PLSC ligand adopts once coordinated to a metal centre. For example, the first deprotonation of H_2L , forming monoanionic form HL^- , changes the carbonyl C-O from a double to a single bond and the hydrazine N-N from a single to a double bond. Further deprotonation to form L^{2-} leads to a change in the C-N-C angle of the pyridine ring from 125° to 118° . Thus, the carbonyl C-O (1.249 (3) and 1.295 (3) Å) and N-N (1.368 (2) and 1.375 (2) Å) bond lengths, and the pyridine C-N-C angles (124.9 (2) Å and 124.19 (19) °) are the concrete evidence that the PTSC ligands are in their neutral (H_2L) forms in complex **1**.

The environment around the central cobalt cation in **1** can be best described as a distorted octahedral geometry. In fact, the $N6-Co-N21$ angle (171.12 (7)°) is somewhat similar to the theoretical 180° , but the other two, symmetry-related,

supplementary materials

O2–Co1–O16 (154.44 (7)°), and O18–Co1–O25 (162.82 (6)°) angles greatly deviate from linearity due to the chelation rings strain. Furthermore, the angles formed by the phenolic and carbonyl O atoms and Co (O2–Co1–O18 and O16–Co1–O25) differ from 90° (83.21 (6) and 101.92 (7)°, respectively) confirming a distorted octahedral geometry around the central Co cation.

The ring-puckering parameters defined for the atom sequence O16(O25)–C15(C24)–C8(C23)–C7(C22)–N6(N21)–Co are $\Theta = 64.8 (7)^\circ$, $Q1 = 0.142 (2) \text{ \AA}$ and $Q2 = 0.067 (2) \text{ \AA}$ corresponding to a twist (1T_2) conformation while the six-membered ring has and a total puckering amplitude of $0.1572 (18) \text{ \AA}$ corresponding to a skew ($1S_2$) conformation (Cremer & Pople, 1975). The O16(O25) atoms are the only atoms in the chelate ligand which exhibit strong H bonding interactions with O11 of the NO_3^- anions. The geometry of NO_3^- groups do not deviate from the usual literature values and the summary of the H bonding (O–H \cdots O, N–H \cdots O and C–H \cdots O) in **1** are given in Table 1.

Experimental

The title complex was prepared by the reaction of $\text{Co}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ and PLSC in a 1:1 molar ratio, using warm H_2O as the solvent. Brown single crystals of **1** were obtained after allowing the reaction mixture to stand overnight.

Refinement

The H atoms were all located in a difference map, but those attached to carbon atoms were repositioned geometrically. The H atoms were initially refined with soft restraints on the bond lengths and angles to regularise their geometry (C–H in the range 0.93–0.98, N–H in the range 0.86–0.89 N–H to 0.86 O–H = 0.82 \AA) and $U_{\text{iso}}(\text{H})$ (in the range 1.2–1.5 times U_{eq} of the parent atom), after which the positions were refined with riding constraints.

A three term Chebychev polynomial weighting scheme was applied (Watkin, 1994; Prince, 1982).

Figures




Fig. 1. Different forms of the PLSC ligand.




Fig. 2. The asymmetric unit of **1**. The thermal ellipsoids are drawn at the 50% probability level. H atoms are not shown.




Fig. 3. Crystal packing diagram of **1** viewed along the *z* axis showing the layered motif. H atoms are not shown.

Bis[4-(2-carbamoylhydrazin-1-ylidene- κ^2N^1,O)-5-hydroxymethyl-2-methylpyridinium-3-olato- κO^3]cobalt(II) dinitrate dihydrate

Crystal data

$[\text{Co}(\text{C}_9\text{H}_{12}\text{N}_4\text{O}_3)_2](\text{NO}_3)_2 \cdot 2\text{H}_2\text{O}$

$M_r = 667.41$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 11.0358 (1) \text{ \AA}$

$b = 18.4859 (2) \text{ \AA}$

$c = 13.8380 (1) \text{ \AA}$

$\beta = 106.5705 (6)^\circ$

$V = 2705.80 (4) \text{ \AA}^3$

$Z = 4$

$F(000) = 1380$

$D_x = 1.638 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 6300 reflections

$\theta = 5\text{--}27^\circ$

$\mu = 0.72 \text{ mm}^{-1}$

$T = 150 \text{ K}$

Block, green

$0.38 \times 0.08 \times 0.03 \text{ mm}$

Data collection

Nonius KappaCCD
diffractometer

graphite

ω scans

Absorption correction: multi-scan

DENZO/SCALEPACK (Otwinowski & Minor, 1997)

$T_{\min} = 0.92$, $T_{\max} = 0.98$

38405 measured reflections

6133 independent reflections

4121 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.056$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 5.1^\circ$

$h = -14 \rightarrow 14$

$k = -23 \rightarrow 23$

$l = -17 \rightarrow 17$

Refinement

Refinement on F

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.033$

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

Method, part 1, Chebychev polynomial, (Watkin, 1994, Prince, 1982) $[\text{weight}] = 1.0/[A_0 * T_0(x) + A_1 * T_1(x) \dots + A_{n-1} * T_{n-1}(x)]$

where A_i are the Chebychev coefficients listed below and $x = F / F_{\max}$ Method = Robust Weighting

supplementary materials

(Prince, 1982) $W = [\text{weight}] * [1 - (\Delta F / 6 * \sigma - \text{ma} F)^2]^2$ A_i are: 0.444 0.165 0.244
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.56 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\text{min}} = -0.42 \text{ e } \text{Å}^{-3}$
Extinction correction: Larson (1970), Equation 22
Extinction coefficient: 80 (17)

$wR(F^2) = 0.041$
 $S = 1.14$
5212 reflections
389 parameters
0 restraints
Primary atom site location: structure-invariant direct methods

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.97214 (3)	0.479038 (15)	0.21508 (2)	0.0178
O2	1.10949 (15)	0.41005 (8)	0.18293 (11)	0.0221
C3	1.1436 (2)	0.35913 (12)	0.24402 (16)	0.0197
N4	1.22358 (19)	0.30776 (11)	0.23500 (15)	0.0266
N5	1.09857 (18)	0.35397 (10)	0.32629 (14)	0.0237
N6	1.01658 (17)	0.40702 (10)	0.33633 (13)	0.0189
C7	0.9609 (2)	0.39890 (12)	0.40564 (16)	0.0191
C8	0.8744 (2)	0.45254 (11)	0.42558 (16)	0.0183
C9	0.8127 (2)	0.43675 (12)	0.50011 (16)	0.0200
C10	0.8221 (2)	0.36423 (12)	0.55277 (17)	0.0232
O11	0.94816 (15)	0.34273 (8)	0.60528 (11)	0.0235
C12	0.7357 (2)	0.48771 (12)	0.52394 (17)	0.0229
N13	0.71667 (17)	0.55170 (10)	0.47433 (14)	0.0226
C14	0.7688 (2)	0.56970 (12)	0.40180 (16)	0.0201
C15	0.85115 (19)	0.52005 (12)	0.37386 (15)	0.0186
O16	0.90135 (15)	0.54119 (8)	0.30417 (12)	0.0230
C17	0.7403 (2)	0.64164 (12)	0.35176 (18)	0.0256
O18	1.11612 (15)	0.55823 (9)	0.23239 (11)	0.0246
C19	1.1000 (2)	0.59940 (12)	0.15863 (16)	0.0195
N20	0.99748 (17)	0.59111 (10)	0.07567 (14)	0.0211
N21	0.91416 (17)	0.53710 (9)	0.08134 (13)	0.0187
C22	0.8162 (2)	0.52824 (12)	0.00484 (16)	0.0191
C23	0.7192 (2)	0.47389 (12)	0.00103 (16)	0.0197
C24	0.7303 (2)	0.42092 (12)	0.07771 (17)	0.0212
O25	0.82772 (15)	0.41181 (8)	0.15522 (12)	0.0231
C26	0.6286 (2)	0.37144 (13)	0.06785 (18)	0.0271
N27	0.52905 (19)	0.37585 (12)	-0.01369 (16)	0.0308
C28	0.5171 (2)	0.42374 (14)	-0.08929 (18)	0.0288
C29	0.6111 (2)	0.47329 (13)	-0.08471 (16)	0.0225
C30	0.5993 (2)	0.52333 (13)	-0.17260 (16)	0.0253
O31	0.49006 (17)	0.50463 (10)	-0.25181 (14)	0.0361
C32	0.6314 (3)	0.31578 (15)	0.1459 (2)	0.0380
N33	1.17911 (19)	0.65324 (11)	0.15436 (15)	0.0266
O34	0.92284 (17)	0.30192 (9)	0.05176 (13)	0.0298
O35	0.71622 (18)	0.80490 (10)	0.60566 (13)	0.0335

N36	0.75110 (18)	0.79336 (10)	0.52883 (14)	0.0225
O37	0.68648 (16)	0.81769 (9)	0.44575 (12)	0.0291
O38	0.85047 (15)	0.75880 (9)	0.53404 (13)	0.0298
O39	0.41254 (18)	0.62823 (12)	0.31113 (15)	0.0467
O40	0.5576 (2)	0.67311 (12)	0.50711 (15)	0.0484
N41	0.51580 (19)	0.64733 (13)	0.57582 (15)	0.0311
O42	0.5441 (2)	0.58280 (11)	0.59929 (16)	0.0453
O43	0.4521 (2)	0.68264 (14)	0.61762 (18)	0.0566
H71	0.9774	0.3573	0.4460	0.0228*
H101	0.7720	0.3664	0.6016	0.0279*
H102	0.7876	0.3265	0.5016	0.0267*
H121	0.6942	0.4791	0.5765	0.0442*
H172	0.8136	0.6724	0.3747	0.0384*
H173	0.7197	0.6360	0.2794	0.0387*
H221	0.8066	0.5589	-0.0502	0.0232*
H271	0.4677	0.3462	-0.0184	0.0362*
H301	0.5918	0.5725	-0.1507	0.0287*
H302	0.6725	0.5194	-0.1970	0.0290*
H321	0.5552	0.2872	0.1270	0.0565*
H322	0.6385	0.3400	0.2085	0.0563*
H323	0.7039	0.2852	0.1537	0.0561*
H325	0.9802	0.3770	0.6409	0.0357*
H11	0.4517	0.6415	0.3701	0.0682*
H17	0.4888	0.5336	-0.2964	0.0494*
H23	0.6697	0.6643	0.3664	0.0389*
H45	0.4494	0.5912	0.2972	0.0683*
H7	0.9433	0.2609	0.0759	0.0456*
H19	0.9005	0.3256	0.0950	0.0455*
H324	0.4444	0.4228	-0.1444	0.0326*
H131	0.6742	0.5791	0.4884	0.0126*
H2	1.1172	0.3178	0.3671	0.0294*
H3	1.2527	0.3079	0.1833	0.0331*
H4	1.2454	0.6582	0.2043	0.0307*
H5	1.1634	0.6795	0.0993	0.0304*
H6	1.2462	0.2740	0.2810	0.0331*
H9	0.9850	0.6186	0.0235	0.0258*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.02145 (15)	0.01664 (15)	0.01580 (14)	0.00014 (12)	0.00608 (10)	0.00114 (12)
O2	0.0268 (8)	0.0207 (8)	0.0212 (8)	0.0020 (6)	0.0105 (6)	0.0033 (6)
C3	0.0217 (11)	0.0196 (10)	0.0184 (10)	-0.0021 (8)	0.0069 (8)	-0.0006 (8)
N4	0.0329 (11)	0.0264 (10)	0.0243 (10)	0.0087 (9)	0.0145 (8)	0.0036 (8)
N5	0.0311 (10)	0.0208 (9)	0.0223 (9)	0.0101 (8)	0.0127 (8)	0.0064 (8)
N6	0.0211 (9)	0.0188 (9)	0.0178 (9)	0.0039 (7)	0.0071 (7)	0.0009 (7)
C7	0.0238 (11)	0.0176 (10)	0.0164 (10)	0.0019 (8)	0.0062 (8)	0.0022 (8)
C8	0.0186 (10)	0.0196 (10)	0.0167 (10)	-0.0011 (8)	0.0050 (8)	-0.0021 (8)

supplementary materials

C9	0.0193 (10)	0.0224 (11)	0.0185 (10)	-0.0027 (9)	0.0057 (8)	-0.0014 (9)
C10	0.0253 (11)	0.0230 (11)	0.0237 (11)	-0.0030 (9)	0.0107 (9)	0.0013 (9)
O11	0.0276 (8)	0.0198 (7)	0.0226 (8)	-0.0004 (6)	0.0064 (6)	-0.0011 (6)
C12	0.0226 (11)	0.0268 (11)	0.0211 (11)	-0.0017 (9)	0.0091 (9)	0.0013 (9)
N13	0.0198 (9)	0.0240 (9)	0.0254 (10)	0.0046 (8)	0.0089 (8)	-0.0033 (8)
C14	0.0191 (10)	0.0209 (10)	0.0191 (10)	-0.0022 (8)	0.0036 (8)	-0.0036 (8)
C15	0.0195 (10)	0.0194 (10)	0.0167 (10)	-0.0013 (9)	0.0052 (8)	-0.0021 (9)
O16	0.0298 (8)	0.0180 (7)	0.0241 (8)	0.0012 (6)	0.0126 (7)	0.0028 (6)
C17	0.0278 (12)	0.0213 (11)	0.0279 (12)	0.0040 (9)	0.0083 (9)	0.0009 (9)
O18	0.0267 (8)	0.0256 (8)	0.0206 (8)	-0.0039 (7)	0.0052 (6)	0.0002 (7)
C19	0.0205 (10)	0.0186 (10)	0.0202 (10)	-0.0003 (8)	0.0069 (8)	-0.0015 (8)
N20	0.0225 (9)	0.0188 (9)	0.0213 (9)	-0.0057 (7)	0.0052 (7)	0.0038 (7)
N21	0.0206 (9)	0.0167 (9)	0.0203 (9)	-0.0014 (7)	0.0080 (7)	-0.0002 (7)
C22	0.0209 (10)	0.0203 (10)	0.0173 (10)	0.0004 (9)	0.0073 (8)	0.0014 (8)
C23	0.0200 (10)	0.0202 (10)	0.0195 (10)	0.0006 (9)	0.0069 (8)	-0.0020 (9)
C24	0.0215 (11)	0.0189 (10)	0.0240 (11)	-0.0003 (9)	0.0078 (9)	-0.0021 (9)
O25	0.0245 (8)	0.0198 (8)	0.0236 (8)	-0.0008 (6)	0.0045 (6)	0.0030 (6)
C26	0.0262 (12)	0.0242 (11)	0.0318 (12)	-0.0044 (10)	0.0099 (10)	0.0003 (10)
N27	0.0240 (10)	0.0326 (11)	0.0350 (11)	-0.0106 (9)	0.0074 (9)	0.0000 (9)
C28	0.0256 (12)	0.0330 (13)	0.0260 (12)	-0.0037 (10)	0.0044 (10)	-0.0007 (10)
C29	0.0209 (10)	0.0261 (11)	0.0213 (10)	-0.0020 (9)	0.0074 (9)	-0.0039 (9)
C30	0.0243 (11)	0.0282 (12)	0.0196 (10)	-0.0045 (10)	0.0003 (9)	-0.0004 (9)
O31	0.0329 (9)	0.0399 (10)	0.0267 (8)	-0.0085 (8)	-0.0057 (7)	0.0034 (8)
C32	0.0332 (14)	0.0317 (13)	0.0471 (16)	-0.0090 (11)	0.0083 (12)	0.0121 (12)
N33	0.0265 (10)	0.0259 (10)	0.0250 (10)	-0.0084 (8)	0.0035 (8)	0.0008 (8)
O34	0.0397 (10)	0.0207 (8)	0.0301 (9)	0.0015 (7)	0.0116 (8)	0.0012 (7)
O35	0.0469 (11)	0.0349 (10)	0.0232 (8)	0.0128 (8)	0.0175 (8)	0.0032 (7)
N36	0.0275 (10)	0.0190 (9)	0.0227 (10)	0.0016 (8)	0.0101 (8)	0.0030 (7)
O37	0.0326 (9)	0.0322 (9)	0.0234 (8)	0.0104 (7)	0.0094 (7)	0.0086 (7)
O38	0.0265 (9)	0.0322 (9)	0.0323 (9)	0.0096 (7)	0.0111 (7)	0.0091 (7)
O39	0.0335 (10)	0.0616 (14)	0.0390 (11)	0.0035 (9)	0.0009 (8)	-0.0149 (10)
O40	0.0599 (13)	0.0568 (13)	0.0304 (10)	-0.0153 (11)	0.0161 (9)	0.0023 (9)
N41	0.0272 (11)	0.0426 (13)	0.0229 (10)	0.0009 (9)	0.0060 (8)	-0.0030 (9)
O42	0.0497 (12)	0.0437 (12)	0.0462 (12)	0.0081 (10)	0.0199 (10)	0.0034 (9)
O43	0.0491 (13)	0.0671 (15)	0.0602 (14)	0.0178 (11)	0.0264 (11)	-0.0122 (12)

Geometric parameters (Å, °)

Co1—O2	2.1228 (15)	C19—N33	1.336 (3)
Co1—N6	2.0878 (18)	N20—N21	1.375 (2)
Co1—O16	2.0000 (15)	N20—H9	0.861
Co1—O18	2.1232 (16)	N21—C22	1.290 (3)
Co1—N21	2.0763 (18)	C22—C23	1.458 (3)
Co1—O25	2.0044 (16)	C22—H221	0.931
O2—C3	1.249 (3)	C23—C24	1.423 (3)
C3—N4	1.327 (3)	C23—C29	1.422 (3)
C3—N5	1.369 (3)	C24—O25	1.295 (3)
N4—H3	0.864	C24—C26	1.424 (3)
N4—H6	0.875	C26—N27	1.334 (3)

N5—N6	1.368 (2)	C26—C32	1.485 (3)
N5—H2	0.861	N27—C28	1.348 (3)
N6—C7	1.287 (3)	N27—H271	0.859
C7—C8	1.457 (3)	C28—C29	1.372 (3)
C7—H71	0.937	C28—H324	0.935
C8—C9	1.419 (3)	C29—C30	1.504 (3)
C8—C15	1.425 (3)	C30—O31	1.421 (3)
C9—C10	1.515 (3)	C30—H301	0.970
C9—C12	1.370 (3)	C30—H302	0.964
C10—O11	1.430 (3)	O31—H17	0.814
C10—H101	0.988	C32—H321	0.964
C10—H102	0.990	C32—H322	0.959
O11—H325	0.818	C32—H323	0.961
C12—N13	1.354 (3)	N33—H4	0.856
C12—H121	0.977	N33—H5	0.878
N13—C14	1.334 (3)	O34—H7	0.834
N13—H131	0.753	O34—H19	0.833
C14—C15	1.421 (3)	O35—N36	1.248 (2)
C14—C17	1.491 (3)	N36—O37	1.251 (2)
C15—O16	1.302 (3)	N36—O38	1.253 (2)
C17—H172	0.965	O39—H11	0.844
C17—H173	0.967	O39—H45	0.847
C17—H23	0.955	O40—N41	1.262 (3)
O18—C19	1.245 (3)	N41—O42	1.252 (3)
C19—N20	1.371 (3)	N41—O43	1.219 (3)
O2—Co1—N6	76.54 (6)	C14—C17—H23	111.7
O2—Co1—O16	154.44 (7)	H172—C17—H23	109.2
N6—Co1—O16	85.09 (7)	H173—C17—H23	107.6
O2—Co1—O18	83.21 (6)	Co1—O18—C19	113.69 (14)
N6—Co1—O18	110.63 (7)	O18—C19—N20	120.60 (19)
O16—Co1—O18	86.89 (6)	O18—C19—N33	123.3 (2)
O2—Co1—N21	100.41 (6)	N20—C19—N33	116.08 (19)
N6—Co1—N21	171.12 (7)	C19—N20—N21	116.04 (17)
O16—Co1—N21	100.21 (7)	C19—N20—H9	122.1
O18—Co1—N21	76.98 (6)	N21—N20—H9	121.8
O2—Co1—O25	94.13 (6)	Co1—N21—N20	112.65 (13)
N6—Co1—O25	85.04 (7)	Co1—N21—C22	129.72 (15)
O16—Co1—O25	101.92 (7)	N20—N21—C22	117.59 (18)
O18—Co1—O25	162.82 (6)	N21—C22—C23	123.5 (2)
N21—Co1—O25	86.88 (7)	N21—C22—H221	117.8
Co1—O2—C3	113.95 (13)	C23—C22—H221	118.6
O2—C3—N4	123.8 (2)	C22—C23—C24	122.8 (2)
O2—C3—N5	120.31 (19)	C22—C23—C29	117.9 (2)
N4—C3—N5	115.94 (19)	C24—C23—C29	119.3 (2)
C3—N4—H3	119.3	C23—C24—O25	125.6 (2)
C3—N4—H6	120.1	C23—C24—C26	118.2 (2)
H3—N4—H6	120.6	O25—C24—C26	116.2 (2)
C3—N5—N6	116.22 (18)	Co1—O25—C24	129.50 (14)
C3—N5—H2	121.6	C24—C26—N27	118.6 (2)

supplementary materials

N6—N5—H2	122.1	C24—C26—C32	121.5 (2)
Co1—N6—N5	112.70 (13)	N27—C26—C32	119.9 (2)
Co1—N6—C7	128.37 (15)	C26—N27—C28	124.9 (2)
N5—N6—C7	117.54 (18)	C26—N27—H271	117.8
N6—C7—C8	122.68 (19)	C28—N27—H271	117.4
N6—C7—H71	119.1	N27—C28—C29	119.8 (2)
C8—C7—H71	118.2	N27—C28—H324	119.4
C7—C8—C9	118.30 (19)	C29—C28—H324	120.8
C7—C8—C15	122.33 (19)	C23—C29—C28	119.2 (2)
C9—C8—C15	119.37 (19)	C23—C29—C30	121.9 (2)
C8—C9—C10	123.38 (19)	C28—C29—C30	118.9 (2)
C8—C9—C12	119.5 (2)	C29—C30—O31	109.48 (19)
C10—C9—C12	117.05 (19)	C29—C30—H301	108.6
C9—C10—O11	114.45 (18)	O31—C30—H301	109.9
C9—C10—H101	108.5	C29—C30—H302	110.4
O11—C10—H101	108.5	O31—C30—H302	108.9
C9—C10—H102	108.7	H301—C30—H302	109.6
O11—C10—H102	106.7	C30—O31—H17	104.6
H101—C10—H102	110.0	C26—C32—H321	110.3
C10—O11—H325	106.4	C26—C32—H322	108.3
C9—C12—N13	119.7 (2)	H321—C32—H322	109.9
C9—C12—H121	121.5	C26—C32—H323	109.5
N13—C12—H121	118.8	H321—C32—H323	110.0
C12—N13—C14	124.19 (19)	H322—C32—H323	108.8
C12—N13—H131	118.4	C19—N33—H4	117.6
C14—N13—H131	117.4	C19—N33—H5	118.7
N13—C14—C15	119.4 (2)	H4—N33—H5	123.5
N13—C14—C17	119.5 (2)	H7—O34—H19	107.0
C15—C14—C17	121.12 (19)	O35—N36—O37	119.20 (19)
C8—C15—C14	117.81 (19)	O35—N36—O38	121.04 (19)
C8—C15—O16	125.55 (19)	O37—N36—O38	119.76 (18)
C14—C15—O16	116.61 (19)	H11—O39—H45	108.0
Co1—O16—C15	127.43 (14)	O40—N41—O42	115.9 (2)
C14—C17—H172	109.0	O40—N41—O43	122.6 (2)
C14—C17—H173	109.9	O42—N41—O43	121.4 (2)
H172—C17—H173	109.3		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C7—H71 \cdots O11	0.94	2.33	2.992 (3)	127
C17—H172 \cdots O11 ⁱ	0.97	2.58	3.331 (3)	135
C22—H221 \cdots O2 ⁱⁱ	0.93	2.35	3.153 (3)	144
N27—H271 \cdots O35 ⁱⁱⁱ	0.86	2.18	2.950 (3)	149
N27—H271 \cdots N36 ⁱⁱⁱ	0.86	2.57	3.400 (3)	163
N27—H271 \cdots O37 ⁱⁱⁱ	0.86	2.27	2.995 (3)	142
C30—H301 \cdots O37 ^{iv}	0.97	2.49	3.369 (3)	151
O11—H325 \cdots O16 ⁱ	0.82	2.00	2.783 (3)	159

O11—H325…O18 ⁱ	0.82	2.59	3.132 (3)	125
O39—H11…O40	0.84	2.01	2.851 (3)	175
O31—H17…O42 ^v	0.81	1.95	2.718 (3)	158
C17—H23…O40	0.96	2.59	3.391 (3)	141
O39—H45…O31 ^{vi}	0.85	2.05	2.892 (3)	171
O34—H7…O11 ^{vii}	0.83	1.96	2.767 (3)	164
O34—H19…C24	0.83	2.54	3.149 (3)	131
O34—H19…O25	0.83	2.06	2.852 (3)	158
N13—H131…O40	0.75	2.22	2.963 (3)	169
N13—H131…O42	0.75	2.38	2.971 (3)	136
N5—H2…O38 ⁱ	0.86	1.93	2.789 (3)	174
N4—H3…O37 ^{viii}	0.86	2.09	2.948 (3)	174
N33—H4…O39 ^{ix}	0.86	2.08	2.893 (3)	158
N33—H5…O34 ⁱⁱ	0.88	2.06	2.874 (3)	153
N4—H6…O35 ⁱ	0.88	2.09	2.967 (3)	174
N20—H9…O34 ⁱⁱ	0.86	2.21	2.946 (3)	143

Symmetry codes: (i) $-x+2, -y+1, -z+1$; (ii) $-x+2, -y+1, -z$; (iii) $-x+1, y-1/2, -z+1/2$; (iv) $x, -y+3/2, z-1/2$; (v) $x, y, z-1$; (vi) $-x+1, -y+1, -z$; (vii) $x, -y+1/2, z-1/2$; (viii) $-x+2, y-1/2, -z+1/2$; (ix) $x+1, y, z$.

Fig. 1




Fig. 2




Fig. 3

