

[4'-(3-Pyridyl)-2,2';6',2''-terpyridine]-dithiocyanatocopper(II)

Dongcheng Hu,* Hua Feng* and Changqiu Hu

College of Chemistry & Chemical Engineering, Northwest Normal University, Lanzhou, Gansu 730070, People's Republic of China
Correspondence e-mail: hudch@163.com, hfeng@nwnu.edu.cn


Received 14 March 2010; accepted 18 March 2010

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.059; wR factor = 0.160; data-to-parameter ratio = 17.1.

In the title compound, $[\text{Cu}(\text{NCS})_2(\text{C}_{20}\text{H}_{14}\text{N}_4)]$, the Cu atom is five-coordinated in a tetragonal-pyramidal geometry.

Related literature

For details of the synthesis, see: Constable & Thompson (1992). For related structures, see: Feng *et al.* (2006); Hou *et al.* (2004).



Experimental

Crystal data

$[\text{Cu}(\text{NCS})_2(\text{C}_{20}\text{H}_{14}\text{N}_4)]$
 $M_r = 490.05$
Monoclinic, $P2_1/n$
 $a = 8.2171 (6)\text{ \AA}$
 $b = 23.012 (2)\text{ \AA}$
 $c = 11.2279 (8)\text{ \AA}$
 $\beta = 99.079 (1)^\circ$

$V = 2096.5 (3)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 1.26\text{ mm}^{-1}$
 $T = 298\text{ K}$
 $0.27 \times 0.27 \times 0.18\text{ mm}$

Data collection

Bruker SMART APEX area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2002)
 $T_{\min} = 0.429$, $T_{\max} = 0.805$

12626 measured reflections
4781 independent reflections
3371 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.038$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.059$
 $wR(F^2) = 0.160$
 $S = 1.07$
4781 reflections

280 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.64\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.44\text{ e \AA}^{-3}$

Data collection: *SMART* (Bruker, 2002); cell refinement: *SAINT* (Bruker, 2002); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

The authors thank Northwest Normal University for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5216).

References

- Bruker (2002). *SADABS*, *SAINT* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
Constable, E. C. & Thompson, A. M. W. C. (1992). *J. Chem. Soc. Dalton Trans.*, pp. 2947–2950.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Feng, H., Zhou, X. P., Wu, T., Li, D., Yin, Y. G. & Ng, S. W. (2006). *Inorg. Chim. Acta*, **359**, 4027–4035.
Hou, L., Li, D., Wu, T., Yin, Y.-G. & Ng, S. W. (2004). *Acta Cryst. E* **60**, m1181–m1182.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2010). E66, m442 [doi:10.1107/S1600536810010147]

[4'-(3-Pyridyl)-2,2';6',2"-terpyridine]dithiocyanatocopper(II)

D. Hu, H. Feng and C. Hu

Comment

The heterocyclic ligand 4'-(3-pyridyl)-2,2';6'2"-terpyridine was synthesized according to literature methods (Constable *et al.*, 1992). As a excellent tridentate chelating ligand, it can furnish complexes from a large range of metal salts. The structure of the title compound, (I), obtained from copper(I) thiocyanate is shown in Fig. 1. The Cu(II) atom is chelated by a tridentate terpyridine in a *cis-cis* configuration and two independent thiocyanate in a five-coordinate environment and shows distorted tetragonal pyramid geometry (Fig. 1). The Cu—N bonds range from 1.931 (3) Å to 2.143 (4) Å are comparable with those reported (Hou *et al.*, 2004). In the monomer, one thiocyanate was in the apical site while the terpyridine and the other one thiocyanate were at the equatorial positions.

Experimental

The mixture of CuSCN (0.0125 g, 0.1 mmol), 4'-(3-pyridyl)-2,2';6'2"-terpyridine (0.0155 g, 0.05 mmol), acetonitril (6 ml) were placed and sealed in a 10 ml Teflon-lined stainless steel reactor and heated to 140°C for 72 h, then cooled down to room temperature at a rate of 2°C per 20 min. Green block single crystals suitable for X-ray diffraction were obtained in ca. 60% yield

Refinement

Hydrogen atoms were placed in calculated positions (C—H 0.93 Å; U(H) = 1.2U_{eq}C) and were included in the refinement in the riding model approximation.

Figures




Fig. 1. ORTEP-3 plot of the title compound with the atom labeling scheme. Displacement ellipsoids are drawn at the 35% probability level. H atoms are omitted for clarity.

[4'-(3-Pyridyl)-2,2';6',2"-terpyridine]dithiocyanatocopper(II)

Crystal data

[Cu(NCS)₂(C₂₀H₁₄N₄)]

F(000) = 996

M_r = 490.05

D_x = 1.553 Mg m⁻³

supplementary materials

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 8.2171 (6)$ Å

$b = 23.012 (2)$ Å

$c = 11.2279 (8)$ Å

$\beta = 99.079 (1)^\circ$

$V = 2096.5 (3)$ Å³

$Z = 4$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2079 reflections

$\theta = 2.6\text{--}21.4^\circ$

$\mu = 1.26$ mm⁻¹

$T = 298$ K

Block, green

$0.27 \times 0.27 \times 0.18$ mm

Data collection

Bruker SMART APEX area-detector diffractometer

4781 independent reflections

Radiation source: fine-focus sealed tube graphite

3371 reflections with $I > 2\sigma(I)$

φ and ω scan

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 1.8^\circ$

Absorption correction: multi-scan (SADABS; Bruker, 2002)

$h = -10 \rightarrow 5$

$T_{\min} = 0.429$, $T_{\max} = 0.805$

$k = -26 \rightarrow 29$

12626 measured reflections

$l = -14 \rightarrow 14$

Refinement

Refinement on F^2

Primary atom site location: structure-invariant direct methods

Least-squares matrix: full

Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.059$

Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.160$

H-atom parameters constrained

$S = 1.07$

$w = 1/[\sigma^2(F_o^2) + (0.0832P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

4781 reflections

$(\Delta/\sigma)_{\max} = 0.001$

280 parameters

$\Delta\rho_{\max} = 0.64$ e Å⁻³

0 restraints

$\Delta\rho_{\min} = -0.44$ e Å⁻³

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.35786 (5)	0.135609 (18)	0.69095 (4)	0.04072 (18)
S1	0.5487 (2)	0.10524 (7)	0.30631 (12)	0.0930 (5)
S2	0.69634 (16)	0.05042 (5)	1.01610 (12)	0.0705 (4)
N1	0.4648 (4)	0.21527 (13)	0.7228 (3)	0.0395 (7)
N2	0.1730 (3)	0.18662 (12)	0.6333 (3)	0.0376 (7)
N3	0.1684 (4)	0.07761 (13)	0.6675 (3)	0.0477 (8)
N4	0.4448 (5)	0.12136 (17)	0.5232 (4)	0.0681 (11)
N5	0.5115 (4)	0.09198 (15)	0.8049 (3)	0.0567 (9)
N6	-0.4985 (4)	0.30689 (15)	0.3820 (3)	0.0545 (9)
C1	0.6201 (4)	0.22617 (17)	0.7760 (4)	0.0459 (9)

H1	0.6878	0.1950	0.8032	0.055*
C2	0.6827 (5)	0.28129 (18)	0.7918 (4)	0.0517 (10)
H2	0.7899	0.2875	0.8306	0.062*
C3	0.5841 (5)	0.32709 (17)	0.7493 (4)	0.0544 (11)
H3	0.6247	0.3648	0.7579	0.065*
C4	0.4239 (5)	0.31726 (16)	0.6935 (4)	0.0473 (10)
H4	0.3556	0.3481	0.6649	0.057*
C5	0.3680 (4)	0.26104 (15)	0.6813 (3)	0.0369 (8)
C6	0.1981 (4)	0.24383 (15)	0.6274 (3)	0.0362 (8)
C7	0.0730 (4)	0.28084 (15)	0.5800 (3)	0.0394 (8)
H7	0.0929	0.3204	0.5741	0.047*
C8	-0.0849 (4)	0.25821 (15)	0.5408 (3)	0.0378 (8)
C9	-0.1082 (4)	0.19839 (16)	0.5516 (3)	0.0414 (9)
H9	-0.2116	0.1820	0.5272	0.050*
C10	0.0237 (4)	0.16374 (15)	0.5987 (3)	0.0396 (8)
C11	0.0177 (5)	0.10032 (16)	0.6220 (4)	0.0475 (10)
C12	-0.1211 (6)	0.0675 (2)	0.6069 (5)	0.0726 (15)
H12	-0.2220	0.0839	0.5749	0.087*
C13	-0.1112 (7)	0.0091 (2)	0.6397 (6)	0.094 (2)
H13	-0.2052	-0.0140	0.6326	0.113*
C14	0.0431 (7)	-0.0137 (2)	0.6833 (6)	0.097 (2)
H14	0.0545	-0.0529	0.7031	0.116*
C15	0.1769 (6)	0.02138 (17)	0.6969 (5)	0.0661 (13)
H15	0.2791	0.0057	0.7280	0.079*
C16	-0.4894 (5)	0.36367 (19)	0.4057 (4)	0.0554 (11)
H16	-0.5805	0.3866	0.3774	0.066*
C17	-0.3527 (5)	0.39026 (18)	0.4697 (4)	0.0550 (11)
H17	-0.3514	0.4301	0.4835	0.066*
C18	-0.2186 (5)	0.35648 (16)	0.5128 (4)	0.0482 (10)
H18	-0.1246	0.3734	0.5557	0.058*
C19	-0.2235 (4)	0.29699 (15)	0.4923 (3)	0.0374 (8)
C20	-0.3672 (5)	0.27481 (17)	0.4255 (4)	0.0480 (10)
H20	-0.3720	0.2351	0.4102	0.058*
C22	0.4845 (5)	0.11376 (17)	0.4333 (4)	0.0441 (9)
C23	0.5890 (5)	0.07416 (16)	0.8920 (4)	0.0468 (9)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.0325 (3)	0.0397 (3)	0.0473 (3)	0.00621 (19)	-0.0019 (2)	0.00460 (19)
S1	0.1100 (13)	0.1146 (12)	0.0590 (8)	0.0514 (10)	0.0277 (8)	0.0161 (8)
S2	0.0649 (8)	0.0655 (8)	0.0726 (8)	-0.0015 (6)	-0.0159 (6)	0.0226 (6)
N1	0.0289 (15)	0.0434 (17)	0.0445 (17)	0.0030 (13)	0.0009 (13)	0.0023 (13)
N2	0.0282 (15)	0.0349 (15)	0.0471 (18)	0.0007 (12)	-0.0026 (13)	-0.0006 (13)
N3	0.0423 (18)	0.0389 (17)	0.057 (2)	0.0000 (14)	-0.0059 (16)	0.0046 (15)
N4	0.069 (3)	0.073 (3)	0.064 (3)	0.017 (2)	0.015 (2)	0.001 (2)
N5	0.045 (2)	0.055 (2)	0.065 (2)	0.0071 (17)	-0.0078 (18)	0.0152 (17)
N6	0.0328 (17)	0.061 (2)	0.066 (2)	0.0045 (16)	-0.0062 (16)	0.0049 (17)

supplementary materials

C1	0.0250 (18)	0.052 (2)	0.057 (2)	0.0032 (16)	-0.0053 (17)	0.0087 (18)
C2	0.0277 (18)	0.062 (3)	0.062 (3)	-0.0025 (18)	-0.0034 (18)	0.004 (2)
C3	0.039 (2)	0.045 (2)	0.077 (3)	-0.0086 (19)	0.003 (2)	-0.002 (2)
C4	0.0312 (19)	0.040 (2)	0.069 (3)	0.0007 (17)	0.0017 (18)	0.0014 (18)
C5	0.0270 (17)	0.044 (2)	0.0386 (19)	0.0037 (15)	0.0016 (15)	0.0003 (15)
C6	0.0280 (17)	0.0418 (19)	0.0378 (19)	0.0031 (15)	0.0021 (15)	-0.0020 (15)
C7	0.0333 (19)	0.0378 (19)	0.045 (2)	0.0023 (15)	-0.0012 (16)	0.0035 (15)
C8	0.0298 (18)	0.041 (2)	0.041 (2)	0.0018 (15)	-0.0014 (15)	-0.0007 (15)
C9	0.0291 (18)	0.042 (2)	0.049 (2)	-0.0011 (16)	-0.0045 (16)	-0.0009 (16)
C10	0.0327 (19)	0.039 (2)	0.045 (2)	0.0014 (16)	-0.0003 (16)	0.0026 (16)
C11	0.044 (2)	0.039 (2)	0.055 (2)	-0.0036 (17)	-0.0060 (19)	0.0045 (17)
C12	0.048 (3)	0.057 (3)	0.102 (4)	-0.009 (2)	-0.022 (3)	0.015 (3)
C13	0.069 (3)	0.052 (3)	0.146 (5)	-0.022 (3)	-0.029 (4)	0.024 (3)
C14	0.084 (4)	0.047 (3)	0.144 (6)	-0.011 (3)	-0.034 (4)	0.033 (3)
C15	0.061 (3)	0.042 (2)	0.088 (3)	0.003 (2)	-0.012 (3)	0.014 (2)
C16	0.036 (2)	0.063 (3)	0.064 (3)	0.016 (2)	0.000 (2)	0.015 (2)
C17	0.047 (2)	0.041 (2)	0.077 (3)	0.0076 (19)	0.008 (2)	0.007 (2)
C18	0.034 (2)	0.049 (2)	0.057 (2)	-0.0013 (17)	-0.0050 (18)	0.0067 (18)
C19	0.0278 (17)	0.0392 (19)	0.044 (2)	0.0032 (15)	0.0019 (15)	0.0036 (15)
C20	0.035 (2)	0.048 (2)	0.057 (2)	0.0032 (17)	-0.0060 (18)	-0.0001 (18)
C22	0.036 (2)	0.044 (2)	0.050 (2)	0.0082 (17)	0.0011 (18)	0.0021 (18)
C23	0.036 (2)	0.038 (2)	0.065 (3)	-0.0003 (17)	0.0044 (19)	0.0053 (18)

Geometric parameters (\AA , $^\circ$)

Cu1—N1	2.040 (3)	C5—C6	1.485 (4)
Cu1—N2	1.948 (3)	C6—C7	1.375 (5)
Cu1—N3	2.036 (3)	C7—C8	1.403 (5)
Cu1—N4	2.143 (4)	C7—H7	0.9300
Cu1—N5	1.931 (3)	C8—C9	1.398 (5)
S1—C22	1.609 (5)	C8—C19	1.482 (5)
S2—C23	1.622 (4)	C9—C10	1.382 (5)
N1—C1	1.344 (4)	C9—H9	0.9300
N1—C5	1.357 (4)	C10—C11	1.485 (5)
N2—C10	1.335 (4)	C11—C12	1.356 (6)
N2—C6	1.336 (4)	C12—C13	1.395 (6)
N3—C15	1.334 (5)	C12—H12	0.9300
N3—C11	1.366 (5)	C13—C14	1.388 (7)
N4—C22	1.122 (5)	C13—H13	0.9300
N5—C23	1.154 (5)	C14—C15	1.353 (6)
N6—C16	1.333 (5)	C14—H14	0.9300
N6—C20	1.334 (5)	C15—H15	0.9300
C1—C2	1.370 (5)	C16—C17	1.378 (6)
C1—H1	0.9300	C16—H16	0.9300
C2—C3	1.369 (5)	C17—C18	1.373 (5)
C2—H2	0.9300	C17—H17	0.9300
C3—C4	1.384 (5)	C18—C19	1.388 (5)
C3—H3	0.9300	C18—H18	0.9300
C4—C5	1.373 (5)	C19—C20	1.392 (5)

C4—H4	0.9300	C20—H20	0.9300
N1—Cu1—N2	79.0 (1)	C8—C7—H7	120.4
N1—Cu1—N3	155.7 (1)	C9—C8—C7	118.1 (3)
N1—Cu1—N4	95.6 (1)	C9—C8—C19	121.2 (3)
N1—Cu1—N5	97.6 (1)	C7—C8—C19	120.8 (3)
N2—Cu1—N3	79.1 (1)	C10—C9—C8	119.5 (3)
N2—Cu1—N4	98.7 (1)	C10—C9—H9	120.3
N2—Cu1—N5	158.0 (2)	C8—C9—H9	120.3
N3—Cu1—N4	98.1 (2)	N2—C10—C9	120.9 (3)
N3—Cu1—N5	98.7 (1)	N2—C10—C11	113.0 (3)
N4—Cu1—N5	103.2 (2)	C9—C10—C11	126.1 (3)
C1—N1—C5	118.1 (3)	C12—C11—N3	121.9 (4)
C1—N1—Cu1	126.7 (3)	C12—C11—C10	125.1 (4)
C5—N1—Cu1	115.1 (2)	N3—C11—C10	112.9 (3)
C10—N2—C6	121.1 (3)	C11—C12—C13	119.4 (4)
C10—N2—Cu1	119.5 (2)	C11—C12—H12	120.3
C6—N2—Cu1	119.5 (2)	C13—C12—H12	120.3
C15—N3—C11	118.2 (3)	C14—C13—C12	118.0 (5)
C15—N3—Cu1	126.5 (3)	C14—C13—H13	121.0
C11—N3—Cu1	115.3 (2)	C12—C13—H13	121.0
C22—N4—Cu1	177.4 (4)	C15—C14—C13	119.7 (4)
C23—N5—Cu1	164.0 (4)	C15—C14—H14	120.2
C16—N6—C20	116.9 (3)	C13—C14—H14	120.2
N1—C1—C2	122.8 (3)	N3—C15—C14	122.7 (4)
N1—C1—H1	118.6	N3—C15—H15	118.6
C2—C1—H1	118.6	C14—C15—H15	118.6
C3—C2—C1	118.6 (3)	N6—C16—C17	123.7 (4)
C3—C2—H2	120.7	N6—C16—H16	118.2
C1—C2—H2	120.7	C17—C16—H16	118.2
C2—C3—C4	120.0 (4)	C18—C17—C16	118.4 (4)
C2—C3—H3	120.0	C18—C17—H17	120.8
C4—C3—H3	120.0	C16—C17—H17	120.8
C5—C4—C3	118.6 (4)	C17—C18—C19	120.0 (4)
C5—C4—H4	120.7	C17—C18—H18	120.0
C3—C4—H4	120.7	C19—C18—H18	120.0
N1—C5—C4	121.9 (3)	C18—C19—C20	116.8 (3)
N1—C5—C6	113.3 (3)	C18—C19—C8	122.2 (3)
C4—C5—C6	124.8 (3)	C20—C19—C8	121.0 (3)
N2—C6—C7	121.2 (3)	N6—C20—C19	124.2 (4)
N2—C6—C5	112.6 (3)	N6—C20—H20	117.9
C7—C6—C5	126.1 (3)	C19—C20—H20	117.9
C6—C7—C8	119.3 (3)	N4—C22—S1	177.1 (5)
C6—C7—H7	120.4	N5—C23—S2	178.6 (4)
N5—Cu1—N1—C1	18.8 (3)	N1—C5—C6—N2	1.7 (4)
N2—Cu1—N1—C1	176.8 (3)	C4—C5—C6—N2	-176.0 (3)
N3—Cu1—N1—C1	150.6 (3)	N1—C5—C6—C7	178.9 (3)
N4—Cu1—N1—C1	-85.4 (3)	C4—C5—C6—C7	1.2 (6)
N5—Cu1—N1—C5	-163.6 (3)	N2—C6—C7—C8	2.0 (5)

supplementary materials

N2—Cu1—N1—C5	-5.6 (2)	C5—C6—C7—C8	-175.0 (3)
N3—Cu1—N1—C5	-31.8 (4)	C6—C7—C8—C9	-0.1 (5)
N4—Cu1—N1—C5	92.2 (3)	C6—C7—C8—C19	178.4 (3)
N5—Cu1—N2—C10	-91.2 (4)	C7—C8—C9—C10	-0.7 (5)
N3—Cu1—N2—C10	-4.8 (3)	C19—C8—C9—C10	-179.1 (3)
N1—Cu1—N2—C10	-174.1 (3)	C6—N2—C10—C9	2.3 (5)
N4—Cu1—N2—C10	91.8 (3)	Cu1—N2—C10—C9	-176.6 (3)
N5—Cu1—N2—C6	89.9 (4)	C6—N2—C10—C11	-175.0 (3)
N3—Cu1—N2—C6	176.3 (3)	Cu1—N2—C10—C11	6.0 (4)
N1—Cu1—N2—C6	6.9 (3)	C8—C9—C10—N2	-0.4 (6)
N4—Cu1—N2—C6	-87.1 (3)	C8—C9—C10—C11	176.6 (4)
N5—Cu1—N3—C15	-16.7 (4)	C15—N3—C11—C12	0.3 (7)
N2—Cu1—N3—C15	-174.5 (4)	Cu1—N3—C11—C12	-176.8 (4)
N1—Cu1—N3—C15	-148.3 (4)	C15—N3—C11—C10	177.2 (4)
N4—Cu1—N3—C15	88.1 (4)	Cu1—N3—C11—C10	0.0 (4)
N5—Cu1—N3—C11	160.2 (3)	N2—C10—C11—C12	173.0 (4)
N2—Cu1—N3—C11	2.4 (3)	C9—C10—C11—C12	-4.1 (7)
N1—Cu1—N3—C11	28.5 (5)	N2—C10—C11—N3	-3.7 (5)
N4—Cu1—N3—C11	-95.0 (3)	C9—C10—C11—N3	179.1 (4)
N2—Cu1—N5—C23	-15.6 (15)	N3—C11—C12—C13	0.7 (8)
N3—Cu1—N5—C23	-98.2 (12)	C10—C11—C12—C13	-175.7 (5)
N1—Cu1—N5—C23	63.7 (12)	C11—C12—C13—C14	-2.1 (10)
N4—Cu1—N5—C23	161.3 (12)	C12—C13—C14—C15	2.5 (11)
C5—N1—C1—C2	1.3 (6)	C11—N3—C15—C14	0.1 (8)
Cu1—N1—C1—C2	178.8 (3)	Cu1—N3—C15—C14	176.9 (5)
N1—C1—C2—C3	-1.5 (7)	C13—C14—C15—N3	-1.5 (10)
C1—C2—C3—C4	1.1 (7)	C20—N6—C16—C17	1.2 (7)
C2—C3—C4—C5	-0.5 (7)	N6—C16—C17—C18	-0.7 (7)
C1—N1—C5—C4	-0.7 (5)	C16—C17—C18—C19	-0.5 (6)
Cu1—N1—C5—C4	-178.5 (3)	C17—C18—C19—C20	1.2 (6)
C1—N1—C5—C6	-178.5 (3)	C17—C18—C19—C8	-177.6 (4)
Cu1—N1—C5—C6	3.7 (4)	C9—C8—C19—C18	160.4 (4)
C3—C4—C5—N1	0.4 (6)	C7—C8—C19—C18	-18.0 (6)
C3—C4—C5—C6	177.9 (4)	C9—C8—C19—C20	-18.2 (6)
C10—N2—C6—C7	-3.1 (5)	C7—C8—C19—C20	163.3 (4)
Cu1—N2—C6—C7	175.8 (3)	C16—N6—C20—C19	-0.4 (6)
C10—N2—C6—C5	174.2 (3)	C18—C19—C20—N6	-0.7 (6)
Cu1—N2—C6—C5	-6.8 (4)	C8—C19—C20—N6	178.0 (4)

Fig. 1

