

4-Methylbenzyl 4-aminobenzoate

Ali Haider,^a Zareen Akhter,^{a*} Mohammad Saif Ullah Khan,^a Michael Bolte^b and Humaira M. Siddiqi^a

^aDepartment of Chemistry, Quaid-I-Azam University, Islamabad 45320, Pakistan, and ^bInstitut für Anorganische Chemie, J. W. Goethe-Universität Frankfurt, Max-von-Laue-Strasse 7, 60438 Frankfurt/Main, Germany
Correspondence e-mail: zareenakhter@yahoo.com


Received 10 February 2010; accepted 24 February 2010

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.043; wR factor = 0.094; data-to-parameter ratio = 7.6.

The dihedral angle between the two benzene rings in the title compound, $\text{C}_{15}\text{H}_{15}\text{NO}_2$, is $65.28(12)^\circ$. The crystal structure is stabilized by $\text{N}-\text{H}\cdots\text{N}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, leading to the formation of supramolecular chains along the a -axis direction.

Related literature

For the reduction of aryl-nitro compounds, see: Tafesh & Weiguny (1996); Vass *et al.* (2001); Entwistle *et al.* (1977); Bavín (1958); Yuste *et al.* (1982); Idrees *et al.* (2009). For the uses of amines, see: Kumarraja & Pitchumani (2004).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{15}\text{NO}_2$
 $M_r = 241.28$
Monoclinic, $P2_1$
 $a = 8.2097(12)$ Å
 $b = 5.5344(5)$ Å
 $c = 14.293(2)$ Å
 $\beta = 98.531(12)^\circ$

$V = 642.24(14)$ Å³
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.08$ mm⁻¹
 $T = 173$ K
 $0.27 \times 0.13 \times 0.13$ mm

Data collection

Stoe IPDSII two-circle diffractometer
4021 measured reflections

1322 independent reflections
960 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.093$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.094$
 $S = 0.91$
1322 reflections
173 parameters
1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.14$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.13$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1A}\cdots\text{O2}^{\text{i}}$	0.88 (4)	2.12 (5)	2.977 (4)	164 (4)
$\text{N1}-\text{H1B}\cdots\text{N1}^{\text{ii}}$	0.96 (6)	2.37 (6)	3.278 (3)	158 (4)

Symmetry codes: (i) $x - 1, y, z$; (ii) $-x, y - \frac{1}{2}, -z + 2$.

Data collection: *X-AREA* (Stoe & Cie, 2001); cell refinement: *X-AREA*; data reduction: *X-AREA*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors are grateful to the Department of Chemistry, Quaid-I-Azam University, Islamabad, Pakistan, and the Institute for Inorganic Chemistry, University of Frankfurt, Germany, for providing laboratory and analytical facilities. They also thank the Higher Education Commission of Pakistan for financial support through Project No. 20-723/ R&D/ 06/191.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2629).

References

- Bavín, P. M. G. (1958). *Can. J. Chem.* **36**, 238–241.
Entwistle, I. D., Jackson, A. E., Johnston, R. A. & Telford, R. P. (1977). *J. Chem. Soc. Perkin Trans. 1*, pp. 443–444.
Idrees, G. A., Abuo-Rahma, G., Aly, O. M. & Radwan, M. F. (2009). *Eur. J. Med. Chem.* **44**, 2679–2684.
Kumarraja, M. & Pitchumani, K. (2004). *J. Appl. Catal. A*, **265**, 135–139.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Stoe & Cie (2001). *X-AREA*. Stoe & Cie, Darmstadt, Germany.
Tafesh, M. & Weiguny, J. (1996). *Chem. Rev.* **96**, 2035–2052.
Vass, A., Dudas, J., Toth, J. & Varma, R. S. (2001). *Tetrahedron Lett.* **42**, 5347–5349.
Yuste, F., Saldana, M. & Walls, F. (1982). *Tetrahedron Lett.* **23**, 147–148.

supplementary materials

Acta Cryst. (2010). E66, o736 [doi:10.1107/S1600536810007075]

4-Methylbenzyl 4-aminobenzoate

A. Haider, Z. Akhter, M. Khan, M. Bolte and H. M. Siddiqi

Comment

Reduction of aryl-nitro compounds to their corresponding amines is an important chemical transformation in synthetic organic chemistry mainly due to the fact that the amino group can serve as the site for further derivatization (Tafesh *et al.*, 1996; Vass *et al.*, 2001). Amines are important intermediates in the production of many pharmaceuticals, photographic materials, agrochemicals, polymers, dyes, and rubber materials (Kumarraja & Pitchumani, 2004). Selective reduction nitro-aromatics to amines can be achieved by hydrogen transfer using Pt—C (Entwistle *et al.*, 1977), Pd—C (Bavin *et al.*, 1958) and Raney Ni (Yuste *et al.*, 1982) catalysts. Most commonly applied or reported methods are direct catalytic hydrogenation and catalytic hydrazine reduction. The reduction of 1,4-bis(4-nitrobenzoyloxymethyl) benzene has been carried out using the catalytic hydrogenation method. It is important to note that the process requires much care in the addition of hydrazine, in order to prevent the breakdown of the ester linkage, as hydrazides may be formed from carboxylic esters in the absence of the catalyst or even if the catalyst is not properly charged (Idrees *et al.*, 2009). The limited addition of the hydrazine in the presence of activated catalyst can also cause the breakage of ester linkage not from the aryl carbon but from the acyl carbon as proved by the crystal structure of the title compound, (I). Herein, the synthesis and the crystal structure of (I) are reported.

The dihedral angle between the two benzene rings in (I) is 65.28 (12)°. The crystal structure is stabilized by N—H···N and N—H···O hydrogen bonds, Table 1, which lead to supramolecular chains along the *a* direction.

Experimental

Compound (I) was synthesized in two steps. In the first step, a mixture of 1,4-bis(chloromethyl)benzene Aldrich; 2.00 g, 0.0114 mol), anhydrous K₂CO₃ (3.154 g, 0.0229 mol) and 4-nitrobenzoic acid (3.824 g, 0.0229 mol) were added to a two neck round bottom flask charged with DMF (50 ml). This was heated at 393 K for 12 h under a nitrogen atmosphere. After cooling to room temperature, the reaction mixture was poured into water (800 ml) to precipitate a yellow solid which was washed thoroughly with water and then separated by filtration. In the second step a 250 ml two neck flask was charged with the just synthesised yellow solid (1.00 g, 2.84 mmol) and was refluxed in ethanol with 5% palladium on carbon (Pd/C, 0.06 g), followed by the drop-wise addition of hydrated hydrazine (80%) diluted in ethanol. The mixture was refluxed for 8 h and then filtered to remove Pd/C. The solvent was evaporated and the resulting crude solid was recrystallized from ethanol to afford crystals (yield:68%, m.pt.: 397 K).

Refinement

Hydrogen atoms bonded to C were included in calculated positions [C—H = 0.95–0.99 Å] and refined as riding [$U_{iso}(H) = 1.2–1.5U_{eq}(C)$]. The H atoms bonded to N were isotropically refined. Due to the absence of anomalous scatterers, the absolute structure could not be determined and 773 Friedel pairs were merged.

Figures




Fig. 1. Perspective view of (I) with the atom numbering scheme. The displacement ellipsoids are at the 50% probability level and H atoms are drawn as small spheres of arbitrary radii.

4-Methylbenzyl 4-aminobenzoate

Crystal data

$C_{15}H_{15}NO_2$

$M_r = 241.28$

Monoclinic, $P2_1$

Hall symbol: P 2yb

$a = 8.2097$ (12) Å

$b = 5.5344$ (5) Å

$c = 14.293$ (2) Å

$\beta = 98.531$ (12)°

$V = 642.24$ (14) Å³

$Z = 2$

$F(000) = 256$

$D_x = 1.248$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2492 reflections

$\theta = 4.0$ – 25.9 °

$\mu = 0.08$ mm⁻¹

$T = 173$ K

Prism, colourless

$0.27 \times 0.13 \times 0.13$ mm

Data collection

Stoe IPDSII two-circle diffractometer

Radiation source: fine-focus sealed tube

graphite

ω scans

4021 measured reflections

1322 independent reflections

960 reflections with $I > 2\sigma(I)$

$R_{int} = 0.093$

$\theta_{max} = 25.7$ °, $\theta_{min} = 3.5$ °

$h = -9 \rightarrow 9$

$k = -6 \rightarrow 5$

$l = -17 \rightarrow 17$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.043$

$wR(F^2) = 0.094$

$S = 0.91$

1322 reflections

173 parameters

1 restraint

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0405P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{max} = 0.028$

$\Delta\rho_{max} = 0.14$ e Å⁻³

$\Delta\rho_{min} = -0.13$ e Å⁻³

Extinction correction: SHELXL97 (Sheldrick, 2008),

$F_c^* = kF_c[1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Primary atom site location: structure-invariant direct methods Extinction coefficient: 0.077 (11)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.0100 (4)	0.0360 (6)	0.9396 (2)	0.0442 (8)
H1A	-0.086 (5)	0.090 (9)	0.911 (3)	0.057 (12)*
H1B	0.007 (7)	-0.130 (11)	0.958 (3)	0.090 (17)*
O1	0.5634 (3)	0.6085 (5)	0.77036 (17)	0.0454 (6)
O2	0.7240 (3)	0.3098 (5)	0.83952 (16)	0.0468 (7)
C1	0.7064 (5)	0.7251 (7)	0.7398 (3)	0.0475 (9)
H1C	0.8064	0.6833	0.7845	0.057*
H1D	0.6922	0.9027	0.7412	0.057*
C2	0.5874 (4)	0.4043 (6)	0.8206 (2)	0.0347 (8)
C11	0.7290 (4)	0.6485 (6)	0.6420 (2)	0.0383 (8)
C12	0.8155 (4)	0.4400 (6)	0.6259 (2)	0.0414 (9)
H12	0.8593	0.3410	0.6779	0.050*
C13	0.8388 (4)	0.3743 (6)	0.5354 (2)	0.0405 (9)
H13	0.8994	0.2321	0.5265	0.049*
C14	0.7755 (4)	0.5117 (6)	0.4576 (2)	0.0405 (8)
C15	0.6888 (5)	0.7204 (7)	0.4742 (3)	0.0511 (11)
H15	0.6446	0.8195	0.4224	0.061*
C16	0.6659 (4)	0.7853 (7)	0.5643 (3)	0.0464 (9)
H16	0.6053	0.9276	0.5733	0.056*
C17	0.7999 (6)	0.4368 (9)	0.3599 (3)	0.0601 (12)
H17A	0.6988	0.3612	0.3278	0.090*
H17B	0.8909	0.3210	0.3639	0.090*
H17C	0.8258	0.5792	0.3241	0.090*
C21	0.4371 (4)	0.3093 (6)	0.8495 (2)	0.0337 (8)
C22	0.4413 (4)	0.0941 (6)	0.9016 (2)	0.0345 (8)
H22	0.5425	0.0099	0.9172	0.041*
C23	0.3008 (4)	0.0031 (6)	0.9306 (2)	0.0376 (8)
H23	0.3064	-0.1430	0.9659	0.045*
C24	0.1503 (4)	0.1226 (6)	0.90885 (19)	0.0321 (7)
C25	0.1451 (4)	0.3382 (7)	0.8578 (2)	0.0373 (8)
H25	0.0437	0.4225	0.8429	0.045*

supplementary materials

C26	0.2854 (4)	0.4302 (6)	0.82882 (19)	0.0352 (8)
H26	0.2796	0.5774	0.7943	0.042*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0384 (18)	0.047 (2)	0.0480 (17)	-0.0004 (15)	0.0081 (14)	0.0099 (15)
O1	0.0416 (14)	0.0390 (13)	0.0604 (14)	0.0023 (12)	0.0232 (11)	0.0028 (12)
O2	0.0303 (13)	0.0573 (16)	0.0533 (14)	0.0020 (13)	0.0077 (11)	-0.0018 (13)
C1	0.048 (2)	0.036 (2)	0.064 (2)	-0.0084 (18)	0.0270 (18)	-0.0041 (17)
C2	0.0344 (19)	0.0332 (18)	0.0375 (16)	0.0003 (16)	0.0084 (14)	-0.0069 (15)
C11	0.0316 (18)	0.0346 (19)	0.0508 (18)	-0.0047 (15)	0.0133 (14)	-0.0020 (15)
C12	0.042 (2)	0.040 (2)	0.0413 (16)	0.0086 (17)	0.0056 (14)	0.0014 (15)
C13	0.040 (2)	0.0333 (19)	0.0489 (19)	0.0037 (15)	0.0086 (15)	-0.0030 (15)
C14	0.037 (2)	0.038 (2)	0.0455 (17)	-0.0061 (17)	0.0059 (15)	-0.0006 (16)
C15	0.044 (2)	0.047 (2)	0.060 (2)	-0.0007 (19)	0.0001 (18)	0.0148 (18)
C16	0.040 (2)	0.0305 (18)	0.073 (2)	0.0047 (17)	0.0207 (17)	0.0053 (18)
C17	0.067 (3)	0.069 (3)	0.0439 (19)	-0.011 (2)	0.0063 (18)	-0.004 (2)
C21	0.0356 (18)	0.0357 (18)	0.0300 (14)	0.0023 (16)	0.0059 (13)	-0.0047 (14)
C22	0.0321 (18)	0.0365 (18)	0.0349 (15)	0.0051 (16)	0.0054 (13)	-0.0051 (14)
C23	0.043 (2)	0.0371 (18)	0.0314 (16)	0.0012 (17)	0.0011 (14)	-0.0023 (13)
C24	0.0346 (18)	0.0328 (17)	0.0295 (14)	-0.0026 (16)	0.0069 (13)	-0.0048 (14)
C25	0.0319 (17)	0.045 (2)	0.0349 (16)	0.0060 (17)	0.0061 (13)	0.0029 (15)
C26	0.039 (2)	0.0351 (18)	0.0328 (15)	0.0049 (16)	0.0090 (14)	0.0021 (14)

Geometric parameters (\AA , $^\circ$)

N1—C24	1.378 (5)	C14—C17	1.499 (5)
N1—H1A	0.88 (4)	C15—C16	1.377 (5)
N1—H1B	0.96 (6)	C15—H15	0.9500
O1—C2	1.338 (4)	C16—H16	0.9500
O1—C1	1.462 (4)	C17—H17A	0.9800
O2—C2	1.230 (4)	C17—H17B	0.9800
C1—C11	1.499 (5)	C17—H17C	0.9800
C1—H1C	0.9900	C21—C22	1.402 (5)
C1—H1D	0.9900	C21—C26	1.406 (4)
C2—C21	1.457 (5)	C22—C23	1.378 (5)
C11—C16	1.379 (5)	C22—H22	0.9500
C11—C12	1.392 (5)	C23—C24	1.395 (5)
C12—C13	1.384 (5)	C23—H23	0.9500
C12—H12	0.9500	C24—C25	1.396 (5)
C13—C14	1.383 (5)	C25—C26	1.378 (5)
C13—H13	0.9500	C25—H25	0.9500
C14—C15	1.395 (5)	C26—H26	0.9500
C24—N1—H1A	118 (3)	C15—C16—C11	121.5 (3)
C24—N1—H1B	119 (3)	C15—C16—H16	119.3
H1A—N1—H1B	113 (5)	C11—C16—H16	119.3
C2—O1—C1	118.2 (3)	C14—C17—H17A	109.5

O1—C1—C11	111.7 (3)	C14—C17—H17B	109.5
O1—C1—H1C	109.3	H17A—C17—H17B	109.5
C11—C1—H1C	109.3	C14—C17—H17C	109.5
O1—C1—H1D	109.3	H17A—C17—H17C	109.5
C11—C1—H1D	109.3	H17B—C17—H17C	109.5
H1C—C1—H1D	107.9	C22—C21—C26	117.9 (3)
O2—C2—O1	122.3 (3)	C22—C21—C2	120.1 (3)
O2—C2—C21	124.5 (3)	C26—C21—C2	122.0 (3)
O1—C2—C21	113.2 (3)	C23—C22—C21	121.0 (3)
C16—C11—C12	117.5 (3)	C23—C22—H22	119.5
C16—C11—C1	120.8 (3)	C21—C22—H22	119.5
C12—C11—C1	121.7 (3)	C22—C23—C24	120.8 (3)
C13—C12—C11	121.1 (3)	C22—C23—H23	119.6
C13—C12—H12	119.5	C24—C23—H23	119.6
C11—C12—H12	119.5	N1—C24—C23	121.2 (3)
C14—C13—C12	121.3 (3)	N1—C24—C25	120.1 (3)
C14—C13—H13	119.4	C23—C24—C25	118.6 (3)
C12—C13—H13	119.4	C26—C25—C24	120.7 (3)
C13—C14—C15	117.3 (3)	C26—C25—H25	119.6
C13—C14—C17	120.7 (3)	C24—C25—H25	119.6
C15—C14—C17	122.0 (3)	C25—C26—C21	121.0 (3)
C16—C15—C14	121.3 (3)	C25—C26—H26	119.5
C16—C15—H15	119.4	C21—C26—H26	119.5
C14—C15—H15	119.4		
C2—O1—C1—C11	94.1 (4)	O2—C2—C21—C22	-0.9 (4)
C1—O1—C2—O2	-1.9 (5)	O1—C2—C21—C22	179.1 (3)
C1—O1—C2—C21	178.1 (3)	O2—C2—C21—C26	177.6 (3)
O1—C1—C11—C16	95.1 (4)	O1—C2—C21—C26	-2.4 (4)
O1—C1—C11—C12	-85.7 (4)	C26—C21—C22—C23	0.7 (4)
C16—C11—C12—C13	0.8 (5)	C2—C21—C22—C23	179.2 (3)
C1—C11—C12—C13	-178.5 (4)	C21—C22—C23—C24	0.0 (4)
C11—C12—C13—C14	-0.8 (5)	C22—C23—C24—N1	-178.3 (3)
C12—C13—C14—C15	0.7 (5)	C22—C23—C24—C25	-0.7 (4)
C12—C13—C14—C17	-179.2 (4)	N1—C24—C25—C26	178.2 (3)
C13—C14—C15—C16	-0.6 (5)	C23—C24—C25—C26	0.6 (4)
C17—C14—C15—C16	179.3 (4)	C24—C25—C26—C21	0.2 (4)
C14—C15—C16—C11	0.7 (6)	C22—C21—C26—C25	-0.8 (4)
C12—C11—C16—C15	-0.7 (5)	C2—C21—C26—C25	-179.3 (3)
C1—C11—C16—C15	178.5 (4)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1A \cdots O2 ⁱ	0.88 (4)	2.12 (5)	2.977 (4)	164 (4)
N1—H1B \cdots N1 ⁱⁱ	0.96 (6)	2.37 (6)	3.278 (3)	158 (4)

Symmetry codes: (i) $x-1, y, z$; (ii) $-x, y-1/2, -z+2$.

Fig. 1

