

[2,6-Bis(dimethylaminomethyl)phenyl]-selenium bromide monohydrate

Richard A. Varga,^{a*} Monica Kulcsar^b and Anca Silvestru^a

^aFaculty of Chemistry and Chemical Engineering, Babes-Bolyai University, Arany Janos Str. no. 11, RO-400028, Cluj Napoca, Romania, and ^bDepartamento de Química Inorgánica, Instituto de Ciencia de, Materiales de Aragón, Universidad de Zaragoza—CSIC, E-50009 Zaragoza, Spain
Correspondence e-mail: richy@chem.ubbcluj.ro


Received 4 February 2010; accepted 2 March 2010

Key indicators: single-crystal X-ray study; $T = 297$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.039; wR factor = 0.086; data-to-parameter ratio = 18.5.

In the title hydrated molecular salt, $\text{C}_{12}\text{H}_{19}\text{N}_2\text{Se}^+\cdot\text{Br}^-\cdot\text{H}_2\text{O}$, the two independent bromide anions lie on a twofold rotation axis. Strong intramolecular $\text{N}\rightarrow\text{Se}$ interactions [2.185 (3) and 2.181 (3) Å] are established by both N atoms of the organic group in the cation, in *trans* positions to each other, with an $\text{N}-\text{Se}-\text{N}$ angle of $161.6(1)^\circ$, resulting in a T-shaped (C,N,N')Se core. In the crystal, dimeric associations are formed by $\text{Br}\cdots\text{Se}$ [3.662 (2) Å] and $\text{Br}\cdots\text{H}$ interactions [2.56 (6) and 2.63 (7) Å] involving two bromide anions, two cations and two water molecules.

Related literature

For related selenium and tellurium compounds, see: Drake *et al.* (2001a,b); Deleanu *et al.* (2002); Kulcsar *et al.* (2005, 2007); Beleaga *et al.* (2009); Fujihara *et al.* (1995). For van der Waals radii, see: Emsley (1994).



Experimental

Crystal data

$\text{C}_{12}\text{H}_{19}\text{N}_2\text{Se}^+\cdot\text{Br}^-\cdot\text{H}_2\text{O}$

$M_r = 368.18$

Monoclinic, $C2/c$

$a = 15.1494(14)$ Å

$b = 11.3182(10)$ Å

$c = 18.8083(17)$ Å

$\beta = 110.475(2)^\circ$

$V = 3021.2(5)$ Å³

$Z = 8$

Mo $K\alpha$ radiation

$\mu = 5.12$ mm⁻¹

$T = 297$ K

$0.31 \times 0.29 \times 0.09$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer

Absorption correction: multi-scan (SADABS; Bruker, 2000)

$T_{\min} = 0.300$, $T_{\max} = 0.656$

11856 measured reflections

3093 independent reflections

2696 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.050$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$

$wR(F^2) = 0.086$

$S = 1.09$

3093 reflections

167 parameters

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 0.49$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.53$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1 \cdots Br2	0.78 (6)	2.56 (6)	3.340 (5)	175 (5)
O1—H2 \cdots Br1	0.78 (6)	2.63 (7)	3.406 (5)	176 (7)

Data collection: SMART (Bruker, 2000); cell refinement: SAINT-Plus (Bruker, 2001); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and DIAMOND 3 (Brandenburg & Putz, 2006); software used to prepare material for publication: publCIF (Westrip, 2010).

This work was supported by the Romanian Ministry of Education and Research (PNII Program, grant 2404/2008). We also thank the National Center for X-ray Diffraction, Cluj-Napoca, for the structure determination.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2538).

References

- Beleaga, A., Kulcsar, M., Deleanu, C., Nicolescu, A., Silvestru, C. & Silvestru, A. (2009). *J. Organomet. Chem.* **694**, 1308–1316.
- Brandenburg, K. & Putz, H. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2000). SMART and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2001). SAINT-Plus. Bruker AXS Inc., Madison, Wisconsin, USA.
- Deleanu, C., Drake, J. E., Hursthouse, M. B., Kulcsar, M., Leight, M. E. & Silvestru, A. (2002). *Appl. Organomet. Chem.* **16**, 727–731.
- Drake, J. E., Hursthouse, M. B., Kulcsar, M., Leight, M. E. & Silvestru, A. (2001a). *J. Organomet. Chem.* **623**, 153–160.
- Drake, J. E., Hursthouse, M. B., Kulcsar, M., Leight, M. E. & Silvestru, A. (2001b). *Phosphorus Sulfur*, **169**, 293–296.
- Emsley, J. (1994). In *Die Elemente*. Berlin: Walter de Gruyter.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Fujihara, H., Mima, H. & Furukawa, N. (1995). *J. Am. Chem. Soc.* **117**, 10153–10154.
- Kulcsar, M., Beleaga, A., Silvestru, C., Nicolescu, A., Deleanu, C., Todasca, C. & Silvestru, A. (2007). *Dalton Trans.* pp. 2187–2196.
- Kulcsar, M., Silvestru, A., Silvestru, C., Drake, J. E., Macdonald, C. L. B., Hursthouse, M. B. & Leight, M. E. (2005). *J. Organomet. Chem.* **690**, 3217–3228.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2010). *publCIF*. In preparation.

supplementary materials

Acta Cryst. (2010). E66, o771 [doi:10.1107/S1600536810008019]

[2,6-Bis(dimethylaminomethyl)phenyl]selenium bromide monohydrate

R. A. Varga, M. Kulcsar and A. Silvestru

Comment

In the last years our interest was focused on the synthesis, structural characterization and chemical reactivity of some new hypervalent organoselenium and organotellurium derivatives containing aryl groups with pendant arms, e.g. 2-(Me₂NCH₂)C₆H₄, 2-[O(CH₂CH₂)₂NCH₂]C₆H₄ and 2-[MeN(CH₂CH₂)₂NCH₂]C₆H₄ (Drake *et al.*, 2001a,b, Deleanu *et al.*, 2002, Kulcsar *et al.*, 2005, 2007, Beleaga *et al.*, 2009).

The crystal of the title compound contains a mixture of [2,6-(Me₂NCH₂)₂C₆H₃]Se⁺ cations and [Br]⁻ anions and crystallizes with a water molecule. The two independent bromine anions lie on a two-fold rotation axis. Both pendant arms of the organic group attached to selenium establish strong intramolecular N→Se interactions [Se—N1 = 2.185 (3) Å, Se1—N2 = 2.181 (3) Å], *trans* one to the other [N1—Se1—N2 = 161.6 (1)°], thus resulting in the increase of the coordination number at Se to three (Fig. 1). The Se—N distances are of the same magnitude as found in the cation of [2,6-(Me₂NCH₂)₂C₆H₃]Se⁺[PF₆]⁻ [Se1—N1 = 2.180 Å; Se1—N2 = 2.154 Å] (Fujihara *et al.*, 1995). This results in a distorted T-shaped (C,N,N)Se core [C1—Se1—N1 = 80.5 (1)°, C1—Se1—N2 = 81.2 (1)°], the distortion being mainly due to the constraints imposed by the two SeC₃N five-membered chelate rings.

An unusual dimer association is formed between two cations, two bromine atoms and two water molecules (Fig. 2). The Br2 atom is involved in bridging two selenium atoms and the interatomic Se1—Br2 distances [3.662 (2) Å] are much longer than the sum of the corresponding covalent radii [$\Sigma r_{\text{cov}}(\text{Se},\text{Br})$ ca. 2.31 Å], but shorter than the sum of the van der Waals radii [$\Sigma r_{\text{vdW}}(\text{Se},\text{Br})$ ca. 3.95 Å] (Emsley, 1994), consistent with an electrostatic anion-cation interaction. This interaction is directed *trans* to the selenium-carbon bond in the cation [C1—Se1—Br2 = 154.4 (1)°], thus resulting in a distorted square-planar environment around the chalcogen atom. The water molecules bridge the Br1 and Br2 anions through Br⋯H hydrogen bonding [Br1⋯H1 = 2.56 (6) Å, Br2⋯H2 = 2.63 (7) Å].

Experimental

[2,6-(Me₂NCH₂)₂C₆H₃]SeBr was obtained by oxidizing [2,6-(Me₂NCH₂)₂C₆H₃]₂Se₂ with elemental bromine.

The attempt to grow crystals of [2,6-(Me₂NCH₂)₂C₆H₃]SeBr from a methylene dichloride / n-hexane mixture (1:5, v/v) in open atmosphere led to the isolation of [2,6-(Me₂NCH₂)₂C₆H₃]SeBr.H₂O.

Refinement

All hydrogen atoms were placed in calculated positions using a riding model, with C—H = 0.93–0.97 Å and with U_{iso} = 1.5U_{eq}(C) for methyl H and U_{iso} = 1.2U_{eq}(C) for aryl H. The methyl groups were allowed to rotate but not to tip. Hydrogen atoms from the water molecule were found from difference map and refined to O—H distances of 0.78 (6) Å.

Figures




Fig. 1. : View of title compound showing the atom-numbering scheme at 30% probability thermal ellipsoids.




Fig. 2. : Intra- and intermolecular interactions in the title compound (dashed lines; only H atoms involved in interactions are shown) [symmetry code: (i) = 1-x, y, 1.5-z].

[2,6-Bis(dimethylaminomethyl)phenyl]selenium bromide monohydrate

Crystal data

$C_{12}H_{19}N_2Se^+ \cdot Br^- \cdot H_2O$

$M_r = 368.18$

Monoclinic, $C2/c$

Hall symbol: $-C\ 2yc$

$a = 15.1494\ (14)\ \text{\AA}$

$b = 11.3182\ (10)\ \text{\AA}$

$c = 18.8083\ (17)\ \text{\AA}$

$\beta = 110.475\ (2)^\circ$

$V = 3021.2\ (5)\ \text{\AA}^3$

$Z = 8$

$F(000) = 1472$

$D_x = 1.619\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 3226 reflections

$\theta = 2.3\text{--}25.2^\circ$

$\mu = 5.12\ \text{mm}^{-1}$

$T = 297\ \text{K}$

Block, colourless

$0.31 \times 0.29 \times 0.09\ \text{mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

phi and ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2000)

$T_{\min} = 0.300$, $T_{\max} = 0.656$

11856 measured reflections

3093 independent reflections

2696 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.050$

$\theta_{\max} = 26.4^\circ$, $\theta_{\min} = 2.3^\circ$

$h = -18 \rightarrow 18$

$k = -14 \rightarrow 14$

$l = -23 \rightarrow 23$

Refinement

Refinement on F^2

Least-squares matrix: full

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

$$R[F^2 > 2\sigma(F^2)] = 0.039$$

$$wR(F^2) = 0.086$$

$$S = 1.09$$

3093 reflections

167 parameters

0 restraints

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0287P)^2 + 3.8997P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.49 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.53 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.5000	0.26510 (5)	0.7500	0.05461 (17)
Br2	0.5000	0.73811 (6)	0.7500	0.0747 (2)
C1	0.3337 (2)	0.9300 (3)	0.93573 (18)	0.0319 (7)
C2	0.3774 (2)	0.9329 (3)	1.01348 (18)	0.0372 (7)
C6	0.2603 (2)	1.0046 (3)	0.89677 (19)	0.0346 (7)
C10	0.2169 (2)	0.9828 (3)	0.81308 (19)	0.0408 (8)
H10A	0.1640	0.9290	0.8028	0.049*
H10B	0.1939	1.0565	0.7868	0.049*
C7	0.4589 (3)	0.8503 (3)	1.04538 (19)	0.0444 (8)
H7A	0.5173	0.8906	1.0505	0.053*
H7B	0.4622	0.8232	1.0952	0.053*
C8	0.3799 (3)	0.6604 (3)	1.0050 (3)	0.0572 (11)
H8A	0.4077	0.6257	1.0544	0.086*
H8B	0.3677	0.5999	0.9670	0.086*
H8C	0.3218	0.6984	1.0014	0.086*
C9	0.5358 (3)	0.6906 (4)	0.9999 (2)	0.0561 (10)
H9A	0.5780	0.7482	0.9919	0.084*
H9B	0.5246	0.6294	0.9624	0.084*
H9C	0.5635	0.6570	1.0496	0.084*
C12	0.3495 (3)	1.0235 (3)	0.7709 (2)	0.0495 (9)
H12A	0.4006	0.9872	0.7599	0.074*
H12B	0.3744	1.0733	0.8147	0.074*
H12C	0.3126	1.0702	0.7282	0.074*

supplementary materials

C11	0.2469 (3)	0.8570 (4)	0.7181 (2)	0.0525 (9)
H11A	0.2098	0.9056	0.6767	0.079*
H11B	0.2073	0.7982	0.7286	0.079*
H11C	0.2957	0.8189	0.7050	0.079*
C5	0.2301 (3)	1.0860 (3)	0.9381 (2)	0.0454 (9)
H5	0.1810	1.1374	0.9134	0.054*
C4	0.2730 (3)	1.0911 (3)	1.0160 (2)	0.0505 (9)
H4	0.2525	1.1462	1.0434	0.061*
C3	0.3460 (3)	1.0155 (3)	1.0540 (2)	0.0473 (9)
H3	0.3740	1.0198	1.1065	0.057*
N1	0.4453 (2)	0.7484 (2)	0.99307 (16)	0.0385 (7)
N2	0.28962 (19)	0.9310 (2)	0.78604 (15)	0.0372 (6)
O1	0.4714 (3)	0.5053 (4)	0.8489 (2)	0.0793 (11)
Se1	0.37778 (2)	0.82185 (3)	0.879443 (18)	0.03542 (12)
H1	0.480 (4)	0.562 (5)	0.828 (3)	0.09 (2)*
H2	0.475 (5)	0.451 (6)	0.825 (4)	0.12 (3)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0623 (4)	0.0495 (3)	0.0552 (4)	0.000	0.0246 (3)	0.000
Br2	0.0914 (5)	0.0596 (4)	0.0986 (6)	0.000	0.0651 (5)	0.000
C1	0.0345 (17)	0.0321 (16)	0.0323 (17)	-0.0077 (13)	0.0157 (14)	-0.0033 (13)
C2	0.0410 (19)	0.0415 (18)	0.0301 (18)	-0.0085 (15)	0.0139 (15)	0.0012 (14)
C6	0.0337 (17)	0.0329 (16)	0.0367 (18)	-0.0046 (13)	0.0117 (14)	-0.0018 (13)
C10	0.0355 (18)	0.0434 (19)	0.0379 (19)	0.0018 (15)	0.0057 (15)	0.0008 (15)
C7	0.043 (2)	0.057 (2)	0.0302 (18)	-0.0040 (17)	0.0092 (15)	0.0064 (16)
C8	0.045 (2)	0.050 (2)	0.073 (3)	-0.0058 (17)	0.016 (2)	0.021 (2)
C9	0.040 (2)	0.065 (3)	0.061 (3)	0.0149 (19)	0.0143 (19)	0.014 (2)
C12	0.056 (2)	0.053 (2)	0.040 (2)	-0.0079 (18)	0.0165 (18)	0.0080 (17)
C11	0.058 (2)	0.055 (2)	0.036 (2)	-0.0003 (19)	0.0076 (18)	-0.0115 (17)
C5	0.047 (2)	0.0364 (18)	0.056 (2)	0.0009 (16)	0.0221 (18)	-0.0025 (16)
C4	0.063 (2)	0.043 (2)	0.055 (2)	-0.0016 (19)	0.033 (2)	-0.0114 (18)
C3	0.060 (2)	0.054 (2)	0.0317 (19)	-0.0089 (19)	0.0210 (18)	-0.0050 (16)
N1	0.0319 (15)	0.0426 (16)	0.0395 (16)	0.0017 (12)	0.0108 (13)	0.0081 (12)
N2	0.0412 (16)	0.0396 (15)	0.0273 (14)	-0.0040 (12)	0.0076 (12)	-0.0005 (12)
O1	0.106 (3)	0.077 (3)	0.046 (2)	-0.007 (2)	0.0145 (19)	0.0004 (19)
Se1	0.03654 (19)	0.03788 (19)	0.03126 (19)	0.00273 (14)	0.01113 (14)	-0.00128 (14)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.379 (4)	C9—H9B	0.9600
C1—C6	1.384 (5)	C9—H9C	0.9600
C1—Se1	1.887 (3)	C12—N2	1.478 (4)
C2—C3	1.391 (5)	C12—H12A	0.9600
C2—C7	1.497 (5)	C12—H12B	0.9600
C6—C5	1.382 (5)	C12—H12C	0.9600
C6—C10	1.499 (5)	C11—N2	1.474 (4)
C10—N2	1.487 (4)	C11—H11A	0.9600

C10—H10A	0.9700	C11—H11B	0.9600
C10—H10B	0.9700	C11—H11C	0.9600
C7—N1	1.483 (5)	C5—C4	1.381 (5)
C7—H7A	0.9700	C5—H5	0.9300
C7—H7B	0.9700	C4—C3	1.383 (5)
C8—N1	1.476 (4)	C4—H4	0.9300
C8—H8A	0.9600	C3—H3	0.9300
C8—H8B	0.9600	N1—Se1	2.185 (3)
C8—H8C	0.9600	N2—Se1	2.180 (3)
C9—N1	1.484 (5)	O1—H1	0.78 (6)
C9—H9A	0.9600	O1—H2	0.78 (6)
C2—C1—C6	123.1 (3)	H12A—C12—H12B	109.5
C2—C1—Se1	118.5 (2)	N2—C12—H12C	109.5
C6—C1—Se1	118.4 (2)	H12A—C12—H12C	109.5
C1—C2—C3	117.9 (3)	H12B—C12—H12C	109.5
C1—C2—C7	115.7 (3)	N2—C11—H11A	109.5
C3—C2—C7	126.3 (3)	N2—C11—H11B	109.5
C5—C6—C1	118.1 (3)	H11A—C11—H11B	109.5
C5—C6—C10	126.1 (3)	N2—C11—H11C	109.5
C1—C6—C10	115.6 (3)	H11A—C11—H11C	109.5
N2—C10—C6	108.6 (3)	H11B—C11—H11C	109.5
N2—C10—H10A	110.0	C4—C5—C6	119.9 (3)
C6—C10—H10A	110.0	C4—C5—H5	120.1
N2—C10—H10B	110.0	C6—C5—H5	120.1
C6—C10—H10B	110.0	C5—C4—C3	121.2 (3)
H10A—C10—H10B	108.4	C5—C4—H4	119.4
N1—C7—C2	108.4 (3)	C3—C4—H4	119.4
N1—C7—H7A	110.0	C4—C3—C2	119.8 (3)
C2—C7—H7A	110.0	C4—C3—H3	120.1
N1—C7—H7B	110.0	C2—C3—H3	120.1
C2—C7—H7B	110.0	C8—N1—C7	111.6 (3)
H7A—C7—H7B	108.4	C8—N1—C9	109.8 (3)
N1—C8—H8A	109.5	C7—N1—C9	112.1 (3)
N1—C8—H8B	109.5	C8—N1—Se1	107.2 (2)
H8A—C8—H8B	109.5	C7—N1—Se1	105.19 (19)
N1—C8—H8C	109.5	C9—N1—Se1	110.7 (2)
H8A—C8—H8C	109.5	C11—N2—C12	110.4 (3)
H8B—C8—H8C	109.5	C11—N2—C10	111.5 (3)
N1—C9—H9A	109.5	C12—N2—C10	111.4 (3)
N1—C9—H9B	109.5	C11—N2—Se1	109.6 (2)
H9A—C9—H9B	109.5	C12—N2—Se1	108.2 (2)
N1—C9—H9C	109.5	C10—N2—Se1	105.46 (19)
H9A—C9—H9C	109.5	H1—O1—H2	107 (6)
H9B—C9—H9C	109.5	C1—Se1—N2	81.21 (12)
N2—C12—H12A	109.5	C1—Se1—N1	80.47 (12)
N2—C12—H12B	109.5	N2—Se1—N1	161.56 (11)
C6—C1—C2—C3	-0.4 (5)	C2—C7—N1—Se1	34.9 (3)
Se1—C1—C2—C3	178.1 (2)	C6—C10—N2—C11	152.9 (3)

supplementary materials

C6—C1—C2—C7	-176.8 (3)	C6—C10—N2—C12	-83.3 (3)
Se1—C1—C2—C7	1.8 (4)	C6—C10—N2—Se1	33.9 (3)
C2—C1—C6—C5	0.4 (5)	C2—C1—Se1—N2	-166.9 (3)
Se1—C1—C6—C5	-178.1 (2)	C6—C1—Se1—N2	11.7 (2)
C2—C1—C6—C10	-175.7 (3)	C2—C1—Se1—N1	15.2 (2)
Se1—C1—C6—C10	5.7 (4)	C6—C1—Se1—N1	-166.2 (3)
C5—C6—C10—N2	155.8 (3)	C11—N2—Se1—C1	-146.0 (3)
C1—C6—C10—N2	-28.4 (4)	C12—N2—Se1—C1	93.5 (2)
C1—C2—C7—N1	-26.9 (4)	C10—N2—Se1—C1	-25.8 (2)
C3—C2—C7—N1	157.1 (3)	C11—N2—Se1—N1	-139.3 (3)
C1—C6—C5—C4	-0.1 (5)	C12—N2—Se1—N1	100.2 (4)
C10—C6—C5—C4	175.6 (3)	C10—N2—Se1—N1	-19.1 (4)
C6—C5—C4—C3	-0.2 (6)	C8—N1—Se1—C1	90.8 (2)
C5—C4—C3—C2	0.2 (6)	C7—N1—Se1—C1	-28.1 (2)
C1—C2—C3—C4	0.1 (5)	C9—N1—Se1—C1	-149.4 (3)
C7—C2—C3—C4	176.0 (3)	C8—N1—Se1—N2	84.2 (4)
C2—C7—N1—C8	-81.0 (3)	C7—N1—Se1—N2	-34.8 (4)
C2—C7—N1—C9	155.3 (3)	C9—N1—Se1—N2	-156.1 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1 \cdots Br2	0.78 (6)	2.56 (6)	3.340 (5)	175 (5)
O1—H2 \cdots Br1	0.78 (6)	2.63 (7)	3.406 (5)	176 (7)

Fig. 1




Fig. 2

