

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-[(*E*)-(3,4-Dimethylisoxazol-5-yl)imino-methyl]phenolHoong-Kun Fun,^{a*} Madhukar Hemamalini,^a Abdullah M. Asiri,^{b§} Salman A. Khan^b and Khalid A. Khan^b^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bDepartment of Chemistry, Faculty of Science, King Abdul Aziz University, Jeddah, Saudi Arabia

Correspondence e-mail: hkfun@usm.my


Received 24 February 2010; accepted 3 March 2010

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.070; wR factor = 0.203; data-to-parameter ratio = 16.2.

The title compound, $\text{C}_{12}\text{H}_{12}\text{N}_2\text{O}_2$, has been synthesized by the reaction of 5-amino-3,4-dimethylisoxazole and salicylaldehyde. The molecule adopts an *E* configuration about the central $\text{C}=\text{N}$ double bond. The dihedral angle between the isoxazole and phenyl rings is $4.2(2)^\circ$ and an intramolecular $\text{O}-\text{H}\cdots\text{N}$ hydrogen bond generates an *S*(6) ring motif. The crystal studied was a non-merohedral twin with a domain ratio of 0.834(4):0.166(4).

Related literature

For background to the biological and pharmacological properties of oxazole derivatives, see: Spinelli (1999); Conti *et al.* (1998); Mishra *et al.* (1998); Ko *et al.* (1998); Kang *et al.* (2000); Huang & Chen (2005). For details of hydrogen bonding and hydrogen-bond motifs, see: Jeffrey & Saenger (1991); Bernstein *et al.* (1995); Jeffrey (1997); Scheiner (1997). For the stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986).



Experimental

Crystal data

 $\text{C}_{12}\text{H}_{12}\text{N}_2\text{O}_2$
 $M_r = 216.24$ Triclinic, $P\bar{1}$
 $a = 5.3475(14)$ Å

* Thomson Reuters ResearcherID: A-3561-2009.

§ On secondment to: The Center of Excellence for Advanced Materials Research, King Abdul Aziz University, Jeddah 21589, Saudi Arabia.

 $b = 8.615(2)$ Å
 $c = 12.321(3)$ Å
 $\alpha = 103.696(5)^\circ$
 $\beta = 91.486(5)^\circ$
 $\gamma = 94.059(5)^\circ$
 $V = 549.6(2)$ Å³ $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 100$ K
 $0.56 \times 0.14 \times 0.08$ mm

Data collection

Bruker APEX DUO CCD area-detector diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2009)
 $T_{\min} = 0.951$, $T_{\max} = 0.993$ 2467 measured reflections
2467 independent reflections
1946 reflections with $I > 2\sigma(I)$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.070$
 $wR(F^2) = 0.203$
 $S = 1.06$
2467 reflections
152 parametersH atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.40$ e Å⁻³
 $\Delta\rho_{\min} = -0.34$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

| $D-\text{H}\cdots A$ | $D-\text{H}$ | $\text{H}\cdots A$ | $D\cdots A$ | $D-\text{H}\cdots A$ |
|--|--------------|--------------------|-------------|----------------------|
| $\text{O2}-\text{H1O2}\cdots\text{N1}$ | 1.00 (9) | 1.71 (8) | 2.648 (5) | 154 (8) |

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

HKF and MH thank the Malaysian Government and Universiti Sains Malaysia for the Research University Golden Goose grant No. 1001/PFIZIK/811012. MH thanks Universiti Sains Malaysia for a post-doctoral research fellowship. AMA, SAK and KAK thank the Chemistry Department, King Abdul Aziz University, Jeddah, Saudi Arabia, for providing research facilities. AMA would also like to thank the deanship of scientific research at KAU for grant No. 171/428.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ2738).

References

- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Conti, P., Dallanoce, C., Amici, M. D., Micheli, C. D. & Klotz, K. N. (1998). *Bioorg. Med. Chem.* **6**, 401–409.
Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
Huang, L. & Chen, D.-B. (2005). *Acta Cryst.* **E61**, o4169–o4170.
Jeffrey, G. A. (1997). *An Introduction to Hydrogen Bonding*. Oxford University Press.
Jeffrey, G. A. & Saenger, W. (1991). *Hydrogen Bonding in Biological Structures*. Berlin: Springer.
Kang, Y. Y., Shin, K. L., Yoo, K. H., Seo, K. J., Hong, C. Y., Lee, C. S., Park, S. Y., Kim, D. J. & Park, S. W. (2000). *Bioorg. Med. Chem. Lett.* **10**, 95–99.
Ko, D. H., Maponya, M. F., Khalil, M. A., Oriaku, E. T. & You, Z. (1998). *J. Med. Chem. Res.* **8**, 313–318.
Mishra, A., Jain, K. J. & Asthana, J. G. (1998). *Oriental J. Chem.* **14**, 151–152.

Scheiner, S. (1997). *Hydrogen Bonding, A Theoretical Perspective*. Oxford University Press.

Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

Spinelli, O. A. (1999). *Ital. Soc. Chem.* **3** 301–339.

supplementary materials

Acta Cryst. (2010). E66, o773-o774 [doi:10.1107/S1600536810008160]

2-[(*E*)-(3,4-Dimethylisoxazol-5-yl)iminomethyl]phenol

H.-K. Fun, M. Hemamalini, A. M. Asiri, S. A. Khan and K. A. Khan

Comment

Heterocyclic compounds, especially isoxazoles, are one of the key building elements of natural products. Among the numerous heterocyclic systems of biological and pharmacological interest, the oxazole ring is endowed with various activities, including hypoglycemic (Spinelli, 1999), analgesic (Conti *et al.*, 1998), anti-inflammatory (Mishra *et al.*, 1998), anti-bacterial (Ko *et al.*, 1998) and anti-tumor (Kang *et al.*, 2000) properties. In view of the importance of the title compound as a pharmaceutical intermediate, the paper reports its synthesis and crystal structure.

In the title compound (Fig. 1), the isoxazole ring is essentially planar with a maximum deviation of 0.002 (2) Å for atom C8. The dihedral angle between the isoxazole ring (N2/O1/C8–C10) and the phenyl ring (C1–C6) is 4.30 (15)°. The methyl groups at C9 and C10 deviate from the isoxazole mean plane by 0.056 (3) Å and 0.013 (4) Å, respectively. The C5—O2 and C7=N1 bond lengths are 1.353 (4) Å and 1.293 (4) Å, respectively, and agree with the corresponding values in 4-[(1*E*)-(3,5-dibromo-2-hydroxyphenyl)methylene]-amino}-1,5-dimethyl-2-phenyl-1,2-dihydro-3H-pyrazol-3-one [1.344 (3) and 1.292 (4) Å; Huang & Chen, 2005].

In the crystal structure (Fig. 2), the imino N atoms are linked to the phenol O atoms and act as hydrogen-bond acceptors in intramolecular O2—H1O2···N1 interactions (Table 1) (Jeffrey & Saenger, 1991; Jeffrey, 1997; Scheiner, 1997), which generate S(6) ring motifs (Bernstein *et al.*, 1995).

Experimental

A mixture of 5-amino-3,4-dimethylisoxazole (0.50g, 0.0044 mol) and salicylaldehyde (0.54g, 0.0044 mol) in methanol (15 mL) was refluxed for 5 h with stirring to give a light yellow precipitate. It was then filtered and washed with methanol to give the pure Schiff base. Yield: 68%; mp. 116°C. IR (KBr) ν_{\max} cm⁻¹: 2922(C—H), 1594 (C=O), 1562 (C=C), 1152 (C—N). ¹H NMR (CDCl₃) δ : 8.89 (s, 1H, CH olefinic), 7.42 (d, H3, J=1.8Hz), 7.44 (dd, H4, J=7.8Hz), 7.02 (dd, H5, J=7.8Hz), 6.97 (d, H6, J=1.2 Hz), 2.25 (s, CH3), 2.05 (s, CH3).

Refinement

Atom H1O2 was located from the difference Fourier map and refined freely. The remaining hydrogen atoms were positioned geometrically [C—H = 0.93 Å or 0.96 Å] and were refined using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5 U_{\text{eq}}(\text{C})$. A rotating group model was applied to the methyl groups. The crystal is a non-merohedral twin with BASF = 0.166 (4).

Figures




Fig. 1. The asymmetric unit of the title compound, showing 50% probability displacement ellipsoids and the atom-numbering scheme.




Fig. 2. The crystal packing of the title compound (I).

2-[(E)-(3,4-Dimethylisoxazol-5-yl)iminomethyl]phenol

Crystal data

$C_{12}H_{12}N_2O_2$

$M_r = 216.24$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 5.3475$ (14) Å

$b = 8.615$ (2) Å

$c = 12.321$ (3) Å

$\alpha = 103.696$ (5)°

$\beta = 91.486$ (5)°

$\gamma = 94.059$ (5)°

$V = 549.6$ (2) Å³

$Z = 2$

$F(000) = 228$

$D_x = 1.307$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2862 reflections

$\theta = 2.4$ – 29.7 °

$\mu = 0.09$ mm⁻¹

$T = 100$ K

Plate, yellow

$0.56 \times 0.14 \times 0.08$ mm

Data collection

Bruker APEX DUO CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

φ and ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2009)

$T_{\min} = 0.951$, $T_{\max} = 0.993$

2467 measured reflections

2467 independent reflections

1946 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.000$

$\theta_{\max} = 27.5$ °, $\theta_{\min} = 1.7$ °

$h = -6 \rightarrow 6$

$k = -11 \rightarrow 10$

$l = -6 \rightarrow 15$

Refinement

Refinement on F^2

Least-squares matrix: full

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

$$R[F^2 > 2\sigma(F^2)] = 0.070$$

$$wR(F^2) = 0.203$$

$$S = 1.06$$

2467 reflections

152 parameters

0 restraints

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0657P)^2 + 1.1217P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.40 \text{ e } \text{Å}^{-3}$$

$$\Delta\rho_{\min} = -0.34 \text{ e } \text{Å}^{-3}$$

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|------|------------|------------|--------------|----------------------------------|
| O1 | 1.0387 (4) | 1.1305 (2) | 0.41739 (17) | 0.0228 (5) |
| O2 | 0.6135 (4) | 0.7350 (3) | 0.08828 (18) | 0.0262 (5) |
| N1 | 0.8330 (5) | 0.9521 (3) | 0.2591 (2) | 0.0194 (5) |
| N2 | 1.2481 (5) | 1.2478 (3) | 0.4414 (2) | 0.0252 (6) |
| C1 | 0.2930 (5) | 0.7387 (3) | 0.3499 (3) | 0.0201 (6) |
| H1A | 0.3082 | 0.7892 | 0.4256 | 0.024* |
| C2 | 0.0960 (5) | 0.6240 (3) | 0.3108 (3) | 0.0221 (6) |
| H2A | -0.0224 | 0.5990 | 0.3593 | 0.026* |
| C3 | 0.0782 (6) | 0.5468 (4) | 0.1976 (3) | 0.0249 (7) |
| H3A | -0.0529 | 0.4691 | 0.1710 | 0.030* |
| C4 | 0.2512 (6) | 0.5831 (4) | 0.1235 (3) | 0.0250 (7) |
| H4A | 0.2362 | 0.5296 | 0.0484 | 0.030* |
| C5 | 0.4491 (5) | 0.7008 (3) | 0.1626 (2) | 0.0199 (6) |
| C6 | 0.4711 (5) | 0.7802 (3) | 0.2771 (2) | 0.0186 (6) |
| C7 | 0.6723 (5) | 0.9024 (3) | 0.3225 (2) | 0.0191 (6) |
| H7A | 0.6859 | 0.9462 | 0.3992 | 0.023* |
| C8 | 1.0225 (5) | 1.0682 (3) | 0.3052 (2) | 0.0176 (6) |
| C9 | 1.2091 (5) | 1.1366 (3) | 0.2539 (2) | 0.0192 (6) |
| C10 | 1.3428 (5) | 1.2479 (3) | 0.3446 (2) | 0.0195 (6) |
| C11 | 1.2629 (6) | 1.1010 (4) | 0.1327 (3) | 0.0276 (7) |
| H11A | 1.1457 | 1.0163 | 0.0921 | 0.041* |

supplementary materials

| | | | | |
|------|------------|------------|------------|-------------|
| H11C | 1.4305 | 1.0685 | 0.1229 | 0.041* |
| H11D | 1.2470 | 1.1953 | 0.1051 | 0.041* |
| C12 | 1.5687 (5) | 1.3582 (4) | 0.3386 (3) | 0.0243 (7) |
| H12A | 1.6191 | 1.4202 | 0.4124 | 0.036* |
| H12D | 1.5285 | 1.4287 | 0.2920 | 0.036* |
| H12B | 1.7034 | 1.2963 | 0.3075 | 0.036* |
| H1O2 | 0.722 (11) | 0.820 (7) | 0.132 (5) | 0.080 (18)* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|-------------|-------------|-------------|--------------|--------------|-------------|
| O1 | 0.0194 (11) | 0.0238 (11) | 0.0233 (11) | -0.0060 (8) | 0.0019 (8) | 0.0042 (8) |
| O2 | 0.0250 (12) | 0.0281 (11) | 0.0232 (11) | -0.0040 (9) | 0.0061 (9) | 0.0028 (9) |
| N1 | 0.0170 (12) | 0.0148 (11) | 0.0261 (13) | 0.0009 (9) | 0.0020 (9) | 0.0041 (9) |
| N2 | 0.0207 (13) | 0.0246 (13) | 0.0290 (14) | -0.0061 (10) | -0.0003 (10) | 0.0062 (10) |
| C1 | 0.0155 (14) | 0.0176 (13) | 0.0270 (15) | 0.0054 (10) | 0.0036 (11) | 0.0036 (11) |
| C2 | 0.0152 (14) | 0.0204 (14) | 0.0318 (16) | 0.0021 (11) | 0.0054 (11) | 0.0078 (12) |
| C3 | 0.0159 (14) | 0.0225 (14) | 0.0353 (17) | -0.0013 (11) | -0.0012 (12) | 0.0063 (12) |
| C4 | 0.0247 (16) | 0.0227 (15) | 0.0246 (15) | -0.0006 (12) | -0.0007 (12) | 0.0004 (12) |
| C5 | 0.0164 (14) | 0.0194 (14) | 0.0242 (14) | 0.0026 (11) | 0.0021 (11) | 0.0052 (11) |
| C6 | 0.0154 (13) | 0.0155 (13) | 0.0256 (14) | 0.0033 (10) | 0.0019 (11) | 0.0054 (11) |
| C7 | 0.0199 (14) | 0.0158 (13) | 0.0216 (14) | 0.0058 (11) | 0.0019 (11) | 0.0034 (10) |
| C8 | 0.0140 (13) | 0.0163 (13) | 0.0237 (14) | 0.0051 (10) | 0.0029 (10) | 0.0056 (10) |
| C9 | 0.0159 (13) | 0.0169 (13) | 0.0262 (15) | 0.0037 (10) | 0.0033 (11) | 0.0074 (11) |
| C10 | 0.0136 (13) | 0.0169 (13) | 0.0296 (15) | 0.0038 (10) | 0.0020 (11) | 0.0079 (11) |
| C11 | 0.0272 (16) | 0.0291 (16) | 0.0260 (16) | 0.0009 (13) | 0.0080 (12) | 0.0055 (12) |
| C12 | 0.0137 (13) | 0.0215 (14) | 0.0392 (17) | 0.0005 (11) | 0.0038 (12) | 0.0100 (12) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|------------|-----------|-----------|-----------|
| O1—C8 | 1.357 (3) | C4—H4A | 0.9300 |
| O1—N2 | 1.429 (3) | C5—C6 | 1.412 (4) |
| O2—C5 | 1.353 (4) | C6—C7 | 1.452 (4) |
| O2—H1O2 | 0.95 (6) | C7—H7A | 0.9300 |
| N1—C7 | 1.293 (4) | C8—C9 | 1.367 (4) |
| N1—C8 | 1.381 (4) | C9—C10 | 1.426 (4) |
| N2—C10 | 1.308 (4) | C9—C11 | 1.491 (4) |
| C1—C2 | 1.385 (4) | C10—C12 | 1.498 (4) |
| C1—C6 | 1.410 (4) | C11—H11A | 0.9600 |
| C1—H1A | 0.9300 | C11—H11C | 0.9600 |
| C2—C3 | 1.394 (4) | C11—H11D | 0.9600 |
| C2—H2A | 0.9300 | C12—H12A | 0.9600 |
| C3—C4 | 1.387 (4) | C12—H12D | 0.9600 |
| C3—H3A | 0.9300 | C12—H12B | 0.9600 |
| C4—C5 | 1.404 (4) | | |
| C8—O1—N2 | 107.7 (2) | N1—C7—H7A | 119.1 |
| C5—O2—H1O2 | 103 (3) | C6—C7—H7A | 119.1 |
| C7—N1—C8 | 120.3 (2) | O1—C8—C9 | 110.8 (2) |

| | | | |
|--------------|------------|----------------|------------|
| C10—N2—O1 | 105.3 (2) | O1—C8—N1 | 119.7 (2) |
| C2—C1—C6 | 121.2 (3) | C9—C8—N1 | 129.5 (3) |
| C2—C1—H1A | 119.4 | C8—C9—C10 | 103.2 (3) |
| C6—C1—H1A | 119.4 | C8—C9—C11 | 128.4 (3) |
| C1—C2—C3 | 118.9 (3) | C10—C9—C11 | 128.4 (3) |
| C1—C2—H2A | 120.6 | N2—C10—C9 | 112.9 (3) |
| C3—C2—H2A | 120.6 | N2—C10—C12 | 119.8 (3) |
| C4—C3—C2 | 121.6 (3) | C9—C10—C12 | 127.3 (3) |
| C4—C3—H3A | 119.2 | C9—C11—H11A | 109.5 |
| C2—C3—H3A | 119.2 | C9—C11—H11C | 109.5 |
| C3—C4—C5 | 119.7 (3) | H11A—C11—H11C | 109.5 |
| C3—C4—H4A | 120.2 | C9—C11—H11D | 109.5 |
| C5—C4—H4A | 120.2 | H11A—C11—H11D | 109.5 |
| O2—C5—C4 | 118.4 (3) | H11C—C11—H11D | 109.5 |
| O2—C5—C6 | 121.9 (3) | C10—C12—H12A | 109.5 |
| C4—C5—C6 | 119.6 (3) | C10—C12—H12D | 109.5 |
| C1—C6—C5 | 119.0 (3) | H12A—C12—H12D | 109.5 |
| C1—C6—C7 | 118.8 (3) | C10—C12—H12B | 109.5 |
| C5—C6—C7 | 122.2 (3) | H12A—C12—H12B | 109.5 |
| N1—C7—C6 | 121.8 (3) | H12D—C12—H12B | 109.5 |
| C8—O1—N2—C10 | -0.3 (3) | N2—O1—C8—C9 | 0.3 (3) |
| C6—C1—C2—C3 | 1.3 (4) | N2—O1—C8—N1 | -179.4 (2) |
| C1—C2—C3—C4 | -0.6 (5) | C7—N1—C8—O1 | -0.3 (4) |
| C2—C3—C4—C5 | -0.3 (5) | C7—N1—C8—C9 | -179.9 (3) |
| C3—C4—C5—O2 | -179.3 (3) | O1—C8—C9—C10 | -0.2 (3) |
| C3—C4—C5—C6 | 0.5 (4) | N1—C8—C9—C10 | 179.4 (3) |
| C2—C1—C6—C5 | -1.1 (4) | O1—C8—C9—C11 | 178.7 (3) |
| C2—C1—C6—C7 | 179.4 (3) | N1—C8—C9—C11 | -1.7 (5) |
| O2—C5—C6—C1 | 180.0 (3) | O1—N2—C10—C9 | 0.1 (3) |
| C4—C5—C6—C1 | 0.2 (4) | O1—N2—C10—C12 | -179.8 (2) |
| O2—C5—C6—C7 | -0.5 (4) | C8—C9—C10—N2 | 0.0 (3) |
| C4—C5—C6—C7 | 179.7 (3) | C11—C9—C10—N2 | -178.9 (3) |
| C8—N1—C7—C6 | -179.7 (2) | C8—C9—C10—C12 | 180.0 (3) |
| C1—C6—C7—N1 | -176.1 (3) | C11—C9—C10—C12 | 1.1 (5) |
| C5—C6—C7—N1 | 4.4 (4) | | |

Hydrogen-bond geometry (\AA , $^\circ$)

| <i>D</i> —H \cdots <i>A</i> | <i>D</i> —H | H \cdots <i>A</i> | <i>D</i> \cdots <i>A</i> | <i>D</i> —H \cdots <i>A</i> |
|-------------------------------|-------------|---------------------|----------------------------|-------------------------------|
| O2—H1O2 \cdots N1 | 1.00 (9) | 1.71 (8) | 2.648 (5) | 154 (8) |

Fig. 2

