

Ethyl 1-*tert*-butyl-2-(4-methoxyphenyl)-1*H*-benzimidazole-5-carboxylate

Natarajan Arumugam,^a Shafida Abd Hamid,^b Aisyah Saad Abdul Rahim,^{a,‡} Madhukar Hemamalini^c and Hoong-Kun Fun^{c,*§}

^aSchool of Pharmaceutical Sciences, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, ^bKulliyah of Science, International Islamic University Malaysia (IIUM), Jalan Istana, Bandar Indera Mahkota, 25200 Kuantan, Pahang, Malaysia, and ^cX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia

Correspondence e-mail: hkfun@usm.my


Received 25 February 2010; accepted 2 March 2010

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.037; wR factor = 0.113; data-to-parameter ratio = 17.6.

In the title molecule, $\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_3$, the imidazole ring is essentially planar, with a maximum deviation of 0.015 (1) Å. The dihedral angle between the benzene and imidazole rings is 65.47 (6)°. The crystal packing is stabilized by weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds, forming zigzag chains along the c axis. The crystal structure is further stabilized by $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For background to benzimidazole derivatives, their biological activity and medical applications, see: Orjales *et al.* (1997); Andrzejewska *et al.* (2002); Garuti *et al.* (2000); Lukevics *et al.* (2001); Komazin *et al.* (2003). For details of hydrogen bonding, see: Jeffrey & Saenger (1991); Jeffrey (1997); Scheiner (1997). For the stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986).



Experimental

Crystal data

$\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_3$
 $M_r = 352.42$

Orthorhombic, $Pna2_1$
 $a = 14.3963$ (7) Å

‡ Additional correspondence author, e-mail: aisyah@usm.my.
§ Thomson Reuters ResearcherID: A-3561-2009.

$b = 8.6206$ (5) Å
 $c = 15.1609$ (8) Å
 $V = 1881.54$ (17) Å³
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.08$ mm⁻¹
 $T = 100$ K
 $0.53 \times 0.42 \times 0.27$ mm

Data collection

Bruker APEX DUO CCD area-detector diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2009)
 $T_{\min} = 0.957$, $T_{\max} = 0.978$

17426 measured reflections
4233 independent reflections
3951 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.113$
 $S = 1.13$
4233 reflections
240 parameters

1 restraint
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.60$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.57$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$Cg1$, $Cg2$ and $Cg3$ are the centroids of the rings N1,N2,C7–C9 and C1–C6 and C8–C13, respectively.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$C5-H5A\cdots O2^i$	0.93	2.57	3.3822 (16)	146
$C13-H13A\cdots O2^{ii}$	0.93	2.52	3.4116 (16)	160
$C19-H19C\cdots N2^{iii}$	0.96	2.57	3.4976 (19)	164
$C2-H2A\cdots Cg3^{iv}$	0.93	2.75	3.5594 (14)	146
$C18-H18C\cdots Cg2^v$	0.96	2.74	3.6878 (16)	168
$C21-H21B\cdots Cg1^{iv}$	0.96	2.78	3.4703 (16)	130

Symmetry codes: (i) $-x + \frac{1}{2}, y + \frac{1}{2}, z - \frac{1}{2}$; (ii) $x - \frac{1}{2}, -y + \frac{1}{2}, z$; (iii) $x - \frac{1}{2}, -y + \frac{3}{2}, z$; (iv) $x, y + 1, z$; (v) $-x - \frac{1}{2}, y + \frac{3}{2}, z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

NA, SAH and ASAR are grateful to Universiti Sains Malaysia (USM) and the International Islamic University Malaysia (IIUM) for funding the synthetic chemistry work under the USM Research University Grant (1001/PFAR-MASI/815026) and the IIUM Research Endowment Grant (EDW B 0902-206). NA thanks USM for the award of a post-doctoral fellowship. HKF and MH thank the Malaysian Government and USM for the Research University Golden Goose grant No. 1001/PFIZIK/811012. MH also thanks USM for a post-doctoral research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WN2378).

References

- Andrzejewska, M., Yépez-Mulia, L., Cedillo-Rivera, R., Tapia, A., Vilpo, L., Vilpo, J. & Kazimierczuk, Z. (2002). *Eur. J. Med. Chem.* **37**, 973–978.
Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
Garuti, L., Roberti, M., Malagoli, M., Rossi, T. & Castelli, M. (2000). *Bioorg. Med. Chem. Lett.* **10**, 2193–2195.

- Jeffrey, G. A. (1997). *An Introduction to Hydrogen Bonding*. Oxford University Press.
- Jeffrey, G. A. & Saenger, W. (1991). *Hydrogen Bonding in Biological Structures*. Berlin: Springer.
- Komazin, G., Ptak, R. G., Emmer, B. T., Townsend, L. B. & Drach, J. C. (2003). *Nucleosides Nucleotides Nucleic Acids*, **22**, 1725–1727.
- Lukevics, E., Arsenyan, P., Shestakova, I., Domracheva, I., Nesterova, A. & Pudova, O. (2001). *Eur. J. Med. Chem.* **36**, 507–515.
- Orjales, A., Mosquera, R., Labeaga, L. & Rodes, R. (1997). *J. Med. Chem.* **40**, 586–593.
- Scheiner, S. (1997). *Hydrogen Bonding, A Theoretical Perspective*. Oxford University Press.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2010). E66, o776-o777 [doi:10.1107/S1600536810007956]

Ethyl 1-*tert*-butyl-2-(4-methoxyphenyl)-1*H*-benzimidazole-5-carboxylate

N. Arumugam, S. Abd Hamid, A. S. Abdul Rahim, M. Hemamalini and H.-K. Fun

Comment

Benzimidazole derivatives are reported to be physiologically and pharmacologically active and find applications in the treatment of several diseases, such as epilepsy, diabetes and infertility (Orjales *et al.*, 1997). In addition, they also show clinical benefit toward breast cancer (Andrzejewska *et al.*, 2002), leukemia (Garuti *et al.*, 2000), tumor cells (Lukevics *et al.*, 2001) and possess potent antiviral activities (Komazin *et al.*, 2003). We present here the crystal structure of the title compound.

In the asymmetric unit of the title compound (Fig. 1), the imidazole ring is essentially planar, with a maximum deviation of 0.015 (1) Å for atom C8. The dihedral angle between the imidazole ring (N1/N2/C7–C9) and the benzene ring (C1–C6) is 65.47 (6)°.

In the crystal structure (Fig. 2), neighbouring molecules are connected by weak intermolecular C5—H5A···O2, C13—H13A···O2 and C19—H19C···N2 hydrogen bonds (Jeffrey & Saenger, 1991; Jeffrey, 1997; Scheiner, 1997), forming zigzag chains along the *c*-axis. The crystal structure is further stabilized by C—H··· π interactions (Table 1), involving the N1/N2/C7–C9 (centroid Cg1), C1–C6 (centroid Cg2) and C8–C13 (centroid Cg3) rings.

Experimental

Ethyl-3-amino-4-(*tert*-butylamino) benzoate (200 mg, 0.84 mmol) and the sodium metabisulfite adduct of 4-methoxybenzaldehyde (406 mg, 1.68 mmol) were dissolved in DMF. The reaction mixture was irradiated under microwave conditions at 130 °C for 2 minutes. After completion, the reaction mixture was diluted in EtOAc (20 ml) and washed with H₂O (20 ml). The organic layer was collected, dried over Na₂SO₄ and then evaporated in vacuo to yield the crude product. The product was recrystallised from EtOAc as colourless crystals.

Refinement

All hydrogen atoms were positioned geometrically [C—H = 0.93 - 0.97 Å] and were refined using a riding model, with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.5$ for methyl H and 1.2 for all other H atoms. A rotating group model was applied to the methyl groups. In the absence of significant anomalous scattering effects, 3066 Friedel pairs were merged.

Figures




Fig. 1. The asymmetric unit of the title compound, showing 50% probability displacement ellipsoids and the atom-numbering scheme. Hydrogen atoms are shown as spheres of arbitrary radius.




Fig. 2. The crystal packing of the title compound, showing the hydrogen-bonded (dashed lines) network. H atoms not involved in the hydrogen bond interactions have been omitted for clarity.

Ethyl 1-*tert*-butyl-2-(4-methoxyphenyl)-1*H*-benzimidazole-5-carboxylate

Crystal data

$C_{21}H_{24}N_2O_3$

$M_r = 352.42$

Orthorhombic, $Pna2_1$

Hall symbol: P 2c -2n

$a = 14.3963$ (7) Å

$b = 8.6206$ (5) Å

$c = 15.1609$ (8) Å

$V = 1881.54$ (17) Å³

$Z = 4$

$F(000) = 752$

$D_x = 1.244$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 6765 reflections

$\theta = 2.7$ – 36.9°

$\mu = 0.08$ mm⁻¹

$T = 100$ K

Block, colourless

$0.53 \times 0.42 \times 0.27$ mm

Data collection

Bruker APEX DUO CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

φ and ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2009)

$T_{\min} = 0.957$, $T_{\max} = 0.978$

17426 measured reflections

4233 independent reflections

3951 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.032$

$\theta_{\max} = 35.0^\circ$, $\theta_{\min} = 2.7^\circ$

$h = -23 \rightarrow 12$

$k = -12 \rightarrow 13$

$l = -24 \rightarrow 21$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.037$

$wR(F^2) = 0.113$

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$S = 1.13$	$w = 1/[\sigma^2(F_o^2) + (0.0773P)^2]$
4233 reflections	where $P = (F_o^2 + 2F_c^2)/3$
240 parameters	$(\Delta/\sigma)_{\max} = 0.001$
1 restraint	$\Delta\rho_{\max} = 0.60 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.57 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.12374 (8)	1.21773 (11)	0.18521 (7)	0.02061 (19)
O2	0.25619 (7)	0.09272 (11)	0.59077 (8)	0.02019 (19)
O3	0.11256 (7)	0.01389 (12)	0.63146 (8)	0.02132 (19)
N1	0.00387 (7)	0.60189 (11)	0.40798 (7)	0.01307 (17)
N2	0.16125 (7)	0.58244 (12)	0.40296 (7)	0.01439 (18)
C1	0.07671 (9)	0.95165 (14)	0.37017 (8)	0.0159 (2)
H1A	0.0560	0.9522	0.4283	0.019*
C2	0.08488 (9)	1.09163 (14)	0.32461 (9)	0.0158 (2)
H2A	0.0694	1.1849	0.3518	0.019*
C3	0.11669 (9)	1.08946 (13)	0.23757 (8)	0.01465 (19)
C4	0.14432 (9)	0.95031 (14)	0.19836 (8)	0.0156 (2)
H4A	0.1685	0.9503	0.1415	0.019*
C5	0.13556 (8)	0.81218 (14)	0.24449 (8)	0.01461 (19)
H5A	0.1541	0.7195	0.2184	0.018*
C6	0.09897 (8)	0.81128 (13)	0.33013 (8)	0.01312 (18)
C7	0.08802 (8)	0.66340 (14)	0.37851 (8)	0.01293 (19)
C8	0.02836 (8)	0.47467 (13)	0.45937 (8)	0.01276 (18)
C9	0.12598 (8)	0.46271 (13)	0.45378 (8)	0.01270 (18)
C10	0.17417 (8)	0.34437 (14)	0.49659 (8)	0.01382 (19)
H10A	0.2384	0.3360	0.4921	0.017*
C11	0.12284 (8)	0.23910 (13)	0.54632 (8)	0.01365 (18)
C12	0.02572 (8)	0.25487 (14)	0.55447 (8)	0.0156 (2)
H12A	-0.0068	0.1845	0.5892	0.019*
C13	-0.02274 (8)	0.37223 (14)	0.51221 (8)	0.0157 (2)

supplementary materials

H13A	-0.0867	0.3826	0.5187	0.019*
C14	0.17224 (8)	0.10981 (14)	0.59068 (9)	0.01472 (19)
C15	0.14999 (11)	-0.12009 (16)	0.67719 (10)	0.0227 (2)
H15A	0.1157	-0.1375	0.7314	0.027*
H15B	0.2145	-0.1017	0.6923	0.027*
C16	0.14263 (14)	-0.26002 (18)	0.61893 (12)	0.0300 (3)
H16A	0.1631	-0.3501	0.6507	0.045*
H16B	0.1809	-0.2458	0.5677	0.045*
H16C	0.0792	-0.2738	0.6010	0.045*
C17	-0.09468 (8)	0.64670 (15)	0.38647 (9)	0.0157 (2)
C18	-0.14919 (10)	0.50058 (19)	0.36114 (13)	0.0279 (3)
H18A	-0.1544	0.4337	0.4115	0.042*
H18B	-0.1172	0.4471	0.3146	0.042*
H18C	-0.2101	0.5294	0.3414	0.042*
C19	-0.13666 (11)	0.7233 (2)	0.46773 (11)	0.0290 (3)
H19A	-0.0993	0.8106	0.4846	0.044*
H19B	-0.1386	0.6499	0.5153	0.044*
H19C	-0.1985	0.7578	0.4545	0.044*
C20	-0.10132 (11)	0.7557 (2)	0.30761 (12)	0.0305 (4)
H20A	-0.0752	0.8545	0.3229	0.046*
H20B	-0.1653	0.7688	0.2914	0.046*
H20C	-0.0677	0.7124	0.2588	0.046*
C21	0.09176 (12)	1.36157 (16)	0.22065 (10)	0.0232 (3)
H21A	0.0965	1.4409	0.1765	0.035*
H21B	0.1291	1.3894	0.2707	0.035*
H21C	0.0281	1.3511	0.2387	0.035*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0309 (5)	0.0144 (4)	0.0165 (4)	-0.0019 (3)	0.0026 (4)	0.0018 (3)
O2	0.0146 (4)	0.0221 (4)	0.0239 (4)	0.0031 (3)	-0.0012 (3)	0.0034 (4)
O3	0.0189 (4)	0.0182 (4)	0.0269 (5)	0.0009 (3)	0.0004 (4)	0.0079 (4)
N1	0.0101 (4)	0.0151 (4)	0.0141 (4)	0.0004 (3)	0.0000 (3)	0.0013 (3)
N2	0.0117 (4)	0.0141 (4)	0.0174 (4)	-0.0014 (3)	0.0000 (3)	0.0012 (3)
C1	0.0196 (5)	0.0144 (5)	0.0138 (5)	-0.0004 (4)	0.0016 (4)	-0.0007 (4)
C2	0.0195 (5)	0.0135 (5)	0.0145 (5)	-0.0007 (4)	0.0013 (4)	-0.0010 (4)
C3	0.0165 (5)	0.0137 (4)	0.0137 (5)	-0.0022 (4)	-0.0007 (4)	-0.0003 (4)
C4	0.0172 (5)	0.0162 (5)	0.0134 (5)	-0.0025 (4)	0.0015 (4)	-0.0012 (4)
C5	0.0140 (4)	0.0142 (4)	0.0156 (5)	-0.0008 (4)	0.0012 (4)	-0.0019 (4)
C6	0.0126 (4)	0.0127 (4)	0.0140 (4)	-0.0014 (3)	-0.0001 (4)	0.0000 (4)
C7	0.0114 (4)	0.0134 (4)	0.0140 (5)	-0.0009 (3)	-0.0008 (3)	-0.0013 (4)
C8	0.0105 (4)	0.0142 (4)	0.0136 (4)	-0.0001 (3)	0.0007 (3)	0.0006 (4)
C9	0.0102 (4)	0.0130 (4)	0.0149 (4)	-0.0006 (3)	-0.0001 (3)	0.0000 (3)
C10	0.0101 (4)	0.0142 (4)	0.0172 (5)	0.0002 (3)	-0.0010 (4)	-0.0003 (4)
C11	0.0128 (4)	0.0140 (4)	0.0142 (4)	0.0012 (3)	-0.0004 (4)	0.0000 (4)
C12	0.0127 (4)	0.0170 (5)	0.0170 (5)	0.0005 (4)	0.0017 (4)	0.0030 (4)
C13	0.0113 (4)	0.0181 (5)	0.0177 (5)	0.0007 (4)	0.0016 (4)	0.0028 (4)

C14	0.0151 (5)	0.0143 (4)	0.0147 (5)	0.0008 (3)	-0.0012 (4)	-0.0005 (4)
C15	0.0261 (6)	0.0183 (5)	0.0236 (6)	0.0020 (5)	-0.0020 (5)	0.0058 (5)
C16	0.0397 (8)	0.0215 (6)	0.0288 (7)	0.0028 (6)	0.0070 (6)	-0.0002 (5)
C17	0.0101 (4)	0.0204 (5)	0.0165 (5)	0.0025 (4)	-0.0005 (4)	0.0039 (4)
C18	0.0166 (5)	0.0265 (6)	0.0407 (8)	-0.0035 (5)	-0.0102 (5)	0.0018 (6)
C19	0.0214 (6)	0.0409 (8)	0.0248 (7)	0.0142 (6)	0.0012 (5)	-0.0047 (6)
C20	0.0176 (6)	0.0428 (9)	0.0310 (8)	-0.0059 (6)	-0.0073 (5)	0.0231 (7)
C21	0.0339 (7)	0.0139 (5)	0.0220 (6)	0.0002 (5)	0.0013 (5)	0.0024 (4)

Geometric parameters (Å, °)

O1—C3	1.3649 (15)	C11—C12	1.4102 (16)
O1—C21	1.4277 (17)	C11—C14	1.4833 (16)
O2—C14	1.2174 (15)	C12—C13	1.3860 (16)
O3—C14	1.3432 (16)	C12—H12A	0.9300
O3—C15	1.4509 (16)	C13—H13A	0.9300
N1—C8	1.3908 (15)	C15—C16	1.499 (2)
N1—C7	1.3958 (15)	C15—H15A	0.9700
N1—C17	1.5061 (15)	C15—H15B	0.9700
N2—C7	1.3176 (15)	C16—H16A	0.9600
N2—C9	1.3845 (15)	C16—H16B	0.9600
C1—C6	1.3912 (16)	C16—H16C	0.9600
C1—C2	1.3954 (17)	C17—C19	1.523 (2)
C1—H1A	0.9300	C17—C20	1.5236 (19)
C2—C3	1.3970 (18)	C17—C18	1.5330 (19)
C2—H2A	0.9300	C18—H18A	0.9600
C3—C4	1.3966 (17)	C18—H18B	0.9600
C4—C5	1.3868 (16)	C18—H18C	0.9600
C4—H4A	0.9300	C19—H19A	0.9600
C5—C6	1.4011 (17)	C19—H19B	0.9600
C5—H5A	0.9300	C19—H19C	0.9600
C6—C7	1.4792 (16)	C20—H20A	0.9600
C8—C13	1.4010 (16)	C20—H20B	0.9600
C8—C9	1.4117 (15)	C20—H20C	0.9600
C9—C10	1.3941 (16)	C21—H21A	0.9600
C10—C11	1.3921 (16)	C21—H21B	0.9600
C10—H10A	0.9300	C21—H21C	0.9600
C3—O1—C21	117.44 (11)	O2—C14—O3	124.05 (11)
C14—O3—C15	118.20 (11)	O2—C14—C11	124.57 (11)
C8—N1—C7	105.00 (9)	O3—C14—C11	111.38 (10)
C8—N1—C17	124.22 (9)	O3—C15—C16	109.43 (13)
C7—N1—C17	130.60 (10)	O3—C15—H15A	109.8
C7—N2—C9	104.95 (10)	C16—C15—H15A	109.8
C6—C1—C2	121.12 (11)	O3—C15—H15B	109.8
C6—C1—H1A	119.4	C16—C15—H15B	109.8
C2—C1—H1A	119.4	H15A—C15—H15B	108.2
C1—C2—C3	118.93 (11)	C15—C16—H16A	109.5
C1—C2—H2A	120.5	C15—C16—H16B	109.5
C3—C2—H2A	120.5	H16A—C16—H16B	109.5

supplementary materials

O1—C3—C4	115.29 (11)	C15—C16—H16C	109.5
O1—C3—C2	124.26 (11)	H16A—C16—H16C	109.5
C4—C3—C2	120.46 (11)	H16B—C16—H16C	109.5
C5—C4—C3	119.80 (11)	N1—C17—C19	108.05 (11)
C5—C4—H4A	120.1	N1—C17—C20	112.78 (10)
C3—C4—H4A	120.1	C19—C17—C20	110.03 (13)
C4—C5—C6	120.42 (11)	N1—C17—C18	109.01 (10)
C4—C5—H5A	119.8	C19—C17—C18	110.85 (13)
C6—C5—H5A	119.8	C20—C17—C18	106.14 (12)
C1—C6—C5	119.08 (11)	C17—C18—H18A	109.5
C1—C6—C7	120.58 (10)	C17—C18—H18B	109.5
C5—C6—C7	120.29 (10)	H18A—C18—H18B	109.5
N2—C7—N1	113.78 (10)	C17—C18—H18C	109.5
N2—C7—C6	120.71 (10)	H18A—C18—H18C	109.5
N1—C7—C6	125.36 (10)	H18B—C18—H18C	109.5
N1—C8—C13	133.18 (10)	C17—C19—H19A	109.5
N1—C8—C9	106.04 (10)	C17—C19—H19B	109.5
C13—C8—C9	120.74 (10)	H19A—C19—H19B	109.5
N2—C9—C10	128.47 (10)	C17—C19—H19C	109.5
N2—C9—C8	110.13 (10)	H19A—C19—H19C	109.5
C10—C9—C8	121.39 (10)	H19B—C19—H19C	109.5
C11—C10—C9	117.71 (10)	C17—C20—H20A	109.5
C11—C10—H10A	121.1	C17—C20—H20B	109.5
C9—C10—H10A	121.1	H20A—C20—H20B	109.5
C10—C11—C12	120.73 (10)	C17—C20—H20C	109.5
C10—C11—C14	118.74 (10)	H20A—C20—H20C	109.5
C12—C11—C14	120.53 (10)	H20B—C20—H20C	109.5
C13—C12—C11	121.94 (11)	O1—C21—H21A	109.5
C13—C12—H12A	119.0	O1—C21—H21B	109.5
C11—C12—H12A	119.0	H21A—C21—H21B	109.5
C12—C13—C8	117.39 (10)	O1—C21—H21C	109.5
C12—C13—H13A	121.3	H21A—C21—H21C	109.5
C8—C13—H13A	121.3	H21B—C21—H21C	109.5
C6—C1—C2—C3	-0.48 (19)	C7—N2—C9—C8	-0.18 (13)
C21—O1—C3—C4	176.78 (12)	N1—C8—C9—N2	2.09 (13)
C21—O1—C3—C2	-3.48 (19)	C13—C8—C9—N2	-175.77 (11)
C1—C2—C3—O1	177.04 (12)	N1—C8—C9—C10	-178.71 (11)
C1—C2—C3—C4	-3.22 (18)	C13—C8—C9—C10	3.43 (18)
O1—C3—C4—C5	-176.84 (11)	N2—C9—C10—C11	178.14 (12)
C2—C3—C4—C5	3.40 (18)	C8—C9—C10—C11	-0.90 (18)
C3—C4—C5—C6	0.13 (18)	C9—C10—C11—C12	-1.53 (18)
C2—C1—C6—C5	3.95 (18)	C9—C10—C11—C14	178.49 (11)
C2—C1—C6—C7	-178.63 (11)	C10—C11—C12—C13	1.53 (19)
C4—C5—C6—C1	-3.76 (17)	C14—C11—C12—C13	-178.48 (12)
C4—C5—C6—C7	178.81 (11)	C11—C12—C13—C8	0.94 (19)
C9—N2—C7—N1	-1.88 (14)	N1—C8—C13—C12	179.47 (12)
C9—N2—C7—C6	173.81 (11)	C9—C8—C13—C12	-3.35 (18)
C8—N1—C7—N2	3.19 (14)	C15—O3—C14—O2	-1.2 (2)
C17—N1—C7—N2	-172.05 (11)	C15—O3—C14—C11	179.23 (11)

C8—N1—C7—C6	-172.26 (11)	C10—C11—C14—O2	4.03 (19)
C17—N1—C7—C6	12.49 (19)	C12—C11—C14—O2	-175.95 (13)
C1—C6—C7—N2	-110.78 (14)	C10—C11—C14—O3	-176.36 (11)
C5—C6—C7—N2	66.61 (15)	C12—C11—C14—O3	3.65 (16)
C1—C6—C7—N1	64.39 (16)	C14—O3—C15—C16	-97.76 (15)
C5—C6—C7—N1	-118.22 (13)	C8—N1—C17—C19	76.40 (15)
C7—N1—C8—C13	174.46 (13)	C7—N1—C17—C19	-109.15 (15)
C17—N1—C8—C13	-9.9 (2)	C8—N1—C17—C20	-161.76 (13)
C7—N1—C8—C9	-3.02 (12)	C7—N1—C17—C20	12.69 (19)
C17—N1—C8—C9	172.61 (11)	C8—N1—C17—C18	-44.15 (16)
C7—N2—C9—C10	-179.31 (12)	C7—N1—C17—C18	130.30 (14)

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1, Cg2 and Cg3 are the centroids of the rings N1,N2,C7–C9 and C1–C6 and C8–C13, respectively.

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C5—H5A \cdots O2 ⁱ	0.93	2.57	3.3822 (16)	146.
C13—H13A \cdots O2 ⁱⁱ	0.93	2.52	3.4116 (16)	160.
C19—H19C \cdots N2 ⁱⁱⁱ	0.96	2.57	3.4976 (19)	164.
C2—H2A \cdots Cg3 ^{iv}	0.93	2.75	3.5594 (14)	146.
C18—H18C \cdots Cg2 ^v	0.96	2.74	3.6878 (16)	168.
C21—H21B \cdots Cg1 ^{iv}	0.96	2.78	3.4703 (16)	130.

Symmetry codes: (i) $-x+1/2, y+1/2, z-1/2$; (ii) $x-1/2, -y+1/2, z$; (iii) $x-1/2, -y+3/2, z$; (iv) $x, y+1, z$; (v) $-x-1/2, y+3/2, z+1/2$.

Fig. 1




Fig. 2

