

N-(3,5-Dimethylphenyl)-2-methylbenzamide

B. Thimme Gowda,^{a*} Miroslav Tokarčík,^b Jozef Kožíšek,^b
Vinola Zeena Rodrigues^a and Hartmut Fuess^c

^aDepartment of Chemistry, Mangalore University, Mangalagangothri 574 199, Mangalore, India, ^bFaculty of Chemical and Food Technology, Slovak Technical University, Radlinského 9, SK-812 37 Bratislava, Slovak Republic, and ^cInstitute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany

Correspondence e-mail: gowdabt@yahoo.com


Received 9 March 2010; accepted 10 March 2010

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(C-C) = 0.002$ Å; R factor = 0.037; wR factor = 0.120; data-to-parameter ratio = 15.0.

In the molecular structure of the title compound, $C_{16}H_{17}NO$, the amide group is twisted by 41.8 (2) and 29.0 (2)° out of the planes of the 2-methylphenyl and 3,5-dimethylphenyl rings, respectively. The two aromatic rings make a dihedral angle of 69.5 (1)°. In the crystal, intermolecular $N-H \cdots O$ hydrogen bonds connect the molecules into $C(4)$ chains running along the c axis.

Related literature

For our study of the effect of the substituents on the structures of benzanilides and for related structures, see: Gowda, *Foro et al.* (2008a,b); Gowda, Tokarčík *et al.* (2009). For synthesis, see: Gowda, *Foro et al.* (2008b).



Experimental

Crystal data

$C_{16}H_{17}NO$
 $M_r = 239.31$
Monoclinic, $P2_1/c$
 $a = 10.5174$ (5) Å

$b = 14.9616$ (7) Å
 $c = 8.9209$ (4) Å
 $\beta = 105.373$ (4)°
 $V = 1353.54$ (11) Å³

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.07$ mm⁻¹

$T = 295$ K
 $0.54 \times 0.08 \times 0.04$ mm

Data collection

Oxford Diffraction Xcalibur Ruby Gemini diffractometer
Absorption correction: multi-scan (*CrysAlis PRO*; Oxford Diffraction, 2009)
 $T_{min} = 0.957$, $T_{max} = 0.992$
14671 measured reflections
2548 independent reflections
1628 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.041$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.120$
 $S = 0.97$
2548 reflections

170 parameters
H-atom parameters constrained
 $\Delta\rho_{max} = 0.13$ e Å⁻³
 $\Delta\rho_{min} = -0.12$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$N1-H1N \cdots O1^i$	0.86	2.06	2.8935 (16)	163

Symmetry code: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2009); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2002); software used to prepare material for publication: *SHELXL97*, *PLATON* (Spek, 2009) and *WinGX* (Farrugia, 1999).

MT and JK thank the Grant Agency of the Slovak Republic (VEGA 1/0817/08) and the Structural Funds, Interreg IIIA, for financial support in purchasing the diffractometer. VZR thanks the University Grants Commission, Government of India, New Delhi, for the award of a research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2640).

References

- Brandenburg, K. (2002). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Gowda, B. T., Foro, S., Sowmya, B. P. & Fuess, H. (2008a). *Acta Cryst.* **E64**, o383.
Gowda, B. T., Foro, S., Sowmya, B. P. & Fuess, H. (2008b). *Acta Cryst.* **E64**, o1605.
Gowda, B. T., Tokarčík, M., Kožíšek, J., Rodrigues, V. Z. & Fuess, H. (2009). *Acta Cryst.* **E65**, o826.
Oxford Diffraction (2009). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2010). E66, o843 [doi:10.1107/S1600536810009116]

N-(3,5-Dimethylphenyl)-2-methylbenzamide

B. T. Gowda, M. Tokarčík, J. Kozísek, V. Z. Rodrigues and H. Fues

Comment

As a part of our efforts to explore the effect of the substituents on the structures of benzanilides (Gowda, Foro *et al.*, 2008a, b, Gowda, Tokarčík *et al.*, 2009), in the present work, the structure of 2-methyl-*N*-(3,5-dimethylphenyl)benzamide (I) has been determined. In the structure of (I) (Fig. 1), the N—H and C=O groups are in an antiperiplanar conformation. This conformation is similar to those observed, *e.g.* in 2-methyl-*N*-(phenyl)benzamide (II) (Gowda, Foro *et al.*, 2008a), 2-methyl-*N*-(2,6-dimethylphenyl)benzamide (III) (Gowda, Foro *et al.*, 2008b), and 2-methyl-*N*-(2,4-dimethylphenyl)benzamide (IV) (Gowda, Tokarčík *et al.*, 2009). Further, in (I) the conformation of the C=O group to the methyl substituent in the 2-methylphenyl ring is *syn*. This conformation is similar to those observed in (II) and (IV). The bond parameters in (I) are similar to those in (II), (III), (IV), and other benzanilides.

In the molecule, the amido group is twisted 41.8 (2)° and 29.0 (2)° out of the planes of the 2-methylphenyl and the 3,5-dimethylphenyl rings, respectively. The two aromatic rings make the dihedral angle of 69.5 (1)°. Intermolecular N—H···O hydrogen bonds (Table 1) connect the molecules into chains running along the *c*-axis (Fig. 2).

Experimental

Compound (I) as prepared according to the method described by Gowda, Foro *et al.* (2008b). Colourless blocks of (I) were obtained by slow evaporation from an ethanol solution (0.5 g in about 25 ml of ethanol) held at room temperature.

Refinement

All hydrogen atoms were positioned with idealized geometry using a riding model with C—H = 0.93–0.96 Å and N—H = 0.86 Å. The $U_{\text{iso}}(\text{H})$ values were set at $1.2U_{\text{eq}}(\text{C-aromatic, N})$ and $1.5U_{\text{eq}}(\text{C-methyl})$. The methyl groups with the carbon atoms C14, C15 and C16 exhibit orientational disorder in the positions of their H atoms, and each group was modelled by two sets of methyl hydrogen atoms. The refined occupancies are 0.82 (3) and 0.18 (3) for the C14-methyl group, 0.60 (3) and 0.40 (3) for the C15 group, 0.73 (2) and 0.27 (2) for the C16 group.

Figures




Fig. 1. Molecular structure of (I) showing the atom labelling scheme. Displacement ellipsoids are drawn at the 30% probability level and H atoms are represented as small spheres of arbitrary radii.




Fig. 2. Supramolecular chain formation in (I) with hydrogen bonds shown as dashed lines. Symmetry code (i): $x, -y+1/2, z+1/2$. H atoms not involved in hydrogen bonding were omitted for reasons of clarity.

N-(3,5-Dimethylphenyl)-2-methylbenzamide

Crystal data

$C_{16}H_{17}NO$	$F(000) = 512$
$M_r = 239.31$	$D_x = 1.174 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: $-P 2ybc$	Cell parameters from 5268 reflections
$a = 10.5174 (5) \text{ \AA}$	$\theta = 2.0\text{--}29.4^\circ$
$b = 14.9616 (7) \text{ \AA}$	$\mu = 0.07 \text{ mm}^{-1}$
$c = 8.9209 (4) \text{ \AA}$	$T = 295 \text{ K}$
$\beta = 105.373 (4)^\circ$	Needle, colourless
$V = 1353.54 (11) \text{ \AA}^3$	$0.54 \times 0.08 \times 0.04 \text{ mm}$
$Z = 4$	

Data collection

Oxford Diffraction Xcalibur Ruby Gemini diffractometer	2548 independent reflections
graphite	1628 reflections with $I > 2\sigma(I)$
Detector resolution: $10.434 \text{ pixels mm}^{-1}$	$R_{\text{int}} = 0.041$
ω scans	$\theta_{\text{max}} = 25.6^\circ, \theta_{\text{min}} = 2.0^\circ$
Absorption correction: multi-scan (CrysAlis Pro; Oxford Diffraction, 2009)	$h = -12 \rightarrow 12$
$T_{\text{min}} = 0.957, T_{\text{max}} = 0.992$	$k = -18 \rightarrow 18$
14671 measured reflections	$l = -10 \rightarrow 10$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.037$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.120$	H-atom parameters constrained
$S = 0.97$	$[\exp(2.50(\sin\theta/\lambda)^2)]/[\sigma^2(F_o^2) + (0.094P)^2]$, where $P = 0.33333F_o^2 + 0.66667F_c^2$
2548 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
170 parameters	$\Delta\rho_{\text{max}} = 0.13 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.12 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.20367 (15)	0.19330 (10)	0.08175 (15)	0.0348 (4)	
C2	0.09976 (16)	0.14838 (10)	0.14205 (17)	0.0392 (4)	
C3	-0.02376 (17)	0.12792 (12)	0.0426 (2)	0.0503 (5)	
C4	-0.1141 (2)	0.08515 (16)	0.1073 (3)	0.0724 (6)	
H4	-0.1967	0.0705	0.0435	0.087*	
C5	-0.0850 (2)	0.06389 (17)	0.2623 (3)	0.0835 (7)	
H5	-0.148	0.0359	0.3021	0.1*	
C6	0.0366 (2)	0.08381 (16)	0.3589 (3)	0.0760 (7)	
H6	0.0564	0.0695	0.464	0.091*	
C7	0.1290 (2)	0.12525 (12)	0.29829 (19)	0.0527 (5)	
H7	0.2121	0.1379	0.3629	0.063*	
C8	0.38098 (14)	0.30756 (10)	0.15513 (15)	0.0354 (4)	
C9	0.46745 (15)	0.27564 (11)	0.07407 (16)	0.0396 (4)	
H9	0.4536	0.2198	0.0266	0.048*	
C10	0.57512 (15)	0.32734 (12)	0.06388 (17)	0.0454 (4)	
C11	0.59283 (16)	0.41058 (12)	0.13406 (18)	0.0489 (4)	
H11	0.6637	0.4456	0.1256	0.059*	
C12	0.50798 (17)	0.44351 (11)	0.21670 (17)	0.0449 (4)	
C13	0.40205 (16)	0.39107 (11)	0.22612 (16)	0.0398 (4)	
H13	0.344	0.4119	0.2807	0.048*	
C14	-0.0633 (2)	0.15265 (17)	-0.1269 (2)	0.0743 (6)	
H14A	-0.1568	0.1457	-0.1673	0.112*	0.82 (3)
H14B	-0.0187	0.1144	-0.1829	0.112*	0.82 (3)
H14C	-0.0395	0.2137	-0.1386	0.112*	0.82 (3)
H14D	0.0135	0.1703	-0.1585	0.112*	0.18 (3)
H14E	-0.1248	0.2014	-0.143	0.112*	0.18 (3)
H14F	-0.1037	0.1021	-0.1872	0.112*	0.18 (3)
C15	0.67092 (19)	0.29122 (17)	-0.0193 (2)	0.0651 (6)	
H15A	0.7143	0.34	-0.0554	0.098*	0.60 (3)
H15B	0.6244	0.2558	-0.1063	0.098*	0.60 (3)
H15C	0.7353	0.2548	0.0507	0.098*	0.60 (3)
H15D	0.6684	0.2271	-0.0186	0.098*	0.40 (3)
H15E	0.7583	0.3112	0.0324	0.098*	0.40 (3)

supplementary materials

H15F	0.6474	0.3123	-0.1247	0.098*	0.40 (3)
C16	0.5318 (2)	0.53329 (13)	0.2949 (2)	0.0644 (5)	
H16A	0.5572	0.5754	0.227	0.097*	0.73 (2)
H16B	0.6009	0.5283	0.3896	0.097*	0.73 (2)
H16C	0.4525	0.5535	0.3182	0.097*	0.73 (2)
H16D	0.5166	0.5294	0.3962	0.097*	0.27 (2)
H16E	0.4728	0.5765	0.2336	0.097*	0.27 (2)
H16F	0.6213	0.5513	0.305	0.097*	0.27 (2)
N1	0.27245 (13)	0.25733 (9)	0.17550 (13)	0.0399 (3)	
H1N	0.2474	0.2691	0.2576	0.048*	
O1	0.22394 (11)	0.17203 (8)	-0.04280 (11)	0.0456 (3)	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0372 (8)	0.0346 (8)	0.0317 (7)	0.0014 (7)	0.0078 (6)	0.0008 (6)
C2	0.0428 (9)	0.0315 (8)	0.0444 (8)	-0.0029 (7)	0.0137 (7)	-0.0046 (6)
C3	0.0426 (10)	0.0445 (10)	0.0629 (11)	-0.0037 (8)	0.0122 (8)	-0.0086 (8)
C4	0.0485 (11)	0.0649 (14)	0.1019 (16)	-0.0159 (10)	0.0167 (11)	-0.0023 (12)
C5	0.0744 (16)	0.0734 (16)	0.1135 (19)	-0.0260 (13)	0.0438 (15)	0.0144 (13)
C6	0.0901 (17)	0.0724 (15)	0.0719 (12)	-0.0242 (13)	0.0327 (12)	0.0169 (11)
C7	0.0602 (11)	0.0481 (10)	0.0506 (10)	-0.0128 (8)	0.0163 (8)	0.0044 (8)
C8	0.0360 (8)	0.0395 (9)	0.0290 (7)	-0.0040 (7)	0.0056 (6)	0.0048 (6)
C9	0.0396 (9)	0.0428 (9)	0.0341 (7)	-0.0008 (7)	0.0055 (6)	0.0004 (6)
C10	0.0359 (9)	0.0604 (11)	0.0377 (8)	-0.0007 (8)	0.0058 (6)	0.0074 (7)
C11	0.0387 (9)	0.0580 (11)	0.0463 (8)	-0.0133 (8)	0.0050 (7)	0.0113 (8)
C12	0.0486 (10)	0.0413 (9)	0.0399 (8)	-0.0071 (8)	0.0031 (7)	0.0065 (7)
C13	0.0444 (9)	0.0393 (9)	0.0354 (7)	-0.0013 (7)	0.0101 (6)	0.0020 (6)
C14	0.0483 (11)	0.1004 (18)	0.0638 (12)	-0.0075 (12)	-0.0037 (9)	-0.0105 (11)
C15	0.0441 (10)	0.0897 (16)	0.0651 (11)	0.0000 (10)	0.0209 (9)	-0.0018 (10)
C16	0.0745 (14)	0.0471 (11)	0.0683 (11)	-0.0196 (10)	0.0134 (10)	-0.0037 (9)
N1	0.0461 (8)	0.0433 (8)	0.0341 (6)	-0.0087 (6)	0.0173 (5)	-0.0051 (5)
O1	0.0505 (7)	0.0517 (7)	0.0363 (6)	-0.0048 (6)	0.0148 (5)	-0.0068 (5)

Geometric parameters (\AA , $^\circ$)

C1—O1	1.2276 (17)	C12—C13	1.383 (2)
C1—N1	1.3490 (19)	C12—C16	1.504 (2)
C1—C2	1.499 (2)	C13—H13	0.93
C2—C7	1.389 (2)	C14—H14A	0.96
C2—C3	1.399 (2)	C14—H14B	0.96
C3—C4	1.391 (3)	C14—H14C	0.96
C3—C14	1.504 (3)	C14—H14D	0.96
C4—C5	1.372 (3)	C14—H14E	0.96
C4—H4	0.93	C14—H14F	0.96
C5—C6	1.372 (3)	C15—H15A	0.96
C5—H5	0.93	C15—H15B	0.96
C6—C7	1.378 (3)	C15—H15C	0.96
C6—H6	0.93	C15—H15D	0.96

C7—H7	0.93	C15—H15E	0.96
C8—C9	1.388 (2)	C15—H15F	0.96
C8—C13	1.392 (2)	C16—H16A	0.96
C8—N1	1.4180 (19)	C16—H16B	0.96
C9—C10	1.394 (2)	C16—H16C	0.96
C9—H9	0.93	C16—H16D	0.96
C10—C11	1.384 (2)	C16—H16E	0.96
C10—C15	1.501 (3)	C16—H16F	0.96
C11—C12	1.390 (2)	N1—H1N	0.86
C11—H11	0.93		
O1—C1—N1	123.50 (13)	C11—C12—C16	120.75 (16)
O1—C1—C2	121.78 (13)	C12—C13—C8	121.01 (15)
N1—C1—C2	114.72 (12)	C12—C13—H13	119.5
C7—C2—C3	120.21 (16)	C8—C13—H13	119.5
C7—C2—C1	118.90 (15)	C3—C14—H14A	109.5
C3—C2—C1	120.87 (14)	C3—C14—H14B	109.5
C4—C3—C2	117.28 (17)	C3—C14—H14C	109.5
C4—C3—C14	119.57 (18)	C3—C14—H14D	109.5
C2—C3—C14	123.12 (16)	C3—C14—H14E	109.5
C5—C4—C3	122.1 (2)	H14D—C14—H14E	109.5
C5—C4—H4	119	C3—C14—H14F	109.5
C3—C4—H4	119	H14D—C14—H14F	109.5
C4—C5—C6	120.27 (19)	H14E—C14—H14F	109.5
C4—C5—H5	119.9	C10—C15—H15A	109.5
C6—C5—H5	119.9	C10—C15—H15B	109.5
C5—C6—C7	119.2 (2)	C10—C15—H15C	109.5
C5—C6—H6	120.4	C10—C15—H15D	109.5
C7—C6—H6	120.4	C10—C15—H15E	109.5
C6—C7—C2	120.94 (19)	H15D—C15—H15E	109.5
C6—C7—H7	119.5	C10—C15—H15F	109.5
C2—C7—H7	119.5	H15D—C15—H15F	109.5
C9—C8—C13	119.91 (14)	H15E—C15—H15F	109.5
C9—C8—N1	123.04 (14)	C12—C16—H16A	109.5
C13—C8—N1	117.00 (13)	C12—C16—H16B	109.5
C8—C9—C10	119.92 (15)	C12—C16—H16C	109.5
C8—C9—H9	120	C12—C16—H16D	109.5
C10—C9—H9	120	C12—C16—H16E	109.5
C11—C10—C9	118.94 (15)	H16D—C16—H16E	109.5
C11—C10—C15	121.33 (16)	C12—C16—H16F	109.5
C9—C10—C15	119.71 (17)	H16D—C16—H16F	109.5
C10—C11—C12	122.04 (15)	H16E—C16—H16F	109.5
C10—C11—H11	119	C1—N1—C8	127.92 (12)
C12—C11—H11	119	C1—N1—H1N	116
C13—C12—C11	118.16 (15)	C8—N1—H1N	116
C13—C12—C16	121.09 (16)		
O1—C1—C2—C7	136.94 (16)	N1—C8—C9—C10	-177.47 (13)
N1—C1—C2—C7	-42.4 (2)	C8—C9—C10—C11	-0.7 (2)
O1—C1—C2—C3	-41.5 (2)	C8—C9—C10—C15	177.92 (15)

supplementary materials

N1—C1—C2—C3	139.08 (15)	C9—C10—C11—C12	1.2 (2)
C7—C2—C3—C4	0.7 (3)	C15—C10—C11—C12	-177.41 (16)
C1—C2—C3—C4	179.16 (17)	C10—C11—C12—C13	-0.9 (2)
C7—C2—C3—C14	178.64 (19)	C10—C11—C12—C16	178.48 (16)
C1—C2—C3—C14	-2.9 (3)	C11—C12—C13—C8	0.1 (2)
C2—C3—C4—C5	0.4 (3)	C16—C12—C13—C8	-179.26 (15)
C14—C3—C4—C5	-177.6 (2)	C9—C8—C13—C12	0.3 (2)
C3—C4—C5—C6	-0.7 (4)	N1—C8—C13—C12	177.91 (13)
C4—C5—C6—C7	-0.1 (4)	O1—C1—N1—C8	-3.4 (2)
C5—C6—C7—C2	1.2 (3)	C2—C1—N1—C8	176.00 (14)
C3—C2—C7—C6	-1.5 (3)	C9—C8—N1—C1	-27.9 (2)
C1—C2—C7—C6	-179.97 (18)	C13—C8—N1—C1	154.64 (15)
C13—C8—C9—C10	0.0 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1N \cdots O1 ⁱ	0.86	2.06	2.8935 (16)	163

Symmetry codes: (i) $x, -y+1/2, z+1/2$.

Fig. 1




Fig. 2

