

2-Amino-5-methylpyridinium 4-hydroxybenzoate

Madhukar Hemamalini and Hoong-Kun Fun*‡

X-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia

Correspondence e-mail: hkfun@usm.my


Received 4 March 2010; accepted 12 March 2010

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.001\text{ \AA}$; R factor = 0.039; wR factor = 0.137; data-to-parameter ratio = 24.3.

In the title salt, $\text{C}_6\text{H}_9\text{N}_2^+\cdot\text{C}_7\text{H}_5\text{O}_3^-$, the carboxylate mean plane of the 4-hydroxybenzoate anion is twisted by $13.07(4)^\circ$ from the attached ring. In the crystal structure, the ions are linked into a two-dimensional network by $\text{N}-\text{H}\cdots\text{O}$, $\text{O}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds. Within this network, the $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds generate $R_2^2(8)$ ring motifs. In addition, $\pi-\pi$ interactions involving the pyridinium rings, with a centroid-centroid distance of $3.7599(4)\text{ \AA}$, are observed.

Related literature

For background to the chemistry of substituted pyridines, see: Pozharski *et al.* (1997); Katritzky *et al.* (1996). For related structures, see: Hemamalini & Fun (2010a,b,c). For 4-hydroxybenzoic acid, see: Vishweshwar *et al.* (2003). For details of hydrogen bonding, see: Jeffrey & Saenger (1991); Jeffrey (1997); Scheiner (1997); Aakeröy *et al.* (2002). For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For bond-length data, see: Allen *et al.* (1987). For the stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986).



Experimental

Crystal data

$\text{C}_6\text{H}_9\text{N}_2^+\cdot\text{C}_7\text{H}_5\text{O}_3^-$
 $M_r = 246.26$
 Monoclinic, $P2_1/c$

$a = 12.9562(6)\text{ \AA}$
 $b = 8.7876(4)\text{ \AA}$
 $c = 11.3276(5)\text{ \AA}$

‡ Thomson Reuters ResearcherID: A-3561-2009.

$\beta = 108.397(1)^\circ$
 $V = 1223.78(10)\text{ \AA}^3$
 $Z = 4$
 Mo $K\alpha$ radiation

$\mu = 0.10\text{ mm}^{-1}$
 $T = 100\text{ K}$
 $0.39 \times 0.33 \times 0.27\text{ mm}$

Data collection

Bruker APEX DUO CCD area-detector diffractometer
 Absorption correction: multi-scan (*SADABS*; Bruker, 2009)
 $T_{\min} = 0.963$, $T_{\max} = 0.975$

20102 measured reflections
 5326 independent reflections
 4662 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.137$
 $S = 1.15$
 5326 reflections
 219 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.60\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.32\text{ e \AA}^{-3}$

Table 1
 Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1N1 \cdots O3	0.957 (17)	1.726 (17)	2.6738 (9)	170.4 (15)
N2—H1N2 \cdots O3 ⁱ	0.905 (14)	1.967 (14)	2.8443 (9)	162.9 (13)
N2—H2N2 \cdots O2	0.922 (14)	1.876 (14)	2.7962 (9)	176.1 (13)
O1—H1O1 \cdots O2 ⁱⁱ	0.892 (18)	1.779 (18)	2.6635 (8)	170.7 (19)
C3—H3A \cdots O2 ⁱⁱⁱ	1.016 (16)	2.476 (15)	3.1887 (9)	126.7 (10)

Symmetry codes: (i) $x, -y + \frac{3}{2}, z - \frac{1}{2}$; (ii) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

MH and HKF thank the Malaysian Government and Universiti Sains Malaysia for the Research University Golden Goose grant No. 1001/PFIZIK/811012. MH also thanks Universiti Sains Malaysia for a post-doctoral research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2424).

References

- Aakeröy, C. B., Beatty, A. M. & Helfrich, B. A. (2002). *J. Am. Chem. Soc.* **124**, 14425–14432.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2009). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
- Hemamalini, M. & Fun, H.-K. (2010a). *Acta Cryst. E66*, o623–o624.
- Hemamalini, M. & Fun, H.-K. (2010b). *Acta Cryst. E66*, o621.
- Hemamalini, M. & Fun, H.-K. (2010c). *Acta Cryst. E66*, o662.
- Jeffrey, G. A. (1997). *An Introduction to Hydrogen Bonding*. Oxford University Press.
- Jeffrey, G. A. & Saenger, W. (1991). *Hydrogen Bonding in Biological Structures*. Berlin: Springer.
- Katritzky, A. R., Rees, C. W. & Scriven, E. F. V. (1996). *Comprehensive Heterocyclic Chemistry II*. Oxford: Pergamon Press.

- Pozharski, A. F., Soldatenkov, A. T. & Katritzky, A. R. (1997). *Heterocycles in Life and Society*. New York: Wiley.
- Scheiner, S. (1997). *Hydrogen Bonding. A Theoretical Perspective*. Oxford University Press.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Vishweshwar, P., Nangia, A. & Lynch, V. M. (2003). *CrystEngComm*. **5**, 164–168.

supplementary materials

Acta Cryst. (2010). E66, o936-o937 [doi:10.1107/S1600536810009396]

2-Amino-5-methylpyridinium 4-hydroxybenzoate

M. Hemamalini and H.-K. Fun

Comment

In recent years, hydrogen bonds have attracted the interest of chemists and have been widely used to design and synthesize one, two and three-dimensional supramolecular compounds (Aakeröy *et al.*, 2002). Pyridine and its derivatives play an important role in heterocyclic chemistry (Pozharski *et al.*, 1997; Katritzky *et al.*, 1996). They are often involved in hydrogen-bond interactions (Jeffrey & Saenger, 1991; Jeffrey, 1997; Scheiner, 1997). 4-Hydroxybenzoic acid is a good hydrogen-bond donor and can form cocrystals with other organic molecules (Vishweshwar *et al.*, 2003). We have recently reported the crystal structures of 2-amino-5-methylpyridinium 3-aminobenzoate (Hemamalini & Fun, 2010a), 2-amino-5-methylpyridinium 4-nitrobenzoate (Hemamalini & Fun, 2010b) and 2-amino-5-methylpyridinium nicotinate (Hemamalini & Fun, 2010c). In continuation of our studies of pyrimidinium derivatives, the crystal structure determination of the title compound has been undertaken.

The asymmetric unit (Fig. 1) contains a 2-amino-5-methylpyridinium cation and a 4-hydroxybenzoate anion. In the 2-amino-5-methylpyridinium cation, a wide angle ($122.65(6)^\circ$) is subtended at the protonated N1 atom. The 2-amino-5-methylpyridinium cation is planar, with a maximum deviation of $0.024(1)\text{\AA}$ for atom N1. The bond lengths are normal (Allen *et al.*, 1987). In the 4-hydroxybenzoate anion, the carboxylate group is twisted slightly from the attached ring; the dihedral angle between C7–C12 and O2/O3/C12–C13 planes is $13.07(4)^\circ$.

In the crystal packing (Fig. 2), the protonated N1 atom and the 2-amino group N2 atom are hydrogen-bonded to the carboxylate oxygen atoms (O2 and O3) via a pair of N—H \cdots O hydrogen bonds forming a ring motif $R_2^2(8)$ (Bernstein *et al.*, 1995). The hydroxyl group hydrogen atom is also hydrogen-bonded to the carboxylate oxygen atom through O1—H1O1 \cdots O2 hydrogen bonds. The packing is further stabilized by weak C—H \cdots O and π – π interactions involving the pyridinium (centroid Cg1) rings, with Cg1–Cg1 = $3.7599(4)\text{\AA}$ [symmetry codes: 1-x, 1-y, 1-z].

Experimental

A hot methanol solution (20 ml) of 2-amino-5-methylpyridine (54 mg, Aldrich) and 4-hydroxybenzoic acid (69 mg, Merck) were mixed and warmed over a heating magnetic stirrer for a few minutes. The resulting solution was allowed to cool slowly at room temperature and crystals of the title compound appeared after a few days.

Refinement

All the H atoms were located in a difference Fourier map and allowed to refine freely [N—H = $0.904(14)$ - $0.956(16)\text{\AA}$, C—H = $0.952(16)$ - $1.020(15)\text{\AA}$ and O—H = $0.893(18)\text{\AA}$].

supplementary materials

Figures




Fig. 1. The asymmetric unit of the title compound. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. The crystal packing of the title compound, showing hydrogen-bonded (dashed lines) networks. H atoms are not involving the hydrogen bond interactions are omitted for clarity.

2-Amino-5-methylpyridinium 4-hydroxybenzoate

Crystal data

$C_6H_9N_2^+ \cdot C_7H_5O_3^-$	$F(000) = 520$
$M_r = 246.26$	$D_x = 1.337 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 9907 reflections
$a = 12.9562 (6) \text{ \AA}$	$\theta = 6.5\text{--}36.2^\circ$
$b = 8.7876 (4) \text{ \AA}$	$\mu = 0.10 \text{ mm}^{-1}$
$c = 11.3276 (5) \text{ \AA}$	$T = 100 \text{ K}$
$\beta = 108.397 (1)^\circ$	Blcok, colourless
$V = 1223.78 (10) \text{ \AA}^3$	$0.39 \times 0.33 \times 0.27 \text{ mm}$
$Z = 4$	

Data collection

Bruker APEX DUO CCD area-detector diffractometer	5326 independent reflections
Radiation source: fine-focus sealed tube graphite	4662 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.019$
Absorption correction: multi-scan (SADABS; Bruker, 2009)	$\theta_{\text{max}} = 35.0^\circ, \theta_{\text{min}} = 6.2^\circ$
$T_{\text{min}} = 0.963, T_{\text{max}} = 0.975$	$h = -19 \rightarrow 20$
20102 measured reflections	$k = -14 \rightarrow 11$
	$l = -18 \rightarrow 15$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.039$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.137$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.15$	$w = 1/[\sigma^2(F_o^2) + (0.0859P)^2 + 0.1288P]$ where $P = (F_o^2 + 2F_c^2)/3$
5326 reflections	$(\Delta/\sigma)_{\max} = 0.001$
219 parameters	$\Delta\rho_{\max} = 0.60 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.32 \text{ e \AA}^{-3}$

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	-0.07705 (4)	1.29454 (6)	0.44060 (5)	0.01923 (12)
O2	0.23879 (4)	0.85691 (6)	0.26656 (5)	0.01752 (11)
O3	0.34273 (4)	0.90704 (7)	0.45979 (5)	0.02001 (12)
C7	0.17088 (6)	1.07237 (8)	0.50635 (6)	0.01751 (13)
C8	0.08861 (6)	1.16244 (9)	0.52300 (6)	0.01815 (13)
C9	0.00128 (5)	1.20628 (7)	0.42014 (6)	0.01493 (12)
C10	-0.00280 (5)	1.15864 (8)	0.30079 (6)	0.01519 (12)
C11	0.07897 (5)	1.06554 (8)	0.28576 (6)	0.01482 (12)
C12	0.16699 (5)	1.02150 (7)	0.38786 (6)	0.01410 (11)
C13	0.25438 (5)	0.92210 (8)	0.37051 (6)	0.01435 (12)
N1	0.48836 (5)	0.71343 (7)	0.42194 (5)	0.01544 (11)
N2	0.39201 (5)	0.66496 (8)	0.21647 (6)	0.01851 (12)
C1	0.48185 (5)	0.64504 (7)	0.31296 (6)	0.01461 (12)
C2	0.57195 (6)	0.55617 (8)	0.30846 (6)	0.01732 (13)
C3	0.65954 (6)	0.53911 (8)	0.41374 (7)	0.01848 (13)
C4	0.66298 (5)	0.60787 (8)	0.52806 (6)	0.01614 (12)
C5	0.57535 (5)	0.69542 (8)	0.52670 (6)	0.01641 (12)
C6	0.75785 (6)	0.58625 (10)	0.64330 (7)	0.02191 (14)
H2A	0.5718 (11)	0.5103 (16)	0.2257 (12)	0.027 (3)*
H3A	0.7256 (12)	0.4762 (18)	0.4148 (13)	0.034 (3)*
H5A	0.5712 (11)	0.7533 (16)	0.5988 (13)	0.027 (3)*

supplementary materials

H6A	0.7657 (12)	0.4804 (19)	0.6662 (14)	0.039 (4)*
H6B	0.8244 (13)	0.6124 (18)	0.6294 (14)	0.038 (4)*
H6C	0.7483 (13)	0.6507 (19)	0.7124 (16)	0.040 (4)*
H7A	0.2334 (11)	1.0425 (17)	0.5794 (13)	0.029 (3)*
H8A	0.0899 (13)	1.2010 (18)	0.6068 (15)	0.038 (4)*
H10A	-0.0628 (12)	1.1968 (16)	0.2289 (14)	0.030 (3)*
H11A	0.0758 (10)	1.0333 (15)	0.2012 (11)	0.021 (3)*
H101	-0.1257 (15)	1.3172 (19)	0.3673 (16)	0.045 (4)*
H1N1	0.4304 (13)	0.7749 (19)	0.4310 (14)	0.038 (4)*
H1N2	0.3819 (12)	0.6231 (15)	0.1407 (13)	0.028 (3)*
H2N2	0.3394 (11)	0.7245 (15)	0.2326 (12)	0.025 (3)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0165 (2)	0.0239 (3)	0.0175 (2)	0.00637 (17)	0.00561 (18)	0.00151 (18)
O2	0.0140 (2)	0.0240 (2)	0.0143 (2)	0.00056 (16)	0.00405 (16)	-0.00257 (17)
O3	0.0149 (2)	0.0292 (3)	0.0137 (2)	0.00667 (18)	0.00134 (16)	-0.00034 (18)
C7	0.0160 (3)	0.0233 (3)	0.0122 (2)	0.0047 (2)	0.0029 (2)	0.0015 (2)
C8	0.0176 (3)	0.0240 (3)	0.0124 (2)	0.0053 (2)	0.0041 (2)	0.0011 (2)
C9	0.0131 (2)	0.0166 (3)	0.0150 (2)	0.00114 (18)	0.00441 (19)	0.00139 (19)
C10	0.0128 (2)	0.0175 (3)	0.0136 (2)	0.00088 (19)	0.00182 (19)	0.00068 (19)
C11	0.0137 (2)	0.0171 (3)	0.0126 (2)	0.00074 (19)	0.00256 (19)	0.00033 (19)
C12	0.0123 (2)	0.0169 (2)	0.0127 (2)	0.00125 (18)	0.00344 (18)	0.00125 (19)
C13	0.0125 (2)	0.0180 (3)	0.0127 (2)	0.00072 (19)	0.00415 (19)	0.00162 (19)
N1	0.0147 (2)	0.0176 (2)	0.0136 (2)	0.00132 (17)	0.00391 (18)	-0.00160 (17)
N2	0.0164 (2)	0.0227 (3)	0.0144 (2)	0.00148 (19)	0.00197 (19)	-0.00235 (19)
C1	0.0152 (2)	0.0154 (2)	0.0135 (2)	-0.00101 (18)	0.0050 (2)	-0.00051 (19)
C2	0.0187 (3)	0.0187 (3)	0.0155 (3)	0.0024 (2)	0.0067 (2)	-0.0014 (2)
C3	0.0174 (3)	0.0202 (3)	0.0185 (3)	0.0030 (2)	0.0066 (2)	-0.0001 (2)
C4	0.0144 (2)	0.0177 (3)	0.0158 (3)	-0.0001 (2)	0.0041 (2)	0.0008 (2)
C5	0.0160 (3)	0.0193 (3)	0.0134 (2)	0.0003 (2)	0.0039 (2)	-0.0015 (2)
C6	0.0163 (3)	0.0271 (3)	0.0193 (3)	0.0001 (2)	0.0013 (2)	0.0036 (2)

Geometric parameters (\AA , $^\circ$)

O1—C9	1.3541 (8)	N1—C5	1.3636 (9)
O1—H101	0.893 (18)	N1—H1N1	0.956 (16)
O2—C13	1.2670 (8)	N2—C1	1.3331 (9)
O3—C13	1.2728 (8)	N2—H1N2	0.904 (14)
C7—C8	1.3875 (9)	N2—H2N2	0.922 (14)
C7—C12	1.4004 (9)	C1—C2	1.4187 (9)
C7—H7A	0.993 (14)	C2—C3	1.3705 (10)
C8—C9	1.3978 (9)	C2—H2A	1.020 (13)
C8—H8A	1.003 (15)	C3—C4	1.4169 (10)
C9—C10	1.4003 (9)	C3—H3A	1.016 (15)
C10—C11	1.3909 (9)	C4—C5	1.3675 (9)
C10—H10A	0.990 (15)	C4—C6	1.4962 (10)
C11—C12	1.3984 (9)	C5—H5A	0.978 (14)

C11—H11A	0.987 (12)	C6—H6A	0.963 (16)
C12—C13	1.4912 (9)	C6—H6B	0.952 (16)
N1—C1	1.3516 (8)	C6—H6C	1.006 (17)
C9—O1—H101	108.4 (11)	C1—N2—H1N2	123.5 (9)
C8—C7—C12	121.04 (6)	C1—N2—H2N2	114.9 (8)
C8—C7—H7A	119.8 (8)	H1N2—N2—H2N2	121.6 (12)
C12—C7—H7A	119.1 (8)	N2—C1—N1	118.50 (6)
C7—C8—C9	119.89 (6)	N2—C1—C2	123.94 (6)
C7—C8—H8A	122.5 (9)	N1—C1—C2	117.56 (6)
C9—C8—H8A	117.6 (9)	C3—C2—C1	119.59 (6)
O1—C9—C8	117.93 (6)	C3—C2—H2A	121.2 (8)
O1—C9—C10	122.31 (6)	C1—C2—H2A	119.2 (8)
C8—C9—C10	119.76 (6)	C2—C3—C4	121.81 (6)
C11—C10—C9	119.73 (6)	C2—C3—H3A	122.2 (8)
C11—C10—H10A	122.0 (9)	C4—C3—H3A	116.0 (8)
C9—C10—H10A	118.2 (9)	C5—C4—C3	116.31 (6)
C10—C11—C12	121.02 (6)	C5—C4—C6	122.13 (6)
C10—C11—H11A	119.2 (7)	C3—C4—C6	121.56 (6)
C12—C11—H11A	119.7 (7)	N1—C5—C4	122.03 (6)
C11—C12—C7	118.53 (6)	N1—C5—H5A	114.7 (8)
C11—C12—C13	120.53 (6)	C4—C5—H5A	123.2 (8)
C7—C12—C13	120.94 (6)	C4—C6—H6A	110.2 (9)
O2—C13—O3	122.08 (6)	C4—C6—H6B	111.2 (9)
O2—C13—C12	118.84 (6)	H6A—C6—H6B	104.7 (13)
O3—C13—C12	119.08 (6)	C4—C6—H6C	109.8 (9)
C1—N1—C5	122.65 (6)	H6A—C6—H6C	111.4 (13)
C1—N1—H1N1	121.6 (9)	H6B—C6—H6C	109.5 (13)
C5—N1—H1N1	115.7 (9)		
C12—C7—C8—C9	-1.35 (11)	C11—C12—C13—O3	166.93 (6)
C7—C8—C9—O1	-179.81 (6)	C7—C12—C13—O3	-13.29 (10)
C7—C8—C9—C10	0.17 (11)	C5—N1—C1—N2	177.71 (6)
O1—C9—C10—C11	-178.73 (6)	C5—N1—C1—C2	-2.43 (10)
C8—C9—C10—C11	1.29 (10)	N2—C1—C2—C3	-178.20 (7)
C9—C10—C11—C12	-1.60 (10)	N1—C1—C2—C3	1.94 (10)
C10—C11—C12—C7	0.44 (10)	C1—C2—C3—C4	0.08 (11)
C10—C11—C12—C13	-179.77 (6)	C2—C3—C4—C5	-1.66 (11)
C8—C7—C12—C11	1.05 (11)	C2—C3—C4—C6	178.74 (7)
C8—C7—C12—C13	-178.74 (6)	C1—N1—C5—C4	0.83 (11)
C11—C12—C13—O2	-12.12 (10)	C3—C4—C5—N1	1.25 (10)
C7—C12—C13—O2	167.67 (6)	C6—C4—C5—N1	-179.16 (6)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1N1···O3	0.957 (17)	1.726 (17)	2.6738 (9)	170.4 (15)
N2—H1N2···O3 ⁱ	0.905 (14)	1.967 (14)	2.8443 (9)	162.9 (13)
N2—H2N2···O2	0.922 (14)	1.876 (14)	2.7962 (9)	176.1 (13)
O1—H101···O2 ⁱⁱ	0.892 (18)	1.779 (18)	2.6635 (8)	170.7 (19)

supplementary materials

C3—H3A···O2ⁱⁱⁱ

1.016 (16)

2.476 (15)

3.1887 (9)

126.7 (10)

Symmetry codes: (i) $x, -y+3/2, z-1/2$; (ii) $-x, y+1/2, -z+1/2$; (iii) $-x+1, y-1/2, -z+1/2$.

Fig. 1




Fig. 2

