

6,8-Diiodo-5,7-dimethoxy-4-methylcoumarin

P. S. Pereira Silva,^a Mehtab Parveen,^b Zakia Khanam,^b Akhtar Ali^b and M. Ramos Silva^{a*}

^aCEMDRX, Physics Department, University of Coimbra, P-3004-516 Coimbra, Portugal, and ^bDepartment of Chemistry, Aligarh Muslim University, Aligarh 202 002, India

Correspondence e-mail: psidonio@pollux.fis.uc.pt


Received 24 March 2010; accepted 25 March 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.023; wR factor = 0.055; data-to-parameter ratio = 23.7.

In the title compound, $\text{C}_{12}\text{H}_{10}\text{I}_2\text{O}_4$, the methoxy groups are twisted considerably with respect to the plane of the aromatic ring [$\text{CH}_3-\text{O}-\text{C}-\text{C}$ torsion angles = $-85.9(3)$ and $-92.8(3)^\circ$]. In the crystal, molecules are linked by weak $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds and $\text{O}\cdots\text{I}$ contacts [3.194(2) \AA].

Related literature

For the medicinal applications of coumarin derivatives, see: Lin *et al.* (2006); Massimo *et al.* (2003); Tyagi *et al.* (2003); Nawrot-Modranka *et al.* (2006); Sardari *et al.* (1999); Huang *et al.* (2005); Elinos-Baez *et al.* (2005). For the synthesis of the title compound, see: Ali & Ilyas (1986).



Experimental

Crystal data

$\text{C}_{12}\text{H}_{10}\text{I}_2\text{O}_4$
 $M_r = 472.00$
Monoclinic, $P2_1/c$
 $a = 10.8681(2)\text{ \AA}$
 $b = 9.1179(2)\text{ \AA}$

$c = 17.2315(3)\text{ \AA}$
 $\beta = 125.395(1)^\circ$
 $V = 1391.95(5)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 4.52\text{ mm}^{-1}$
 $T = 293\text{ K}$

$0.30 \times 0.24 \times 0.16\text{ mm}$

Data collection

Bruker APEXII CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 2003)
 $T_{\min} = 0.396$, $T_{\max} = 0.485$

27048 measured reflections
3928 independent reflections
3453 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.023$
 $wR(F^2) = 0.055$
 $S = 1.06$
3928 reflections

166 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.71\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.96\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C3—H3—O2 ⁱ	0.93	2.53	3.460 (3)	175

Symmetry code: (i) $-x, -y, -z$.

Data collection: *APEX2* (Bruker, 2003); cell refinement: *SAINT* (Bruker, 2003); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

This work was supported by the Fundação para a Ciência e a Tecnologia (FCT) under scholarship SFRH/BD/38387/2008.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5225).

References

- Ali, S. M. & Ilyas, M. (1986). *J. Org. Chem.* **51**, 5415–5417.
Bruker (2003). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Elinos-Baez, C. M., Leon, F. & Santos, E. (2005). *Cell Biol. Int.* **29**, 703–708.
Huang, L., Yuon, X., Yu, D., Lee, K. H. & Chin, H. C. (2005). *Virology*, **332**, 623–628.
Lin, C. M., Huang, S. T., Lee, F. W., Sawkuo, H. & Lin, M. H. (2006). *Bioorg. Med. Chem.* **14**, 4402–4409.
Massimo, C., Francesco, E., Federica, M., Carla, M. M., Prieto, G. S. & Carlos, R. J. (2003). *Aust. J. Chem.* **56**, 59–60.
Nawrot-Modranka, J., Nawrot, E. & Graczyk, J. (2006). *Eur. J. Med. Chem.* **41**, 1301–1309.
Sardari, S., Mori, Y., Horita, K., Micetich, R. G., Nishibe, S. & Daneshtalab, M. (1999). *Bioorg. Med. Chem.* **7**, 1933–1940.
Sheldrick, G. M. (2003). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
Tyagi, A. K., Raj, H. G., Vohra, P., Gupta, G., Kumari, R., Kumar, P. & Gupta, R. K. (2003). *Eur. J. Med. Chem.* **40**, 413–420.

supplementary materials

Acta Cryst. (2010). E66, o988 [doi:10.1107/S1600536810011360]

6,8-Diido-5,7-dimethoxy-4-methylcoumarin

P. S. Pereira Silva, M. Parveen, Z. Khanam, A. Ali and M. Ramos Silva

Comment

Coumarin is the simplest member of the group of oxygen heterocyclics called benzo-2-pyrone. Coumarins are an important class of compounds due to their presence in natural products as well as their medicinal applications such as anti-inflammatory (Lin *et al.*, 2006), anti-viral (Massimo *et al.*, 2003), antioxidant (Tyagi *et al.*, 2003), antibacterial (Nawrot-Modranka *et al.*, 2006), antifungal (Sardari *et al.*, 1999), anti-HIV (Huang *et al.*, 2005) and as anti-carcinogenic (Elinos-Baez *et al.*, 2005). Besides the wide spectrum of biological applications of coumarin and its derivatives, the chemical literature also embodies their applications as cosmetics, optical brightening agents, and laser dyes. A recent report has revealed the anion sensing ability of some coumarin derivatives. Among various coumarin derivatives, recent pharmacological evaluation of iodocoumarins as cannabinoid receptor antagonists and inverse agonists has been done. Iodocoumarins such as 8-iodo-7-hydroxycoumarin exhibited moderate activity and 8-iodo-5,7-dihydroxycoumarin displayed good antimicrobial properties with MIC values <100 µg/ml. Also, iodocoumarins had been successfully used for the optimization of reaction conditions and kinetic studies in high throughput format. Because of the biological and pharmaceutical importance of iodocoumarins, several protocols for the synthesis have been reported.

In the light of the mentioned above we planned to synthesize iodocoumarins by reaction of 5,7-dimethoxy-4-methylcoumarin with iodine in basic media (Ali & Ilyas, 1986).

In the molecule of the title compound (Fig. 1), the best plane through the aromatic ring shows an r.m.s. deviation of 0.0154 Å; the O1—C2—C3—C4—C10—C9 ring shows a slightly larger deviation from planarity, with an r.m.s. deviation of 0.0189 Å. The angle between these two planes is 4.96 (11)°.

The C2 atom of the carbonyl group has a distorted trigonal geometry with O2—C2—O1 [116.6 (2)°] and O2—C2—C3 [126.5 (2)°] deviating significantly from the ideal sp^2 value of 120°.

The methoxy groups are considerably twisted with respect to the plane of the aromatic ring as indicated by the torsion angles C12—O3—C5—C6 [-85.9 (3)°] and C13—O4—C7—C6 [90.7 (3)°]. The iodine atoms are almost in the plane of the benzene ring and the methyl group is slightly out of the pyrone ring plane.

The molecules are linked across an inversion centre by one weak hydrogen bond of the C—H···O type (Fig. 2, Table 2).

Experimental

To a stirred solution of 5,7-dimethoxy-4-methylcoumarin (2.20 g, 10 mmol) in 15–20 ml of methanol containing 8.2 g KOH was dropwise added to a solution of I₂ (2.56 g, 10 mmol) over a period of 30 min and stirred at room temperature for about 2 hours. The reaction mixture was poured into water and residual iodine was removed by washing with sodium thiosulphate. On treatment with sodium thiosulphate we obtained a precipitate which was filtered and crystallized with CHCl₃—MeOH as white crystals (300 mg, m.p. 490 K).

supplementary materials

Refinement

All H atoms were located in a difference Fourier synthesis, placed in calculated positions and refined as riding on their parent atoms, using SHELXL97 (Sheldrick, 2008) defaults.

Figures




Fig. 1. A plot of the title compound. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. A packing diagram for the title compound, viewed down the *a* axis, with the hydrogen bonds depicted by dashed lines.

6,8-diiodo-5,7-dimethoxy-4-methyl-2*H*-chromen-2-one

Crystal data

C ₁₂ H ₁₀ I ₂ O ₄	<i>F</i> (000) = 880
<i>M_r</i> = 472.00	<i>D_x</i> = 2.252 Mg m ⁻³
Monoclinic, <i>P</i> 2 ₁ / <i>c</i>	Melting point: 490 K
Hall symbol: -P 2ybc	Mo <i>K</i> α radiation, λ = 0.71073 Å
<i>a</i> = 10.8681 (2) Å	Cell parameters from 7934 reflections
<i>b</i> = 9.1179 (2) Å	θ = 2.4–29.5°
<i>c</i> = 17.2315 (3) Å	μ = 4.52 mm ⁻¹
β = 125.395 (1)°	<i>T</i> = 293 K
<i>V</i> = 1391.95 (5) Å ³	Irregular block, pale yellow
<i>Z</i> = 4	0.30 × 0.24 × 0.16 mm

Data collection

Bruker APEXII CCD area-detector diffractometer	3928 independent reflections
Radiation source: fine-focus sealed tube graphite	3453 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.022$

Absorption correction: multi-scan
(SADABS; Sheldrick, 2003)
 $T_{\min} = 0.396$, $T_{\max} = 0.485$
27048 measured reflections

$h = -14 \rightarrow 15$
 $k = -12 \rightarrow 12$
 $l = -24 \rightarrow 23$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.023$
 $wR(F^2) = 0.055$
 $S = 1.06$
3928 reflections
166 parameters
0 restraints

Primary atom site location: structure-invariant direct methods
Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/\sigma^2(F_o^2) + (0.0226P)^2 + 1.1738P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.71 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.96 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
I1	0.51053 (2)	0.82363 (2)	0.156895 (15)	0.05749 (7)
I2	0.11277 (2)	0.593603 (19)	0.266073 (13)	0.04546 (6)
O1	0.06827 (19)	0.35475 (17)	0.12276 (12)	0.0353 (3)
O2	-0.0507 (2)	0.1432 (2)	0.07590 (17)	0.0548 (5)
O3	0.39384 (18)	0.53122 (19)	0.03873 (12)	0.0383 (4)
O4	0.3379 (2)	0.79647 (18)	0.25451 (12)	0.0434 (4)
C8	0.1966 (3)	0.5766 (2)	0.18397 (16)	0.0317 (4)
C7	0.2959 (3)	0.6823 (2)	0.19268 (16)	0.0337 (5)
C6	0.3586 (3)	0.6668 (2)	0.14197 (17)	0.0352 (5)
C5	0.3273 (2)	0.5453 (3)	0.08510 (15)	0.0318 (4)
C10	0.2315 (2)	0.4324 (2)	0.07795 (15)	0.0297 (4)
C9	0.1647 (2)	0.4549 (2)	0.12598 (15)	0.0294 (4)
C4	0.2056 (3)	0.2920 (2)	0.03026 (16)	0.0352 (5)
C3	0.1095 (3)	0.1972 (3)	0.02911 (18)	0.0395 (5)

supplementary materials

H3	0.0905	0.1087	-0.0028	0.047*
C2	0.0351 (3)	0.2248 (3)	0.07400 (18)	0.0379 (5)
C13	0.2462 (4)	0.9256 (3)	0.2126 (2)	0.0621 (8)
H13A	0.2393	0.9513	0.1562	0.093*
H13B	0.2911	1.0051	0.2573	0.093*
H13C	0.1470	0.9066	0.1964	0.093*
C12	0.3139 (3)	0.6047 (3)	-0.05204 (19)	0.0497 (6)
H12A	0.2153	0.5623	-0.0936	0.075*
H12B	0.3687	0.5937	-0.0797	0.075*
H12C	0.3044	0.7070	-0.0433	0.075*
C11	0.2873 (4)	0.2436 (3)	-0.0117 (2)	0.0582 (8)
H11A	0.2607	0.1440	-0.0331	0.087*
H11B	0.3942	0.2502	0.0357	0.087*
H11C	0.2592	0.3057	-0.0646	0.087*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
I1	0.05881 (12)	0.05243 (11)	0.06135 (13)	-0.02587 (9)	0.03486 (10)	-0.01030 (8)
I2	0.05746 (11)	0.04523 (10)	0.04782 (10)	-0.00537 (7)	0.03859 (9)	-0.01056 (7)
O1	0.0400 (9)	0.0303 (7)	0.0438 (9)	-0.0055 (6)	0.0290 (8)	-0.0069 (7)
O2	0.0650 (13)	0.0391 (9)	0.0823 (15)	-0.0165 (9)	0.0552 (12)	-0.0145 (10)
O3	0.0370 (8)	0.0458 (9)	0.0384 (9)	0.0023 (7)	0.0255 (8)	0.0035 (7)
O4	0.0572 (11)	0.0325 (8)	0.0355 (9)	-0.0062 (8)	0.0239 (8)	-0.0092 (7)
C8	0.0355 (11)	0.0304 (10)	0.0307 (10)	0.0031 (8)	0.0200 (9)	-0.0015 (8)
C7	0.0380 (12)	0.0277 (10)	0.0286 (10)	-0.0006 (8)	0.0154 (9)	-0.0023 (8)
C6	0.0355 (11)	0.0325 (11)	0.0330 (11)	-0.0057 (9)	0.0172 (10)	0.0007 (9)
C5	0.0299 (10)	0.0357 (11)	0.0291 (10)	0.0019 (8)	0.0167 (9)	0.0032 (8)
C10	0.0307 (10)	0.0294 (10)	0.0271 (10)	0.0008 (8)	0.0157 (9)	-0.0008 (8)
C9	0.0293 (10)	0.0280 (9)	0.0296 (10)	0.0003 (8)	0.0162 (9)	-0.0005 (8)
C4	0.0400 (12)	0.0334 (11)	0.0338 (11)	0.0017 (9)	0.0224 (10)	-0.0036 (9)
C3	0.0477 (14)	0.0299 (11)	0.0453 (13)	-0.0041 (10)	0.0294 (12)	-0.0077 (9)
C2	0.0409 (12)	0.0290 (10)	0.0453 (13)	-0.0031 (9)	0.0257 (11)	-0.0049 (9)
C13	0.093 (2)	0.0340 (13)	0.0625 (19)	0.0072 (15)	0.0471 (19)	-0.0025 (12)
C12	0.0480 (15)	0.0696 (18)	0.0379 (13)	0.0039 (13)	0.0285 (12)	0.0077 (12)
C11	0.078 (2)	0.0493 (16)	0.077 (2)	-0.0089 (15)	0.0618 (19)	-0.0206 (15)

Geometric parameters (\AA , $^\circ$)

I1—C6	2.084 (2)	C10—C4	1.458 (3)
I2—C8	2.085 (2)	C4—C3	1.347 (3)
O1—C9	1.367 (3)	C4—C11	1.500 (3)
O1—C2	1.375 (3)	C3—C2	1.428 (3)
O2—C2	1.208 (3)	C3—H3	0.9300
O3—C5	1.359 (3)	C13—H13A	0.9600
O3—C12	1.441 (3)	C13—H13B	0.9600
O4—C7	1.365 (3)	C13—H13C	0.9600
O4—C13	1.438 (3)	C12—H12A	0.9600
C8—C7	1.389 (3)	C12—H12B	0.9600

C8—C9	1.396 (3)	C12—H12C	0.9600
C7—C6	1.392 (3)	C11—H11A	0.9600
C6—C5	1.385 (3)	C11—H11B	0.9600
C5—C10	1.418 (3)	C11—H11C	0.9600
C10—C9	1.397 (3)		
C9—O1—C2	121.60 (18)	C4—C3—C2	123.8 (2)
C5—O3—C12	113.74 (18)	C4—C3—H3	118.1
C7—O4—C13	114.3 (2)	C2—C3—H3	118.1
C7—C8—C9	118.9 (2)	O2—C2—O1	116.6 (2)
C7—C8—I2	119.53 (16)	O2—C2—C3	126.5 (2)
C9—C8—I2	121.34 (16)	O1—C2—C3	116.9 (2)
O4—C7—C8	120.0 (2)	O4—C13—H13A	109.5
O4—C7—C6	120.2 (2)	O4—C13—H13B	109.5
C8—C7—C6	119.6 (2)	H13A—C13—H13B	109.5
C5—C6—C7	121.1 (2)	O4—C13—H13C	109.5
C5—C6—I1	119.26 (17)	H13A—C13—H13C	109.5
C7—C6—I1	119.58 (16)	H13B—C13—H13C	109.5
O3—C5—C6	119.7 (2)	O3—C12—H12A	109.5
O3—C5—C10	119.6 (2)	O3—C12—H12B	109.5
C6—C5—C10	120.6 (2)	H12A—C12—H12B	109.5
C9—C10—C5	116.74 (19)	O3—C12—H12C	109.5
C9—C10—C4	117.8 (2)	H12A—C12—H12C	109.5
C5—C10—C4	125.4 (2)	H12B—C12—H12C	109.5
O1—C9—C10	121.85 (19)	C4—C11—H11A	109.5
O1—C9—C8	115.25 (19)	C4—C11—H11B	109.5
C10—C9—C8	122.8 (2)	H11A—C11—H11B	109.5
C3—C4—C10	117.8 (2)	C4—C11—H11C	109.5
C3—C4—C11	118.3 (2)	H11A—C11—H11C	109.5
C10—C4—C11	123.7 (2)	H11B—C11—H11C	109.5
C13—O4—C7—C8	−92.8 (3)	C2—O1—C9—C10	1.7 (3)
C13—O4—C7—C6	90.7 (3)	C2—O1—C9—C8	−175.3 (2)
C9—C8—C7—O4	−175.0 (2)	C5—C10—C9—O1	178.85 (19)
I2—C8—C7—O4	−0.1 (3)	C4—C10—C9—O1	−5.2 (3)
C9—C8—C7—C6	1.5 (3)	C5—C10—C9—C8	−4.3 (3)
I2—C8—C7—C6	176.40 (17)	C4—C10—C9—C8	171.7 (2)
O4—C7—C6—C5	174.5 (2)	C7—C8—C9—O1	178.8 (2)
C8—C7—C6—C5	−2.0 (3)	I2—C8—C9—O1	4.0 (3)
O4—C7—C6—I1	−2.4 (3)	C7—C8—C9—C10	1.7 (3)
C8—C7—C6—I1	−178.96 (17)	I2—C8—C9—C10	−173.05 (16)
C12—O3—C5—C6	−85.9 (3)	C9—C10—C4—C3	5.1 (3)
C12—O3—C5—C10	96.7 (3)	C5—C10—C4—C3	−179.3 (2)
C7—C6—C5—O3	−178.0 (2)	C9—C10—C4—C11	−170.8 (3)
I1—C6—C5—O3	−1.1 (3)	C5—C10—C4—C11	4.8 (4)
C7—C6—C5—C10	−0.7 (3)	C10—C4—C3—C2	−1.8 (4)
I1—C6—C5—C10	176.25 (16)	C11—C4—C3—C2	174.3 (3)
O3—C5—C10—C9	−178.92 (19)	C9—O1—C2—O2	179.6 (2)
C6—C5—C10—C9	3.7 (3)	C9—O1—C2—C3	1.7 (3)
O3—C5—C10—C4	5.4 (3)	C4—C3—C2—O2	−179.2 (3)

supplementary materials

C6—C5—C10—C4

−171.9 (2)

C4—C3—C2—O1

−1.7 (4)

Hydrogen-bond geometry (Å, °)

D—H···A

D—H

H···A

D···A

D—H···A

C3—H3···O2ⁱ

0.93


2.53

3.460 (3)

175.

Symmetry codes: (i) $-x, -y, -z$.

Fig. 1



supplementary materials

Fig. 2

