

(5,10,15,20-Tetraphenylporphyrinato- $\kappa^4\text{N}$)cobalt(II)-18-crown-6 (1/1)

Anissa Mansour,^a Mohamed Salah Belkhiria,^a Jean-Claude Daran^b and Habib Nasri^{a*}

^aDépartement de Chimie, Faculté des Sciences de Monastir, Avenue de l'environnement, 5019 Monastir, Tunisia, and ^bLaboratoire de Chimie de Coordination, CNRS UPR 8241, 205 route de Narbonne, 31077 Toulouse, Cedex 04, France

Correspondence e-mail: hnasri1@gmail.com


Received 18 March 2010; accepted 30 March 2010

Key indicators: single-crystal X-ray study; $T = 180\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; disorder in main residue; R factor = 0.037; wR factor = 0.100; data-to-parameter ratio = 14.2.

The asymmetric unit of the title compound, $[\text{Co}(\text{C}_{44}\text{H}_{28}\text{N}_4)] \cdot \text{C}_{12}\text{H}_{24}\text{O}_6$, contains one half of a Co^{II} (TPP) (TPP is tetraphenylporphyrin) complex and one half of an 18-crown-6 molecule of crystallization, both lying on inversion centers. The Co^{II} (TPP) complex exhibits a nearly planar conformation of the porphyrinate core [maximum deviation = 0.069 (2) \AA] with an average $\text{Co}-\text{N}$ distance of 1.971 (4) \AA . The distance between the Co atom and the closest O atom of the 18-crown-6 molecule is 2.533 (2) \AA , indicating a short non-bonded contact between the Co atom and the crown ether molecule. An ethylene group of the 18-crown-6 molecule is disordered over two sites with occupancies of 0.565 (7) and 0.435 (7).

Related literature

For general background to cobalt porphyrin species and their applications, see: Sanders *et al.* (2000). For the synthesis of $\text{Co}(\text{II})$ tetraphenylporphyrin, see: Madure & Scheidt (1976). For related structures, see: Konarev *et al.* (2003, 2004); Nascimento *et al.* (2007); Smirnov *et al.* (1998); Lee *et al.* (2002); Iimura *et al.* (1988). For a description of the Cambridge Structural Database, see: Allen (2002). For the SIMU/ISOR restraints used in the refinement, see: McArdle (1995).



Experimental

Crystal data

$[\text{Co}(\text{C}_{44}\text{H}_{28}\text{N}_4)] \cdot \text{C}_{12}\text{H}_{24}\text{O}_6$

$M_r = 935.95$

Triclinic, $P\bar{1}$

$a = 10.1464 (4)\text{ \AA}$

$b = 11.0890 (6)\text{ \AA}$

$c = 11.7570 (5)\text{ \AA}$

$\alpha = 104.327 (4)^\circ$

$\beta = 105.842 (4)^\circ$

$\gamma = 108.284 (4)^\circ$

$V = 1125.12 (9)\text{ \AA}^3$

$Z = 1$

Mo $K\alpha$ radiation

$\mu = 0.44\text{ mm}^{-1}$

$T = 180\text{ K}$

$0.25 \times 0.24 \times 0.21\text{ mm}$

Data collection

Oxford Diffraction Xcalibur diffractometer

Absorption correction: multi-scan (*CrysAlis RED*; Oxford Diffraction, 2008)

$T_{\min} = 0.927$, $T_{\max} = 1.000$

8862 measured reflections

4589 independent reflections

3977 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$

$wR(F^2) = 0.100$

$S = 1.10$

4589 reflections

323 parameters

30 restraints

H-atom parameters constrained

$\Delta\rho_{\max} = 0.79\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.44\text{ e \AA}^{-3}$

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2008); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2008); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SIR2004* (Burla *et al.*, 2005); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

The authors gratefully acknowledge financial support from the Ministry of Higher Education, Scientific Research and Technology of Tunisia.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2268).

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
- Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.

metal-organic compounds

- Iimura, Y., Sakurai, T. & Yamamoto, K. (1988). *Bull. Chem. Soc. Jpn.*, **61**, 821–826.
- Konarev, D., Khasanov, S. S., Saito, G., Lybovskaya, R. N., Yoshida, Y. & Otsuka, A. (2003). *Chem. Eur. J.*, **9**, 3837–3848.
- Konarev, D., Neretin, I. S., Saito, G., Slovokhotov, Y. L., Otsuka, A. & Lyubovskaya, R. N. (2004). *Chem. Eur. J.*, **9**, 1794–1998.
- Lee, U., Joo, H.-C. & Cho, M.-A. (2002). *Acta Cryst. E* **58**, m599–m601.
- Madure, P. & Scheidt, W. R. (1976). *Inorg. Chem.* **15**, 3182–3184.
- McArdle, P. (1995). *J. Appl. Cryst.* **28**, 65.
- Nascimento B. F. O, Pineiro M., Gonsalves A. M. d'A. Rocha, Silva M. R., Beja A. M. & Paixão A. J. (2007). *J. Porphyrins Phthalocyanines*, **11**, 77–84.
- Oxford Diffraction (2008). *CrystAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.
- Sanders, J.K.M., Bampas N., Clyde-Watson, Z., Kim, H.-J., Mak C. C. & Webb, J. S. (2000). *The Porphyrin Handbook*, Vol. 3, edited by K. M. Kadish, K. M. Smith & R. Guillard, pp. 1–200. San Diego: Academic Press.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Smirnov, V. V., Woller, E. K. & DiMagno, S. G. (1998). *Inorg. Chem.* **37**, 4771–4778.

supplementary materials

Acta Cryst. (2010). E66, m509-m510 [doi:10.1107/S1600536810012080]

(5,10,15,20-Tetraphenylporphyrinato- κ^4N)cobalt(II)-18-crown-6 (1/1)

A. Mansour, M. S. Belkhiria, J.-C. Daran and H. Nasri

Comment

Cobalt atom in metalloporphyrines is commonly used as a qualitatively acceptable substitute for iron atom in high-spin five-coordinate hemes of deoxyhemoglobin and in the low-spin oxygenated hemes of oxyhemoglobin. The metalation of a porphyrin by cobalt (using $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ salt) yields the stable $[\text{Co}^{II}(\text{Porph})]$ (Porph = porphyrin) complex used as starting material in the preparation of five and six-coordinated Co(II) and Co(III) metalloporphyrines (Sanders *et al.*, 2000). In the Cambridge Structural Database (CSD, version 5.31; Allen 2002) there are three structures of tetra-coordinated cobalt(II) tetraphenylporphyrin (TPP) complexes: IKUDOH (Konarev *et al.*, 2003), IXIKIJ (Konarev *et al.*, 2004) and TPORCP12 (Nascimento *et al.*, 2007). Herein we report the structure of the title compound, (I), which has been prepared in our laboratory.

The asymmetric unit of (I), contains one half $[\text{Co}^{II}(\text{TPP})]$ complex and one half crystallographically independent 18-crown-6 molecule of crystallization both lying on inversion centers (Fig. 1).

The distance between the cobalt(II) ion and the symmetry related O1 and O1' atoms (Fig. 2) of the two closest crown ether molecules is 2.533 (2) Å. This distance is significantly longer than the Co^{II} —O(THF) bond length [2.204 (4) Å] in the $[\text{Co}^{II}(\text{F}_8\text{TPP})(\text{THF})_2]$ derivative (where F_8TPP is the tetrakis(pentafluorophenylporphyrin)) (Smirnov *et al.*, 1998) and the Co^{II} —O(H_2O) bond distance in the $[\text{Co}^{II}(\text{H}_2\text{O})_6]^{2+}$ species [2.062 (4)–2.141 (4) Å] (Lee *et al.*, 2002). This indicates that in (I) there is a short non-bonded contact between the cobalt ion and the crown ether molecule.

It has been noticed that there is a relationship between the ruffling of the porphyrinato core and the mean equatorial Co^{II} —N_p distance (N_p = pyrrol N atom); the porphyrinato core is ruffled as the Co—N_p distance decreases, (Iimura *et al.*, 1988). Thus, the average distance Co—N_p in (I), 1.971 (4) Å, is longer than those of the three other reported $[\text{Co}^{II}(\text{TPP})]$ structures quoted above [1.923 (3)–1.969 (6) Å]. The porphyrin core in (I) presents a planar conformation with maximum and minimum deviations from the C₂₀N₄ least-squares plane of 0.069 (2) and -0.068 (2) Å for C6 and C8 atoms, respectively, while the Co^{2+} cation is basically in the porphyrin plane with a Co—C_t distance of 0.004 (1) Å (where C_t is the center of the C₂₀N₄ plane).

Experimental

To a solution of $[\text{Co}^{II}(\text{TPP})]$ (Madure & Scheidt 1976) (100 mg, 0.067 mmol) in chlorobenzene (10 ml) was added an excess of 18-crown-6 (150 mg, 0.567 mmol) and sodium methanolate (100 mg, 3.225 mmol). The reaction mixture was stirred at room temperature until reddish-green solution was formed. The resulting material was crystallized by diffusion of hexanes through the chlorobenzene solution which yielded $[\text{Co}^{II}(\text{TPP})](18\text{-C-6})$ crystals as an impurity.

supplementary materials

Refinement

All H atoms were placed in geometrically idealized positions ($C-H = 0.93\text{--}0.97 \text{\AA}$) and constrained to ride on their parent atoms, with $U(H) = 1.2U_{eq}(C)$. An ethylene group [C24 – C25] of the 18-crown-6 molecule was disordered over two sites with occupancies of 0.565 (7) and 0.435 (7). For this fragment, some anisotropic displacement ellipsoids were rather elongated which led us to use the SIMU/ISOR restraints (McArdle, 1995, Sheldrick, 2008). Alerts B and C for short intramolecular contacts $H\cdots H$ may be explained by the fact that C24 and C25 carbon atoms are disordered.

Figures




Fig. 1. A view of the structure of (I), with the atom-numbering scheme. Unlabeled atoms are related by an inversion center to the labeled atoms. Displacement ellipsoids are drawn at 50%. The H atoms and the minor positions of the disordered C24 and C25 atoms have been omitted for clarity.




Fig. 2. A unit-cell packing of (I). The short non-bonded contacts between the cobalt and the symmetry related O1 and O1' atoms of the two closest crown ether molecules, are drawn by dashed lines. The H atoms and the minor positions of the disordered C24 and C25 atoms have been omitted for clarity.

(5,10,15,20-Tetraphenylporphyrinato- κ^4N)cobalt(II)-18-crown-6 (1/1)

Crystal data

$[Co(C_{44}H_{28}N_4)] \cdot C_{12}H_{24}O_6$	$Z = 1$
$M_r = 935.95$	$F(000) = 491$
Triclinic, $P\bar{1}$	$D_x = 1.381 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 10.1464 (4) \text{ \AA}$	Cell parameters from 6796 reflections
$b = 11.0890 (6) \text{ \AA}$	$\theta = 3.1\text{--}32.2^\circ$
$c = 11.7570 (5) \text{ \AA}$	$\mu = 0.44 \text{ mm}^{-1}$
$\alpha = 104.327 (4)^\circ$	$T = 180 \text{ K}$
$\beta = 105.842 (4)^\circ$	Prism, dark purple
$\gamma = 108.284 (4)^\circ$	$0.25 \times 0.24 \times 0.21 \text{ mm}$
$V = 1125.12 (9) \text{ \AA}^3$	

Data collection

Oxford Diffraction Xcalibur 4589 independent reflections

diffractometer

Radiation source: fine-focus sealed tube

graphite

Detector resolution: 8.2632 pixels mm⁻¹

ω and φ scans

Absorption correction: multi-scan
(CrysAlis RED; Oxford Diffraction, 2008)

$T_{\min} = 0.927$, $T_{\max} = 1.000$

8862 measured reflections

3977 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.019$

$\theta_{\max} = 26.4^\circ$, $\theta_{\min} = 3.3^\circ$

$h = -12 \rightarrow 12$

$k = -13 \rightarrow 9$

$l = -14 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.037$

$wR(F^2) = 0.100$

$S = 1.10$

4589 reflections

323 parameters

30 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.05322P)^2 + 0.3421P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.79 \text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.44 \text{ e \AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of $F^{\wedge}2\wedge$ against ALL reflections. The weighted R -factor wR and goodness of fit S are based on $F^{\wedge}2\wedge$, conventional R -factors R are based on F , with F set to zero for negative $F^{\wedge}2\wedge$. The threshold expression of $F^{\wedge}2\wedge > \sigma(F^{\wedge}2\wedge)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on $F^{\wedge}2\wedge$ are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Co	0.0000	0.5000	0.5000	0.01992 (11)	
N1	-0.10600 (16)	0.51602 (15)	0.34011 (13)	0.0205 (3)	
N2	-0.19511 (16)	0.39511 (15)	0.50211 (13)	0.0202 (3)	
C1	-0.04208 (19)	0.58671 (18)	0.27400 (16)	0.0207 (4)	
C2	-0.1554 (2)	0.57948 (19)	0.16553 (17)	0.0250 (4)	
H2	-0.1398	0.6187	0.1064	0.030*	
C3	-0.2885 (2)	0.50553 (19)	0.16529 (17)	0.0256 (4)	
H3	-0.3826	0.4838	0.1059	0.031*	
C4	-0.25852 (19)	0.46615 (18)	0.27385 (16)	0.0213 (4)	
C5	-0.36903 (19)	0.39375 (18)	0.30882 (16)	0.0217 (4)	

supplementary materials

C6	-0.33699 (19)	0.36035 (18)	0.41648 (16)	0.0216 (4)	
C7	-0.4490 (2)	0.2741 (2)	0.44746 (18)	0.0268 (4)	
H7	-0.5525	0.2383	0.4042	0.032*	
C8	-0.3764 (2)	0.2551 (2)	0.55053 (18)	0.0276 (4)	
H8	-0.4199	0.2024	0.5917	0.033*	
C9	-0.21943 (19)	0.33090 (18)	0.58552 (16)	0.0217 (4)	
C10	0.1099 (2)	0.66163 (18)	0.30894 (16)	0.0213 (4)	
C11	0.15851 (19)	0.74003 (18)	0.23032 (16)	0.0224 (4)	
C12	0.1820 (2)	0.6798 (2)	0.12439 (18)	0.0293 (4)	
H12	0.1656	0.5883	0.1003	0.035*	
C13	0.2294 (2)	0.7541 (2)	0.05387 (19)	0.0339 (5)	
H13	0.2456	0.7125	-0.0169	0.041*	
C14	0.2529 (2)	0.8891 (2)	0.08771 (19)	0.0322 (4)	
H14	0.2848	0.9390	0.0401	0.039*	
C15	0.2289 (3)	0.9494 (2)	0.1922 (2)	0.0387 (5)	
H15	0.2439	1.0406	0.2153	0.046*	
C16	0.1824 (3)	0.8759 (2)	0.26378 (19)	0.0347 (5)	
H16	0.1672	0.9181	0.3349	0.042*	
C17	-0.52953 (19)	0.34897 (18)	0.22752 (17)	0.0229 (4)	
C18	-0.5921 (2)	0.2539 (2)	0.10469 (18)	0.0277 (4)	
H18	-0.5327	0.2183	0.0722	0.033*	
C19	-0.7417 (2)	0.2114 (2)	0.03001 (19)	0.0346 (5)	
H19	-0.7823	0.1476	-0.0522	0.042*	
C20	-0.8304 (2)	0.2635 (2)	0.0775 (2)	0.0371 (5)	
H20	-0.9312	0.2343	0.0279	0.045*	
C21	-0.7694 (2)	0.3590 (2)	0.1985 (2)	0.0372 (5)	
H21	-0.8289	0.3949	0.2304	0.045*	
C22	-0.6198 (2)	0.4018 (2)	0.27297 (19)	0.0308 (4)	
H22	-0.5793	0.4668	0.3545	0.037*	
O1	1.00507 (17)	0.72687 (17)	0.61845 (14)	0.0415 (4)	
O2	0.7723 (2)	0.8744 (2)	0.46275 (19)	0.0633 (5)	
O3	0.75479 (18)	0.98421 (16)	0.26483 (16)	0.0438 (4)	
C23	1.1145 (3)	0.8022 (2)	0.7438 (2)	0.0399 (5)	
H23A	1.1340	0.7388	0.7828	0.048*	
H23B	1.0739	0.8534	0.7930	0.048*	
C24A	0.8779 (5)	0.7518 (5)	0.5733 (4)	0.0333 (11)	0.565 (7)
H24A	0.8463	0.7817	0.6419	0.040*	0.565 (7)
H24B	0.7962	0.6691	0.5084	0.040*	0.565 (7)
C25A	0.9181 (5)	0.8618 (5)	0.5181 (5)	0.0465 (14)	0.565 (7)
H25A	0.9936	0.9474	0.5837	0.056*	0.565 (7)
H25B	0.9557	0.8353	0.4528	0.056*	0.565 (7)
C24B	0.9331 (11)	0.8256 (10)	0.5946 (9)	0.071 (2)	0.435 (7)
H24C	0.9026	0.8563	0.6639	0.085*	0.435 (7)
H24D	1.0063	0.9049	0.5921	0.085*	0.435 (7)
C25B	0.8162 (10)	0.7669 (8)	0.4875 (8)	0.075 (3)	0.435 (7)
H25C	0.7347	0.6982	0.4942	0.089*	0.435 (7)
H25D	0.8415	0.7234	0.4193	0.089*	0.435 (7)
C26	0.6814 (3)	0.8027 (2)	0.3354 (3)	0.0493 (6)	
H26A	0.5947	0.7262	0.3264	0.059*	

H26B	0.7366	0.7676	0.2905	0.059*
C27	0.6318 (3)	0.8965 (3)	0.2809 (2)	0.0465 (6)
H27A	0.5481	0.8445	0.1998	0.056*
H27B	0.5997	0.9493	0.3378	0.056*
C28	0.7410 (3)	1.1016 (2)	0.2499 (2)	0.0396 (5)
H28A	0.7135	1.1454	0.3164	0.048*
H28B	0.6623	1.0768	0.1687	0.048*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co	0.01575 (17)	0.0281 (2)	0.01649 (18)	0.00707 (14)	0.00525 (13)	0.01245 (14)
N1	0.0169 (7)	0.0242 (8)	0.0186 (7)	0.0062 (6)	0.0053 (6)	0.0095 (6)
N2	0.0202 (7)	0.0235 (8)	0.0168 (7)	0.0083 (6)	0.0053 (6)	0.0097 (6)
C1	0.0227 (9)	0.0237 (9)	0.0171 (8)	0.0096 (7)	0.0075 (7)	0.0100 (7)
C2	0.0272 (9)	0.0296 (10)	0.0191 (9)	0.0111 (8)	0.0067 (7)	0.0136 (8)
C3	0.0230 (9)	0.0310 (10)	0.0207 (9)	0.0104 (8)	0.0037 (7)	0.0120 (8)
C4	0.0204 (8)	0.0243 (9)	0.0177 (8)	0.0092 (7)	0.0043 (7)	0.0086 (7)
C5	0.0185 (8)	0.0234 (9)	0.0194 (8)	0.0072 (7)	0.0039 (7)	0.0073 (7)
C6	0.0173 (8)	0.0242 (9)	0.0199 (9)	0.0063 (7)	0.0050 (7)	0.0077 (7)
C7	0.0176 (8)	0.0329 (10)	0.0262 (9)	0.0054 (8)	0.0068 (7)	0.0132 (8)
C8	0.0220 (9)	0.0337 (10)	0.0260 (9)	0.0057 (8)	0.0097 (8)	0.0161 (8)
C9	0.0201 (8)	0.0248 (9)	0.0199 (9)	0.0070 (7)	0.0084 (7)	0.0098 (7)
C10	0.0244 (9)	0.0231 (9)	0.0187 (8)	0.0094 (7)	0.0095 (7)	0.0100 (7)
C11	0.0193 (8)	0.0274 (9)	0.0194 (8)	0.0066 (7)	0.0056 (7)	0.0124 (7)
C12	0.0378 (11)	0.0312 (10)	0.0279 (10)	0.0178 (9)	0.0161 (8)	0.0163 (8)
C13	0.0407 (12)	0.0461 (13)	0.0285 (10)	0.0223 (10)	0.0206 (9)	0.0215 (10)
C14	0.0288 (10)	0.0374 (11)	0.0275 (10)	0.0053 (9)	0.0080 (8)	0.0211 (9)
C15	0.0509 (13)	0.0244 (10)	0.0315 (11)	0.0065 (9)	0.0108 (10)	0.0123 (9)
C16	0.0486 (12)	0.0299 (11)	0.0245 (10)	0.0126 (9)	0.0159 (9)	0.0103 (8)
C17	0.0196 (8)	0.0249 (9)	0.0233 (9)	0.0071 (7)	0.0049 (7)	0.0138 (8)
C18	0.0258 (9)	0.0305 (10)	0.0242 (9)	0.0102 (8)	0.0063 (8)	0.0113 (8)
C19	0.0292 (10)	0.0346 (11)	0.0259 (10)	0.0033 (9)	-0.0010 (8)	0.0139 (9)
C20	0.0183 (9)	0.0468 (13)	0.0425 (12)	0.0075 (9)	0.0014 (8)	0.0290 (11)
C21	0.0258 (10)	0.0482 (13)	0.0492 (13)	0.0203 (10)	0.0163 (9)	0.0272 (11)
C22	0.0269 (10)	0.0347 (11)	0.0297 (10)	0.0121 (8)	0.0093 (8)	0.0122 (9)
O1	0.0360 (8)	0.0523 (10)	0.0342 (8)	0.0179 (7)	0.0067 (7)	0.0202 (7)
O2	0.0793 (12)	0.0724 (11)	0.0566 (10)	0.0555 (10)	0.0198 (9)	0.0282 (9)
O3	0.0461 (9)	0.0420 (9)	0.0594 (11)	0.0225 (8)	0.0307 (8)	0.0269 (8)
C23	0.0528 (14)	0.0367 (12)	0.0284 (11)	0.0177 (11)	0.0121 (10)	0.0137 (9)
C24A	0.032 (2)	0.036 (2)	0.035 (2)	0.0101 (18)	0.0153 (17)	0.0186 (18)
C25A	0.034 (2)	0.042 (3)	0.051 (3)	0.0053 (19)	0.0043 (19)	0.024 (2)
C24B	0.070 (3)	0.066 (3)	0.075 (3)	0.030 (2)	0.0147 (19)	0.035 (2)
C25B	0.074 (3)	0.068 (3)	0.080 (3)	0.035 (2)	0.0159 (19)	0.032 (2)
C26	0.0573 (15)	0.0349 (13)	0.0629 (17)	0.0176 (11)	0.0343 (13)	0.0181 (12)
C27	0.0293 (11)	0.0565 (15)	0.0486 (14)	0.0107 (10)	0.0121 (10)	0.0230 (12)
C28	0.0430 (13)	0.0389 (12)	0.0328 (11)	0.0201 (10)	0.0071 (10)	0.0104 (10)

supplementary materials

Geometric parameters (\AA , $^\circ$)

Co—N2	1.9669 (14)	C18—C19	1.385 (3)
Co—N2 ⁱ	1.9669 (14)	C18—H18	0.9300
Co—N1 ⁱ	1.9757 (14)	C19—C20	1.377 (3)
Co—N1	1.9757 (14)	C19—H19	0.9300
N1—C1	1.377 (2)	C20—C21	1.375 (3)
N1—C4	1.378 (2)	C20—H20	0.9300
N2—C9	1.377 (2)	C21—C22	1.383 (3)
N2—C6	1.379 (2)	C21—H21	0.9300
C1—C10	1.385 (2)	C22—H22	0.9300
C1—C2	1.430 (2)	O1—C24A	1.394 (4)
C2—C3	1.341 (3)	O1—C23	1.420 (3)
C2—H2	0.9300	O1—C24B	1.534 (8)
C3—C4	1.435 (2)	O2—C26	1.385 (3)
C3—H3	0.9300	O2—C25B	1.462 (7)
C4—C5	1.386 (2)	O2—C25A	1.510 (5)
C5—C6	1.388 (2)	O3—C28	1.399 (3)
C5—C17	1.490 (2)	O3—C27	1.413 (3)
C6—C7	1.434 (2)	C23—C28 ⁱⁱ	1.488 (3)
C7—C8	1.339 (3)	C23—H23A	0.9700
C7—H7	0.9300	C23—H23B	0.9700
C8—C9	1.429 (2)	C24A—C25A	1.517 (6)
C8—H8	0.9300	C24A—H24A	0.9700
C9—C10 ⁱ	1.386 (2)	C24A—H24B	0.9700
C10—C9 ⁱ	1.386 (2)	C25A—H25A	0.9700
C10—C11	1.494 (2)	C25A—H25B	0.9700
C11—C12	1.380 (3)	C24B—C25B	1.308 (12)
C11—C16	1.381 (3)	C24B—H24C	0.9700
C12—C13	1.380 (3)	C24B—H24D	0.9700
C12—H12	0.9300	C25B—H25C	0.9700
C13—C14	1.373 (3)	C25B—H25D	0.9700
C13—H13	0.9300	C26—C27	1.493 (3)
C14—C15	1.370 (3)	C26—H26A	0.9700
C14—H14	0.9300	C26—H26B	0.9700
C15—C16	1.382 (3)	C27—H27A	0.9700
C15—H15	0.9300	C27—H27B	0.9700
C16—H16	0.9300	C28—C23 ⁱⁱ	1.488 (3)
C17—C22	1.384 (3)	C28—H28A	0.9700
C17—C18	1.388 (3)	C28—H28B	0.9700
N2—Co—N2 ⁱ	180.0	C20—C19—C18	119.9 (2)
N2—Co—N1 ⁱ	90.63 (6)	C20—C19—H19	120.0
N2 ⁱ —Co—N1 ⁱ	89.37 (6)	C18—C19—H19	120.0
N2—Co—N1	89.37 (6)	C21—C20—C19	119.82 (18)
N2 ⁱ —Co—N1	90.63 (6)	C21—C20—H20	120.1
N1 ⁱ —Co—N1	180.0	C19—C20—H20	120.1


C1—N1—C4	104.78 (14)	C20—C21—C22	120.2 (2)
C1—N1—Co	126.92 (12)	C20—C21—H21	119.9
C4—N1—Co	128.22 (12)	C22—C21—H21	119.9
C9—N2—C6	104.48 (14)	C21—C22—C17	120.75 (19)
C9—N2—Co	126.91 (12)	C21—C22—H22	119.6
C6—N2—Co	128.44 (12)	C17—C22—H22	119.6
N1—C1—C10	125.90 (16)	C24A—O1—C23	122.1 (2)
N1—C1—C2	110.67 (15)	C23—O1—C24B	102.2 (4)
C10—C1—C2	123.35 (16)	C26—O2—C25B	97.5 (4)
C3—C2—C1	107.06 (16)	C26—O2—C25A	120.3 (3)
C3—C2—H2	126.5	C28—O3—C27	114.26 (18)
C1—C2—H2	126.5	O1—C23—C28 ⁱⁱ	113.86 (18)
C2—C3—C4	107.11 (16)	O1—C23—H23A	108.8
C2—C3—H3	126.4	C28 ⁱⁱ —C23—H23A	108.8
C4—C3—H3	126.4	O1—C23—H23B	108.8
N1—C4—C5	125.58 (16)	C28 ⁱⁱ —C23—H23B	108.8
N1—C4—C3	110.38 (15)	H23A—C23—H23B	107.7
C5—C4—C3	123.97 (16)	O1—C24A—C25A	107.9 (3)
C4—C5—C6	122.88 (16)	O1—C24A—H24A	110.1
C4—C5—C17	118.89 (16)	C25A—C24A—H24A	110.1
C6—C5—C17	118.23 (16)	O1—C24A—H24B	110.1
N2—C6—C5	125.50 (16)	C25A—C24A—H24B	110.1
N2—C6—C7	110.62 (15)	H24A—C24A—H24B	108.4
C5—C6—C7	123.69 (16)	O2—C25A—C24A	104.3 (3)
C8—C7—C6	106.93 (16)	O2—C25A—H25A	110.9
C8—C7—H7	126.5	C24A—C25A—H25A	110.9
C6—C7—H7	126.5	O2—C25A—H25B	110.9
C7—C8—C9	107.22 (16)	C24A—C25A—H25B	110.9
C7—C8—H8	126.4	H25A—C25A—H25B	108.9
C9—C8—H8	126.4	C25B—C24B—O1	111.0 (8)
N2—C9—C10 ⁱ	126.03 (16)	C25B—C24B—H24C	109.4
N2—C9—C8	110.74 (15)	O1—C24B—H24C	109.4
C10 ⁱ —C9—C8	123.23 (16)	C25B—C24B—H24D	109.4
C1—C10—C9 ⁱ	123.25 (16)	O1—C24B—H24D	109.4
C1—C10—C11	118.58 (15)	H24C—C24B—H24D	108.0
C9 ⁱ —C10—C11	118.16 (16)	C24B—C25B—O2	106.7 (7)
C12—C11—C16	118.63 (17)	C24B—C25B—H25C	110.4
C12—C11—C10	121.28 (17)	O2—C25B—H25C	110.4
C16—C11—C10	120.09 (17)	C24B—C25B—H25D	110.4
C11—C12—C13	120.67 (18)	O2—C25B—H25D	110.4
C11—C12—H12	119.7	H25C—C25B—H25D	108.6
C13—C12—H12	119.7	O2—C26—C27	108.8 (2)
C14—C13—C12	120.34 (19)	O2—C26—H26A	109.9
C14—C13—H13	119.8	C27—C26—H26A	109.9
C12—C13—H13	119.8	O2—C26—H26B	109.9
C15—C14—C13	119.39 (18)	C27—C26—H26B	109.9
C15—C14—H14	120.3	H26A—C26—H26B	108.3

supplementary materials

C13—C14—H14	120.3	O3—C27—C26	107.81 (19)
C14—C15—C16	120.54 (19)	O3—C27—H27A	110.1
C14—C15—H15	119.7	C26—C27—H27A	110.1
C16—C15—H15	119.7	O3—C27—H27B	110.1
C11—C16—C15	120.43 (19)	C26—C27—H27B	110.1
C11—C16—H16	119.8	H27A—C27—H27B	108.5
C15—C16—H16	119.8	O3—C28—C23 ⁱⁱ	110.08 (18)
C22—C17—C18	118.42 (17)	O3—C28—H28A	109.6
C22—C17—C5	120.57 (17)	C23 ⁱⁱ —C28—H28A	109.6
C18—C17—C5	121.01 (17)	O3—C28—H28B	109.6
C19—C18—C17	120.81 (19)	C23 ⁱⁱ —C28—H28B	109.6
C19—C18—H18	119.6	H28A—C28—H28B	108.2
C17—C18—H18	119.6		

Symmetry codes: (i) $-x, -y+1, -z+1$; (ii) $-x+2, -y+2, -z+1$.

Fig. 1



supplementary materials

Fig. 2

