

Dichlorido[(*R,R*)-*N*¹,*N*¹,*N*²-tribenzyl-cyclohexane-1,2-diamine- κ^2 *N*¹,*N*²]-copper(II)

Quang Trung Nguyen and Jong Hwa Jeong*

Department of Chemistry, Kyungpook National University, Taegu 702-701, Republic of Korea

Correspondence e-mail: jeongjh@knu.ac.kr


Received 26 March 2010; accepted 16 April 2010

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(C-C) = 0.005$ Å; R factor = 0.031; wR factor = 0.083; data-to-parameter ratio = 16.9.

In the title compound, $[CuCl_2(C_{27}H_{32}N_2)]$, which bears a chiral diamine ligand, *viz* (*R,R*)-*N,N,N'*-tribenzylcyclohexane-1,2-diamine, the Cu^{II} ion is ligated by two N and two Cl atoms in a distorted square-planar geometry. The coordination of the ligands to the Cu^{II} ion results in the formation of a five-membered heterocyclic ring and a chiral center at the monosubstituted nitrogen in an (*S*)-configuration. The catalytic capacity of the complex for the asymmetric nitroaldol reaction is promising (49% ee).

Related literature

For the synthesis of *N,N,N'*-tribenzyl-(*R,R*)-1,2-diaminocyclohexane, see: Tye *et al.* (2002); Boyd *et al.* (2005). For related structures, see: Alexakis *et al.* (2001); Tye *et al.* (2002); Boyd *et al.* (2005, 2006); Arjan *et al.* (2005); Brethon *et al.* (2004); Jones & Mahon (2008); Evans & Seidel (2005); Evans *et al.* (2007); Roh *et al.* (2004); Nguyen & Jeong (2008a,b).



Experimental

Crystal data

$[CuCl_2(C_{27}H_{32}N_2)]$
 $M_r = 519.00$
Orthorhombic, $P2_12_12_1$
 $a = 10.5806$ (7) Å
 $b = 15.4409$ (8) Å
 $c = 16.2579$ (12) Å

$V = 2656.1$ (3) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 1.04$ mm⁻¹
 $T = 295$ K
 $0.40 \times 0.40 \times 0.40$ mm

Data collection

Enraf–Nonius CAD-4 diffractometer
Absorption correction: analytical (*ABSCALC*; McArdle & Daly, 1999)
 $T_{min} = 0.660$, $T_{max} = 0.666$

5793 measured reflections
4931 independent reflections
3885 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.019$
3 standard reflections every 60 min
intensity decay: none

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$
 $wR(F^2) = 0.083$
 $S = 1.06$
4931 reflections
292 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.33$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.24$ e Å⁻³
Absolute structure: Flack (1983)
Flack parameter: -0.017 (13)

Data collection: *CAD4* (Enraf–Nonius, 1989); cell refinement: *CAD4*; data reduction: *XCAD* (McArdle, 1999); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEPIII* (Burnett & Johnson, 1996); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

This research was supported by Kyungpook National University Research Fund, 2008.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2198).

References

- Alexakis, A., Chauvin, A. S., Stouvenel, R., Vrancken, E., Mutti, S. & Mangeney, P. (2001). *Tetrahedron: Asymmetry*, **12**, 1171–1178.
- Arjan, H., Boyd, E., Coumbarides, G. S., Eames, J., Jones, R. V. H., Stenson, R. A. & Suggate, M. J. (2005). *Tetrahedron Lett.* **46**, 1921–1925.
- Boyd, E., Coumbarides, G. S., Eames, J., Jones, R. V. H., Mottevalli, M., Stenson, R. A. & Suggate, M. J. (2006). *J. Chem. Crystallogr.* **36**, 263–269.
- Boyd, E., Coumbarides, G. S., Eames, J., Jones, R. V. H., Stenson, R. A. & Suggate, M. J. (2005). *Tetrahedron Lett.* **46**, 3479–3484.
- Brethon, A., Moreau, J. J. E. & Man, M. W. C. (2004). *Tetrahedron: Asymmetry*, **15**, 495–502.
- Burnett, M. N. & Johnson, C. K. (1996). *ORTEPIII*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
- Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
- Evans, D. A., Mito, S. & Seidel, D. (2007). *J. Am. Chem. Soc.* **129**, 11583–11592.
- Evans, D. A. & Seidel, D. (2005). *J. Am. Chem. Soc.* **127**, 9958–9959.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Jones, M. D. & Mahon, M. F. (2008). *J. Organomet. Chem.* **693**, 2377–2382.
- McArdle, P. (1999). *XCAD*. National University of Ireland, Galway, Ireland.
- McArdle, P. & Daly, P. (1999). *ABSCALC*. National University of Ireland, Galway, Ireland.
- Nguyen, Q. T. & Jeong, J. H. (2008a). *Bull. Korean Chem. Soc.* **29**, 483–486.
- Nguyen, Q. T. & Jeong, J. H. (2008b). *Polyhedron*, **27**, 3227–3230.
- Roh, S. G., Yoon, J. U. & Jeong, J. H. (2004). *Polyhedron*, **23**, 2063–2067.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Tye, H., Eldred, C. & Wills, M. (2002). *Tetrahedron Lett.* **43**, 155–158.

supplementary materials

Acta Cryst. (2010). E66, m565 [doi:10.1107/S1600536810014054]

Dichlorido[(*R,R*)-*N*¹,*N*¹,*N*²-tribenzylcyclohexane-1,2-diamine- κ^2 *N*¹,*N*²]copper(II)

Q. T. Nguyen and J. H. Jeong

Comment

Disubstituted, trisubstituted and tetrasubstituted (*R,R*)-1,2-diaminocyclohexane were synthesized (Alexakis *et al.*, 2001; Tye *et al.*, 2002; Boyd *et al.*, 2005, 2006; Arjan *et al.*, 2005). Especially disubstituted chiral diamine ligands with Rh (Brethon *et al.*, 2004; Jones & Mahon, 2008), Ni (Evans & Seidel, 2005; Evans *et al.*, 2007), Zn (Roh *et al.*, 2004; Nguyen & Jeong, 2008a), Cu (Nguyen & Jeong, 2008b) were extensively applied in asymmetric synthesis. However, the coordination chemistry and application of asymmetric trisubstituted chiral 1,2-diaminocyclohexanes containing a secondary and a tertiary amines had not attended much. In this study, a new complex of Cu(II) containing *N,N,N'*-tribenzyl-(*R,R*)-1,2-diaminocyclohexane (Tye *et al.*, 2002; Boyd *et al.*, 2005) was synthesized and its molecular and crystal structures were determined.

Also, capability of the complex as an enantioselective catalyst for asymmetric nitroaldol reaction was examined. The copper ion was ligated by two nitrogen and two chloride atoms in distorted square-planar geometry. The coordination of the ligands to the Cu ion induced a 5-membered heterocyclic ring and a chiral center at monosubstituted nitrogen in (*S*)-configuration. Catalytic capacity of the complex for asymmetric nitroaldol reaction was promising (49% ee {ee = [R - S / R+S] x 100 or [S - R / R+S] x 100}).

Experimental

A solution of *N,N,N'*-tribenzyl-(*R,R*)-1,2-diaminocyclohexane (1.57 g, 4.08 mmol) in ethanol (5 ml) was added slowly to a solution of CuCl₂·2H₂O (0.69 g, 4.01 mmol) in ethanol (10 ml) Tye *et al.*, (2002); Boyd *et al.*, (2005). The mixture was stirred overnight at ambient temperature. The solvent was removed to yield blue solids. The product was re-crystallized from anhydrous ethanol to afford blue crystals (1.64 g, yield 79%). Anal. Calc. for C₂₇H₃₂Cl₂CuN₂: C 62.48, H 6.21, N 5.40 and found: C 62.20, H 6.30, N 5.46%.

Refinement

H-atom of N—H was refined with U_{iso}(H) = 1.2U_{eq}(N). All H-atoms placed on C atoms were positioned geometrically and refined using a riding model with C—H = 0.97 Å for methylene, C—H = 0.98 Å for methine, C—H = 0.93 Å for aromatic H atoms. For all H atoms U_{iso}(H) = 1.2U_{eq}(C).

In the crystal structure was found 'accessible void' with volume 54.00 Å³.

supplementary materials

Figures




Fig. 1. A view of title compound molecule with the atom numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. Hydrogen atoms are presented as a small spheres of arbitrary radius.

Dichlorido[(*R,R*)-*N*¹,*N*¹,*N*²- tribenzylcyclohexane-1,2-diamine- κ^2 *N*¹,*N*²]copper(II)

Crystal data

[CuCl ₂ (C ₂₇ H ₃₂ N ₂)]	<i>F</i> (000) = 1084
<i>M_r</i> = 519.00	<i>D_x</i> = 1.298 Mg m ⁻³
Orthorhombic, <i>P</i> 2 ₁ 2 ₁ 2 ₁	Mo <i>K</i> α radiation, λ = 0.71073 Å
Hall symbol: P 2ac 2ab	Cell parameters from 25 reflections
<i>a</i> = 10.5806 (7) Å	θ = 10–13°
<i>b</i> = 15.4409 (8) Å	μ = 1.04 mm ⁻¹
<i>c</i> = 16.2579 (12) Å	<i>T</i> = 295 K
<i>V</i> = 2656.1 (3) Å ³	Block, blue
<i>Z</i> = 4	0.40 × 0.40 × 0.40 mm

Data collection

Enraf–Nonius CAD-4 diffractometer	3885 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	R_{int} = 0.019
graphite	$\theta_{\text{max}} = 25.5^\circ$, $\theta_{\text{min}} = 1.8^\circ$
$\omega/2\theta$ scans	$h = -12 \rightarrow 12$
Absorption correction: analytical (<i>ABSCALC</i> ; McArdle & Daly, 1999)	$k = -18 \rightarrow 18$
$T_{\text{min}} = 0.660$, $T_{\text{max}} = 0.666$	$l = -19 \rightarrow 19$
5793 measured reflections	3 standard reflections every 60 min
4931 independent reflections	intensity decay: none

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.031$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.083$	$w = 1/[\sigma^2(F_o^2) + (0.0488P)^2]$
$S = 1.06$	where $P = (F_o^2 + 2F_c^2)/3$
	$(\Delta/\sigma)_{\text{max}} = 0.001$

4931 reflections	$\Delta\rho_{\max} = 0.33 \text{ e \AA}^{-3}$
292 parameters	$\Delta\rho_{\min} = -0.24 \text{ e \AA}^{-3}$
0 restraints	Absolute structure: Flack (1983)
Primary atom site location: structure-invariant direct methods	Flack parameter: -0.017 (13)

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu	0.03338 (3)	0.43742 (2)	0.76295 (2)	0.03853 (10)
Cl1	-0.03484 (8)	0.33981 (5)	0.85529 (5)	0.05231 (19)
Cl2	-0.10262 (9)	0.40812 (6)	0.66326 (6)	0.0609 (2)
N1	0.0744 (2)	0.56146 (15)	0.72591 (14)	0.0361 (5)
N2	0.2032 (2)	0.43863 (17)	0.81961 (17)	0.0403 (5)
H2	0.193 (3)	0.438 (2)	0.871 (2)	0.048*
C1	0.1701 (3)	0.59164 (18)	0.78886 (17)	0.0352 (6)
H1	0.1234	0.5982	0.8406	0.042*
C2	0.2323 (3)	0.67928 (19)	0.7726 (2)	0.0476 (7)
H2A	0.2862	0.6755	0.7245	0.057*
H2B	0.1679	0.7226	0.7621	0.057*
C3	0.3099 (4)	0.7054 (2)	0.8468 (2)	0.0561 (9)
H3A	0.3490	0.7612	0.8368	0.067*
H3B	0.2552	0.7108	0.8944	0.067*
C4	0.4109 (3)	0.6388 (2)	0.8639 (2)	0.0602 (9)
H4A	0.4581	0.6556	0.9125	0.072*
H4B	0.4692	0.6366	0.8179	0.072*
C5	0.3532 (3)	0.5492 (2)	0.8772 (2)	0.0489 (8)
H5A	0.3028	0.5498	0.9271	0.059*
H5B	0.4204	0.5072	0.8845	0.059*
C6	0.2699 (3)	0.52161 (18)	0.8048 (2)	0.0373 (7)
H6	0.3229	0.5156	0.7557	0.045*
C7	-0.0463 (3)	0.61569 (18)	0.73180 (19)	0.0433 (7)
H7A	-0.0267	0.6739	0.7135	0.052*
H7B	-0.1075	0.5919	0.6936	0.052*
C8	-0.1073 (3)	0.62181 (19)	0.8143 (2)	0.0428 (7)
C9	-0.1934 (3)	0.5605 (2)	0.8410 (2)	0.0544 (8)
H9	-0.2095	0.5122	0.8085	0.065*
C10	-0.2553 (3)	0.5697 (3)	0.9143 (3)	0.0645 (10)
H10	-0.3108	0.5267	0.9317	0.077*
C11	-0.2371 (4)	0.6404 (3)	0.9621 (2)	0.0628 (10)
H11	-0.2812	0.6469	1.0112	0.075*

supplementary materials

C12	-0.1515 (4)	0.7031 (3)	0.9366 (2)	0.0609 (10)
H12	-0.1373	0.7517	0.9692	0.073*
C13	-0.0880 (4)	0.6938 (2)	0.8638 (2)	0.0530 (9)
H13	-0.0312	0.7364	0.8473	0.064*
C14	0.1132 (3)	0.5719 (2)	0.63786 (17)	0.0465 (7)
H14A	0.1232	0.6333	0.6273	0.056*
H14B	0.0438	0.5518	0.6038	0.056*
C15	0.2318 (3)	0.5269 (2)	0.60886 (19)	0.0502 (8)
C16	0.2368 (4)	0.4375 (3)	0.59948 (19)	0.0564 (8)
H16	0.1669	0.4039	0.6127	0.068*
C17	0.3453 (4)	0.3988 (3)	0.5705 (3)	0.0759 (12)
H17	0.3495	0.3388	0.5664	0.091*
C18	0.4472 (5)	0.4482 (4)	0.5478 (3)	0.1003 (17)
H18	0.5202	0.4216	0.5284	0.120*
C19	0.4416 (5)	0.5367 (4)	0.5536 (3)	0.0976 (17)
H19	0.5098	0.5703	0.5368	0.117*
C20	0.3350 (4)	0.5754 (3)	0.5844 (2)	0.0744 (12)
H20	0.3320	0.6354	0.5888	0.089*
C21	0.2727 (3)	0.3570 (2)	0.7986 (2)	0.0551 (9)
H21A	0.2184	0.3082	0.8114	0.066*
H21B	0.2874	0.3562	0.7397	0.066*
C22	0.3962 (3)	0.34379 (19)	0.8410 (2)	0.0432 (7)
C23	0.4030 (4)	0.3147 (3)	0.9210 (2)	0.0631 (10)
H23	0.3292	0.3016	0.9495	0.076*
C24	0.5196 (5)	0.3048 (2)	0.9596 (2)	0.0737 (12)
H24	0.5238	0.2849	1.0135	0.088*
C25	0.6279 (4)	0.3247 (3)	0.9178 (3)	0.0705 (12)
H25	0.7056	0.3197	0.9440	0.085*
C26	0.6229 (3)	0.3515 (2)	0.8387 (3)	0.0621 (10)
H26	0.6970	0.3628	0.8099	0.075*
C27	0.5094 (3)	0.3617 (2)	0.8020 (2)	0.0517 (8)
H27	0.5073	0.3816	0.7480	0.062*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu	0.03398 (17)	0.03603 (16)	0.04556 (19)	0.00039 (16)	-0.00646 (17)	-0.00163 (15)
Cl1	0.0463 (4)	0.0478 (4)	0.0629 (5)	-0.0078 (4)	-0.0052 (4)	0.0079 (3)
Cl2	0.0622 (5)	0.0594 (5)	0.0612 (5)	-0.0084 (4)	-0.0245 (4)	-0.0054 (4)
N1	0.0349 (11)	0.0387 (11)	0.0348 (12)	0.0045 (10)	-0.0041 (9)	-0.0005 (11)
N2	0.0347 (12)	0.0368 (12)	0.0494 (14)	0.0047 (12)	-0.0050 (11)	-0.0005 (13)
C1	0.0369 (15)	0.0370 (15)	0.0317 (15)	0.0021 (12)	-0.0037 (12)	-0.0027 (11)
C2	0.0491 (17)	0.0375 (15)	0.0563 (19)	-0.0031 (13)	-0.0058 (16)	-0.0004 (15)
C3	0.060 (2)	0.0404 (18)	0.068 (2)	-0.0031 (16)	-0.0085 (19)	-0.0075 (16)
C4	0.0491 (19)	0.056 (2)	0.075 (2)	-0.0027 (17)	-0.0152 (18)	-0.0130 (18)
C5	0.0405 (17)	0.046 (2)	0.060 (2)	0.0053 (14)	-0.0171 (15)	-0.0075 (15)
C6	0.0308 (15)	0.0376 (15)	0.0435 (17)	0.0020 (13)	-0.0003 (13)	-0.0048 (13)
C7	0.0416 (16)	0.0432 (15)	0.0452 (15)	0.0113 (13)	-0.0069 (16)	0.0001 (13)

C8	0.0360 (16)	0.0383 (16)	0.0540 (19)	0.0081 (14)	-0.0024 (15)	0.0009 (14)
C9	0.0398 (17)	0.0482 (18)	0.075 (2)	-0.0005 (17)	-0.0008 (17)	-0.0117 (19)
C10	0.0439 (19)	0.062 (2)	0.087 (3)	-0.0029 (19)	0.0155 (18)	0.007 (2)
C11	0.058 (2)	0.071 (3)	0.060 (2)	0.016 (2)	0.0131 (18)	0.004 (2)
C12	0.070 (2)	0.055 (2)	0.057 (2)	0.0071 (19)	0.0078 (19)	-0.0099 (17)
C13	0.061 (2)	0.0385 (17)	0.059 (2)	0.0054 (15)	0.0057 (17)	0.0001 (15)
C14	0.0547 (18)	0.0492 (18)	0.0357 (15)	0.0029 (17)	0.0005 (14)	0.0012 (14)
C15	0.061 (2)	0.058 (2)	0.0313 (16)	-0.0050 (17)	0.0085 (15)	-0.0060 (14)
C16	0.065 (2)	0.061 (2)	0.0434 (18)	-0.002 (2)	0.0110 (15)	-0.0125 (18)
C17	0.081 (3)	0.075 (3)	0.072 (3)	0.008 (2)	0.012 (2)	-0.027 (2)
C18	0.075 (3)	0.127 (4)	0.099 (3)	0.005 (3)	0.036 (3)	-0.035 (3)
C19	0.082 (3)	0.115 (4)	0.096 (3)	-0.022 (3)	0.047 (3)	-0.017 (3)
C20	0.085 (3)	0.076 (3)	0.062 (2)	-0.015 (2)	0.029 (2)	-0.006 (2)
C21	0.0438 (19)	0.0413 (18)	0.080 (2)	0.0136 (15)	-0.0093 (18)	-0.0090 (17)
C22	0.0367 (16)	0.0350 (15)	0.058 (2)	0.0057 (13)	0.0038 (15)	0.0010 (14)
C23	0.059 (2)	0.061 (2)	0.070 (3)	0.0144 (19)	0.020 (2)	0.0199 (19)
C24	0.096 (3)	0.071 (3)	0.055 (2)	0.030 (3)	-0.005 (2)	0.0129 (18)
C25	0.054 (2)	0.071 (3)	0.086 (3)	0.019 (2)	-0.017 (2)	-0.008 (2)
C26	0.0404 (19)	0.050 (2)	0.096 (3)	0.0027 (16)	0.008 (2)	-0.001 (2)
C27	0.046 (2)	0.0439 (17)	0.065 (2)	0.0104 (14)	0.0076 (16)	0.0028 (15)

Geometric parameters (Å, °)

Cu—N2	2.019 (2)	C10—H10	0.9300
Cu—N1	2.054 (2)	C11—C12	1.389 (5)
Cu—Cl2	2.2141 (9)	C11—H11	0.9300
Cu—Cl1	2.2463 (8)	C12—C13	1.369 (5)
N1—C14	1.498 (4)	C12—H12	0.9300
N1—C1	1.513 (3)	C13—H13	0.9300
N1—C7	1.530 (3)	C14—C15	1.511 (5)
N2—C6	1.482 (4)	C14—H14A	0.9700
N2—C21	1.499 (4)	C14—H14B	0.9700
N2—H2	0.85 (3)	C15—C20	1.383 (5)
C1—C2	1.528 (4)	C15—C16	1.389 (5)
C1—C6	1.533 (4)	C16—C17	1.377 (5)
C1—H1	0.9800	C16—H16	0.9300
C2—C3	1.513 (5)	C17—C18	1.371 (6)
C2—H2A	0.9700	C17—H17	0.9300
C2—H2B	0.9700	C18—C19	1.371 (7)
C3—C4	1.509 (5)	C18—H18	0.9300
C3—H3A	0.9700	C19—C20	1.371 (6)
C3—H3B	0.9700	C19—H19	0.9300
C4—C5	1.528 (5)	C20—H20	0.9300
C4—H4A	0.9700	C21—C22	1.492 (5)
C4—H4B	0.9700	C21—H21A	0.9700
C5—C6	1.531 (4)	C21—H21B	0.9700
C5—H5A	0.9700	C22—C23	1.379 (5)
C5—H5B	0.9700	C22—C27	1.383 (4)
C6—H6	0.9800	C23—C24	1.392 (6)

supplementary materials

C7—C8	1.492 (4)	C23—H23	0.9300
C7—H7A	0.9700	C24—C25	1.366 (6)
C7—H7B	0.9700	C24—H24	0.9300
C8—C9	1.383 (5)	C25—C26	1.352 (6)
C8—C13	1.388 (5)	C25—H25	0.9300
C9—C10	1.368 (5)	C26—C27	1.350 (5)
C9—H9	0.9300	C26—H26	0.9300
C10—C11	1.354 (6)	C27—H27	0.9300
N2—Cu—N1	86.39 (10)	C10—C9—C8	121.2 (4)
N2—Cu—Cl2	156.09 (8)	C10—C9—H9	119.4
N1—Cu—Cl2	96.50 (7)	C8—C9—H9	119.4
N2—Cu—Cl1	89.27 (8)	C11—C10—C9	121.1 (4)
N1—Cu—Cl1	152.80 (7)	C11—C10—H10	119.5
Cl2—Cu—Cl1	98.24 (4)	C9—C10—H10	119.5
C14—N1—C1	115.5 (2)	C10—C11—C12	118.9 (4)
C14—N1—C7	103.3 (2)	C10—C11—H11	120.5
C1—N1—C7	110.3 (2)	C12—C11—H11	120.5
C14—N1—Cu	115.99 (19)	C13—C12—C11	120.4 (4)
C1—N1—Cu	103.32 (16)	C13—C12—H12	119.8
C7—N1—Cu	108.40 (17)	C11—C12—H12	119.8
C6—N2—C21	117.2 (2)	C12—C13—C8	120.9 (3)
C6—N2—Cu	110.95 (18)	C12—C13—H13	119.6
C21—N2—Cu	108.9 (2)	C8—C13—H13	119.6
C6—N2—H2	103 (2)	N1—C14—C15	118.5 (3)
C21—N2—H2	106 (2)	N1—C14—H14A	107.7
Cu—N2—H2	110 (2)	C15—C14—H14A	107.7
N1—C1—C2	116.4 (2)	N1—C14—H14B	107.7
N1—C1—C6	111.0 (2)	C15—C14—H14B	107.7
C2—C1—C6	111.0 (2)	H14A—C14—H14B	107.1
N1—C1—H1	105.9	C20—C15—C16	118.5 (3)
C2—C1—H1	105.9	C20—C15—C14	119.8 (3)
C6—C1—H1	105.9	C16—C15—C14	121.5 (3)
C3—C2—C1	109.4 (3)	C17—C16—C15	120.0 (4)
C3—C2—H2A	109.8	C17—C16—H16	120.0
C1—C2—H2A	109.8	C15—C16—H16	120.0
C3—C2—H2B	109.8	C18—C17—C16	120.4 (4)
C1—C2—H2B	109.8	C18—C17—H17	119.8
H2A—C2—H2B	108.2	C16—C17—H17	119.8
C4—C3—C2	110.5 (3)	C19—C18—C17	120.1 (5)
C4—C3—H3A	109.6	C19—C18—H18	119.9
C2—C3—H3A	109.6	C17—C18—H18	119.9
C4—C3—H3B	109.6	C18—C19—C20	119.7 (5)
C2—C3—H3B	109.6	C18—C19—H19	120.2
H3A—C3—H3B	108.1	C20—C19—H19	120.2
C3—C4—C5	111.1 (3)	C19—C20—C15	121.2 (4)
C3—C4—H4A	109.4	C19—C20—H20	119.4
C5—C4—H4A	109.4	C15—C20—H20	119.4
C3—C4—H4B	109.4	C22—C21—N2	116.0 (3)
C5—C4—H4B	109.4	C22—C21—H21A	108.3

H4A—C4—H4B	108.0	N2—C21—H21A	108.3
C4—C5—C6	111.9 (3)	C22—C21—H21B	108.3
C4—C5—H5A	109.2	N2—C21—H21B	108.3
C6—C5—H5A	109.2	H21A—C21—H21B	107.4
C4—C5—H5B	109.2	C23—C22—C27	116.9 (3)
C6—C5—H5B	109.2	C23—C22—C21	121.8 (3)
H5A—C5—H5B	107.9	C27—C22—C21	121.3 (3)
N2—C6—C5	112.9 (3)	C22—C23—C24	120.5 (4)
N2—C6—C1	108.0 (2)	C22—C23—H23	119.8
C5—C6—C1	109.3 (2)	C24—C23—H23	119.8
N2—C6—H6	108.8	C25—C24—C23	119.6 (3)
C5—C6—H6	108.8	C25—C24—H24	120.2
C1—C6—H6	108.8	C23—C24—H24	120.2
C8—C7—N1	116.9 (2)	C26—C25—C24	120.6 (4)
C8—C7—H7A	108.1	C26—C25—H25	119.7
N1—C7—H7A	108.1	C24—C25—H25	119.7
C8—C7—H7B	108.1	C27—C26—C25	119.4 (4)
N1—C7—H7B	108.1	C27—C26—H26	120.3
H7A—C7—H7B	107.3	C25—C26—H26	120.3
C9—C8—C13	117.6 (3)	C26—C27—C22	123.0 (3)
C9—C8—C7	121.6 (3)	C26—C27—H27	118.5
C13—C8—C7	120.6 (3)	C22—C27—H27	118.5
N2—Cu—N1—C14	104.1 (2)	C1—N1—C7—C8	−53.7 (3)
Cl2—Cu—N1—C14	−52.04 (19)	Cu—N1—C7—C8	58.8 (3)
Cl1—Cu—N1—C14	−174.54 (15)	N1—C7—C8—C9	−86.8 (3)
N2—Cu—N1—C1	−23.25 (17)	N1—C7—C8—C13	99.2 (3)
Cl2—Cu—N1—C1	−179.41 (15)	C13—C8—C9—C10	−1.4 (5)
Cl1—Cu—N1—C1	58.1 (2)	C7—C8—C9—C10	−175.5 (3)
N2—Cu—N1—C7	−140.32 (18)	C8—C9—C10—C11	2.0 (6)
Cl2—Cu—N1—C7	63.51 (17)	C9—C10—C11—C12	−1.7 (6)
Cl1—Cu—N1—C7	−59.0 (2)	C10—C11—C12—C13	0.8 (6)
N1—Cu—N2—C6	−1.8 (2)	C11—C12—C13—C8	−0.3 (6)
Cl2—Cu—N2—C6	96.1 (3)	C9—C8—C13—C12	0.5 (5)
Cl1—Cu—N2—C6	−154.94 (19)	C7—C8—C13—C12	174.8 (3)
N1—Cu—N2—C21	−132.2 (2)	C1—N1—C14—C15	59.0 (4)
Cl2—Cu—N2—C21	−34.3 (3)	C7—N1—C14—C15	179.6 (3)
Cl1—Cu—N2—C21	74.7 (2)	Cu—N1—C14—C15	−62.0 (3)
C14—N1—C1—C2	45.3 (3)	N1—C14—C15—C20	−114.2 (4)
C7—N1—C1—C2	−71.3 (3)	N1—C14—C15—C16	71.3 (4)
Cu—N1—C1—C2	173.0 (2)	C20—C15—C16—C17	3.5 (5)
C14—N1—C1—C6	−82.8 (3)	C14—C15—C16—C17	178.0 (3)
C7—N1—C1—C6	160.6 (2)	C15—C16—C17—C18	−2.5 (6)
Cu—N1—C1—C6	44.9 (2)	C16—C17—C18—C19	−0.1 (8)
N1—C1—C2—C3	171.8 (3)	C17—C18—C19—C20	1.7 (8)
C6—C1—C2—C3	−60.1 (3)	C18—C19—C20—C15	−0.7 (8)
C1—C2—C3—C4	59.6 (4)	C16—C15—C20—C19	−1.9 (6)
C2—C3—C4—C5	−57.4 (4)	C14—C15—C20—C19	−176.5 (4)
C3—C4—C5—C6	55.2 (4)	C6—N2—C21—C22	57.9 (4)
C21—N2—C6—C5	−86.4 (3)	Cu—N2—C21—C22	−175.2 (3)

supplementary materials

Cu—N2—C6—C5	147.6 (2)	N2—C21—C22—C23	78.9 (4)
C21—N2—C6—C1	152.6 (3)	N2—C21—C22—C27	-100.0 (4)
Cu—N2—C6—C1	26.6 (3)	C27—C22—C23—C24	0.3 (5)
C4—C5—C6—N2	-174.6 (3)	C21—C22—C23—C24	-178.6 (3)
C4—C5—C6—C1	-54.3 (4)	C22—C23—C24—C25	0.4 (6)
N1—C1—C6—N2	-48.7 (3)	C23—C24—C25—C26	-1.8 (6)
C2—C1—C6—N2	-179.7 (2)	C24—C25—C26—C27	2.4 (6)
N1—C1—C6—C5	-171.9 (2)	C25—C26—C27—C22	-1.7 (6)
C2—C1—C6—C5	57.1 (3)	C23—C22—C27—C26	0.3 (5)
C14—N1—C7—C8	-177.7 (3)	C21—C22—C27—C26	179.2 (3)

Fig. 1

