

Bis(1,3-benzothiazole-2-thiolato)[(Z)-methyl 2-(2-amino-1,3-thiazol-4-yl)-2-(methoxyimino)acetate]nickel(II)

Islam Ullah Khan,^{a*} Onur Şahin,^{b*} Shehzada Muhammad Sajid Jillani,^a Shahzad Sharif^a and Orhan Büyükgüngör^b

^aMaterials Chemistry Laboratory, Department of Chemistry, Government College University, Lahore 54000, Pakistan, and ^bDepartment of Physics, Ondokuz Mayıs University, TR-55139 Samsun, Turkey
Correspondence e-mail: iuklodhi@yahoo.com, onurs@omu.edu.tr


Received 14 April 2010; accepted 24 April 2010

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.043; wR factor = 0.131; data-to-parameter ratio = 19.0.

In the title compound, $[\text{Ni}(\text{C}_7\text{H}_4\text{NS}_2)_2(\text{C}_7\text{H}_9\text{N}_3\text{O}_3\text{S})]$, the Ni^{II} ion is in a slightly distorted N_4S_2 octahedral coordination environment. The two benzothiazole-2-thiolate ligands chelate *via* their thiazole N and thiolate S atoms while the methyl 2-(2-aminothiazol-4-yl)-2-(methoxyimino)acetate also acts as a chelate ligand binding through the thiazole and imino N atoms. Intramolecular $\text{N}-\text{H}\cdots\text{N}$, $\text{C}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{O}$ interactions contribute to the molecular conformation. In the crystal structure, intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds produce $R_1^2(6)$ rings and generate chains along the c axis. An extensive one-dimensional supramolecular network of $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds and $\text{C}-\text{H}\cdots\pi$ interactions is responsible for the crystal structure stabilization.

Related literature

For the graph-set analysis of hydrogen-bond patterns, see: Bernstein *et al.* (1995). For related structures, see: Batı *et al.* (2006); Sieroń (2007); Liu & Xu (2004); Sharif *et al.* (2009); Song *et al.* (2005); Tashpulatov *et al.* (1957).



Experimental

Crystal data

$[\text{Ni}(\text{C}_7\text{H}_4\text{NS}_2)_2(\text{C}_7\text{H}_9\text{N}_3\text{O}_3\text{S})]$	$V = 2496.5$ (3) Å ³
$M_r = 606.41$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 17.8387$ (11) Å	$\mu = 1.23$ mm ⁻¹
$b = 7.8701$ (5) Å	$T = 296$ K
$c = 17.9861$ (10) Å	$0.42 \times 0.37 \times 0.34$ mm
$\beta = 98.639$ (2)°	

Data collection

Bruker APEXII CCD area-detector diffractometer	6164 independent reflections
26981 measured reflections	3512 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.056$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.131$	$\Delta\rho_{\text{max}} = 0.39$ e Å ⁻³
$S = 1.03$	$\Delta\rho_{\text{min}} = -0.32$ e Å ⁻³
6164 reflections	
324 parameters	
2 restraints	

Table 1

Selected bond lengths (Å).

N1—Ni1	2.103 (3)	N4—Ni1	2.153 (3)
N2—Ni1	2.108 (2)	S2—Ni1	2.5410 (11)
N3—Ni1	2.042 (3)	S4—Ni1	2.5123 (10)

Table 2

Hydrogen-bond geometry (Å, °).

Cg_1 and Cg_2 are the centroids of the C1–C6 and C8–C13 benzene rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N5—H5A \cdots N2	0.85 (2)	2.25 (3)	3.036 (5)	154 (4)
N5—H5B \cdots O2 ⁱ	0.86 (2)	2.24 (3)	3.025 (5)	150 (4)
N5—H5B \cdots O1 ⁱ	0.86 (2)	2.38 (3)	3.036 (4)	133 (3)
C16—H16 \cdots O3	0.93	2.40	2.898 (4)	114
C21—H21B \cdots N1	0.96	2.41	3.282 (5)	151
C4—H4 \cdots Cg2 ⁱⁱ	0.93	2.93	3.588 (6)	129
C9—H9 \cdots Cg1 ⁱ	0.93	2.99	3.636 (4)	128
C21—H21A \cdots Cg2 ⁱⁱⁱ	0.96	2.76	3.556 (4)	141

Symmetry codes: (i) $x, -y + \frac{1}{2}, z - \frac{1}{2}$; (ii) $-x, -y, -z + 2$; (iii) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997); software used to prepare material for publication: WinGX (Farrugia, 1999).

IUK thanks the Higher Education Commission of Pakistan for financial support under the project 'Strengthening of the Materials Chemistry Laboratory' at GCUL.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ2774).

References

- Batı, H., Yüksektepe, Ç., Çalıskan, N. & Büyükğüngör, O. (2006). *Acta Cryst.* **E62**, m2313–m2315.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Liu, J.-G. & Xu, D.-J. (2004). *Acta Cryst.* **E60**, m403–m405.
- Sharif, S., Khan, I. U., Arshad, M. N., Sheikh, T. A. & Qureshi, M. Z. (2009). *Acta Cryst.* **E65**, o1805.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sieroni, L. (2007). *Acta Cryst.* **E63**, m598–m600.
- Song, R.-F., Liu, J.-H., Qian, H. & Zhao, K.-Y. (2005). *Acta Cryst.* **E61**, m2142–m2144.
- Tashpulatov, Yu., Zvonkova, Z. V. & Zhdanov, G. S. (1957). *Kristallografiya*, **2**, 33.

supplementary materials

Acta Cryst. (2010). E66, m587-m588 [doi:10.1107/S1600536810015072]

**Bis(1,3-benzothiazole-2-thiolato)[(Z)-methyl
(methoxyimino)acetate]nickel(II)**

2-(2-amino-1,3-thiazol-4-yl)-2-

I. U. Khan, O. Sahin, S. M. S. Jillani, S. Sharif and O. Büyükgüngör

Comment

This work was performed to explore the ligand properties of (2Z)-Methyl 2-(2-amino-1,3-thiazol-4-yl)-2-(methoxyimino)ethanoate (Sharif *et al.*, 2009), one of the precursors of *S*-1,3-Benzothiazol-2-yl(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-(methoxyimino)-ethanethioate (MAEM). A study of the ligand behaviour of a second MAEM component 2-mercapto-benzothiazole has also been reported (Tashpulatov *et al.*, 1957). We report here the structure of the title compound, (I), in which hydrogen bonds and C—H \cdots π interactions lead to a one-dimensional supramolecular network.

The molecular structure of (I) and atom-labelling scheme are shown in Fig. 1. The Ni^{II} ion is coordinated by two S atoms (S2 and S4) and four N atoms (N1, N2, N3 and N4). The geometry around the Ni^{II} ion is that of a slightly distorted octahedron. The title compound (I) has got two chelate ring types. In the first of these, atoms N3 and N4 are bonded to Ni1, thus generating five-membered chelate ring (C17/N3/Ni1/N4/C18). The other type, atoms N1, N2, S2 and S4 are bonded to Ni1 to form four-membered chelate rings (N1/Ni1/S2/C7) and (N2/Ni1/S4/C14). The two Ni—S distances are 2.5123 (10) and 2.5410 (11) Å. The Ni—N_{thiazol} distance of 2.042 (3) Å in (I) is almost equal to that in [Ni(NCS)(C₆H₆N₄S₂)(CH₄O)]Cl (Liu and Xu, 2004). The Ni—N_{benzothiazol} distances in (I) are longer than the equivalent Pd—N bond distance (Song *et al.*, 2005), Co—N bond distances (Bati *et al.*, 2006) and Cu—N bond distance (Sieroń, 2007). The N1—C7 and N2—C14 bond lengths are indicative of significant double-bond character. These values are comparable with those found for similar compounds (Bati *et al.*, 2006; Sieroń, 2007). The N4=C18 and C19=O2 bond lengths are 1.281 (4) and 1.194 (4) Å, respectively, and agree with the corresponding distances in *S*-1,3-Benzothiazol-2-yl-(2Z)-2-(2-amino-1,3-thiazol-4-yl)-2-(methoxyimino)-ethanethioate [1.281 (3) and 1.190 (3) Å, respectively; Sharif *et al.*, 2009]. The C15—N3 bond is somewhat shorter than the C17—N3 bond, as a result of pronounced delocalization in the -N=C=N- fragment of the 2-aminothiazole ring. Each benzothiazole ligand is planar; the angles between the mean planes through the five- and six-membered rings of each ligand being 2.18 (14)° and 1.37 (20)°.

The molecules of (I) are linked by intermolecular hydrogen bonding, and we employ graph-set notation (Bernstein *et al.*, 1995) to describe the patterns of hydrogen bonding. Molecules of (I) are linked into sheets by a combination of N—H \cdots O hydrogen bonds (Table 2). Within the selected asymmetric unit, intramolecular N—H \cdots N, C—H \cdots N and C—H \cdots O hydrogen bonds define S(6) motifs (Fig. 1). Amino atom N5 in the reference molecule at (x, y, z) acts as hydrogen-bond donor, via atom H5B, to atom O1 in the molecule at (x, 1/2-y, z-1/2), so forming a C(7) chain running parallel to the [00-1] direction. Similarly, amino atom N5 in the reference molecule at (x, y, z) acts as hydrogen-bond donor, via atom H5B, to atom O2 in the molecule at (x, 1/2-y, z-1/2), so forming a C(8) chain running parallel to the [00-1] direction. The combination of C(7) and C(8) chains generates a chain of edge-fused R₁²(6) rings running parallel to the [001] direction (Fig. 2).

Compound (I) also contains three intermolecular C—H \cdots π interactions. In the first, atom C21 in the molecule at (x, y, z) acts as hydrogen-bond donor to the C8—C13 benzene ring in the molecule at (x, 1/2-y, 1/2+z), so forming a C(8) chain running parallel to the [001] direction. The combination of N—H \cdots O hydrogen bonds and C—H \cdots π interactions produce

$R_2^2(11)$ rings (Fig. 2). In the second, atom C4 in the molecule at (x, y, z) acts as hydrogen-bond donor to the C8—C13 benzene ring in the molecule at $(-x, -y, 2-z)$, so forming a centrosymmetric $R_2^2(20)$ rings centred at $(0, 0, n+1)$ ($n = \text{zero or integer}$) (Fig. 3). Finally, atom C9 in the molecule at (x, y, z) acts as hydrogen-bond donor to the C1—C6 benzene ring in the molecule at $(x, 1/2-y, z-1/2)$, so forming a C(9) chain running parallel to the $[00-1]$ direction. Details of these interactions are given in Table 2. The combination of C—H $\cdots\pi$ interactions generates a chain of edge-fused $R_2^2(20)$ and $R_6^6(30)$ rings running parallel to the $[001]$ direction (Fig. 3).

Experimental

MAEM (0.25 g, 0.71 mmol) was dissolved in 20 ml methanol and refluxed for 10 minutes. A solution of nickel acetate (0.25 g, 0.18 mmol) previously prepared in 5 ml methanol was added dropwise, the mixture was refluxed further for one hour. The resulting solution was filtered. The filtrate was kept for slow evaporation, after three days light yellow crystals suitable for crystallography were obtained.

Refinement

All H-atoms bound to C were refined using a riding model with $d(\text{C—H}) = 0.93 \text{ \AA}$ ($U_{\text{iso}} = 1.2U_{\text{eq}}$ of the parent atom) for aromatic carbon atoms and $d(\text{C—H}) = 0.96 \text{ \AA}$ ($U_{\text{iso}} = 1.5U_{\text{eq}}$ of the parent atom) for methyl carbon atoms. Amino H atoms were located in difference maps and refined subject to the DFIX restraint $\text{N—H} = 0.87 (2) \text{ \AA}$.

Figures




Fig. 1. A view of the molecule of (I), showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level and H atoms are shown as small spheres of arbitrary radii. Hydrogen bonds are indicated by dashed lines.




Fig. 2. Part of the crystal structure of (I), showing the formation of $R_1^2(6)$ and $R_2^2(11)$ rings. N—H \cdots O hydrogen bonds and C—H $\cdots\pi$ interactions are indicated by dashed lines. H atoms not involved in these interactions have been omitted for clarity. (Symmetry codes as in Table 2.)




Fig. 3. Part of the crystal structure of (I), showing the formation of a chain along $[001]$ generated by the C—H $\cdots\pi$ interactions (dashed lines; see Table 2). For the sake of clarity, H atoms not involved in the motif shown have been omitted. (Symmetry codes as in Table 2.)

Bis(1,3-benzothiazole-2-thiolato)[(Z)-methyl 2-(2-amino-1,3-thiazol-4-yl)-2-(methoxyimino)acetate]nickel(II)

Crystal data

[Ni(C ₇ H ₄ NS ₂) ₂ (C ₇ H ₉ N ₃ O ₃ S)]	$F(000) = 1240$
$M_r = 606.41$	$D_x = 1.613 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 4277 reflections
$a = 17.8387 (11) \text{ \AA}$	$\theta = 2.3\text{--}23.3^\circ$
$b = 7.8701 (5) \text{ \AA}$	$\mu = 1.23 \text{ mm}^{-1}$
$c = 17.9861 (10) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 98.639 (2)^\circ$	Needle, colourless
$V = 2496.5 (3) \text{ \AA}^3$	$0.42 \times 0.37 \times 0.34 \text{ mm}$
$Z = 4$	

Data collection

Bruker APEXII CCD area-detector diffractometer	3512 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.056$
graphite	$\theta_{\text{max}} = 28.3^\circ$, $\theta_{\text{min}} = 1.2^\circ$
φ and ω scans	$h = -23 \rightarrow 23$
26981 measured reflections	$k = -10 \rightarrow 10$
6164 independent reflections	$l = -20 \rightarrow 23$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.043$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.131$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.03$	$w = 1/[\sigma^2(F_o^2) + (0.0609P)^2]$
6164 reflections	where $P = (F_o^2 + 2F_c^2)/3$
324 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
2 restraints	$\Delta\rho_{\text{max}} = 0.39 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.31 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

supplementary materials

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.1273 (2)	0.1742 (5)	1.06227 (18)	0.0477 (9)
C2	0.0859 (2)	0.3229 (6)	1.0511 (2)	0.0597 (11)
H2	0.1078	0.4225	1.0367	0.072*
C3	0.0109 (3)	0.3189 (8)	1.0620 (3)	0.0794 (15)
H3	-0.0182	0.4171	1.0540	0.095*
C4	-0.0216 (3)	0.1725 (9)	1.0845 (3)	0.0903 (18)
H4	-0.0722	0.1741	1.0916	0.108*
C5	0.0185 (3)	0.0264 (8)	1.0963 (3)	0.0841 (16)
H5	-0.0040	-0.0717	1.1115	0.101*
C6	0.0942 (2)	0.0261 (6)	1.0853 (2)	0.0607 (11)
C7	0.2253 (2)	-0.0043 (5)	1.06449 (18)	0.0443 (9)
C8	0.1283 (2)	0.1743 (5)	0.80170 (18)	0.0440 (9)
C9	0.0844 (2)	0.0802 (6)	0.7465 (2)	0.0593 (11)
H9	0.0446	0.1301	0.7146	0.071*
C10	0.1009 (2)	-0.0861 (6)	0.7403 (2)	0.0646 (12)
H10	0.0720	-0.1508	0.7032	0.078*
C11	0.1593 (3)	-0.1627 (5)	0.7872 (2)	0.0602 (11)
H11	0.1693	-0.2774	0.7810	0.072*
C12	0.2033 (2)	-0.0723 (5)	0.8431 (2)	0.0474 (9)
H12	0.2420	-0.1248	0.8753	0.057*
C13	0.18813 (18)	0.0992 (4)	0.84986 (16)	0.0342 (7)
C14	0.20073 (18)	0.3672 (4)	0.89389 (18)	0.0396 (8)
C15	0.4252 (2)	0.2366 (5)	0.90876 (18)	0.0449 (9)
C16	0.5309 (2)	0.2658 (5)	1.01051 (19)	0.0511 (10)
H16	0.5764	0.2740	1.0431	0.061*
C17	0.46199 (18)	0.2752 (4)	1.03122 (17)	0.0395 (8)
C18	0.44057 (19)	0.3070 (5)	1.10545 (17)	0.0400 (8)
C19	0.4991 (2)	0.3044 (5)	1.17575 (19)	0.0480 (10)
C20	0.6241 (2)	0.3713 (7)	1.2288 (2)	0.0851 (16)
H20A	0.6677	0.4288	1.2156	0.128*
H20B	0.6367	0.2549	1.2407	0.128*
H20C	0.6085	0.4257	1.2718	0.128*
C21	0.2788 (2)	0.4476 (6)	1.1692 (2)	0.0766 (14)
H21A	0.2668	0.4729	1.2183	0.115*
H21B	0.2407	0.3734	1.1434	0.115*
H21C	0.2801	0.5511	1.1412	0.115*
N1	0.20268 (14)	0.1531 (4)	1.05239 (14)	0.0379 (7)
N2	0.22816 (14)	0.2105 (3)	0.90121 (14)	0.0341 (6)
N3	0.40088 (14)	0.2583 (4)	0.97356 (14)	0.0390 (7)
N4	0.36948 (16)	0.3292 (4)	1.10589 (14)	0.0421 (7)

N5	0.3796 (2)	0.2109 (6)	0.84349 (18)	0.0689 (11)
H5A	0.3330 (13)	0.195 (6)	0.846 (3)	0.099 (18)*
H5B	0.400 (2)	0.197 (5)	0.8034 (15)	0.075 (14)*
O1	0.35067 (14)	0.3667 (4)	1.17628 (12)	0.0565 (7)
O2	0.48854 (16)	0.2341 (4)	1.23206 (15)	0.0793 (10)
O3	0.56225 (14)	0.3783 (4)	1.16549 (13)	0.0603 (7)
S1	0.15776 (7)	-0.14012 (16)	1.09195 (7)	0.0752 (4)
S2	0.31353 (5)	-0.06091 (13)	1.04963 (6)	0.0561 (3)
S3	0.12298 (6)	0.38890 (12)	0.82287 (6)	0.0565 (3)
S4	0.24041 (6)	0.52105 (12)	0.95328 (5)	0.0520 (3)
S5	0.52295 (5)	0.23624 (15)	0.91477 (5)	0.0567 (3)
Ni1	0.29632 (2)	0.24588 (5)	1.00641 (2)	0.03629 (14)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.035 (2)	0.077 (3)	0.0312 (18)	0.0005 (19)	0.0061 (15)	0.0008 (18)
C2	0.038 (2)	0.086 (3)	0.055 (2)	0.015 (2)	0.0075 (18)	-0.003 (2)
C3	0.042 (3)	0.134 (5)	0.065 (3)	0.022 (3)	0.018 (2)	-0.010 (3)
C4	0.035 (3)	0.180 (6)	0.059 (3)	0.001 (3)	0.022 (2)	-0.014 (3)
C5	0.049 (3)	0.143 (5)	0.067 (3)	-0.024 (3)	0.028 (2)	0.000 (3)
C6	0.045 (2)	0.095 (3)	0.045 (2)	-0.013 (2)	0.0139 (18)	0.002 (2)
C7	0.041 (2)	0.057 (2)	0.0363 (18)	-0.0005 (17)	0.0095 (15)	0.0108 (16)
C8	0.043 (2)	0.052 (2)	0.0368 (19)	-0.0017 (17)	0.0051 (16)	0.0007 (16)
C9	0.051 (2)	0.079 (3)	0.045 (2)	-0.006 (2)	-0.0011 (18)	-0.004 (2)
C10	0.069 (3)	0.074 (3)	0.051 (2)	-0.021 (2)	0.011 (2)	-0.019 (2)
C11	0.080 (3)	0.045 (2)	0.061 (3)	-0.016 (2)	0.031 (2)	-0.009 (2)
C12	0.055 (2)	0.043 (2)	0.047 (2)	-0.0021 (18)	0.0171 (18)	0.0041 (17)
C13	0.0340 (18)	0.042 (2)	0.0288 (16)	-0.0045 (15)	0.0114 (13)	0.0041 (14)
C14	0.0390 (19)	0.042 (2)	0.0379 (18)	0.0000 (16)	0.0066 (14)	0.0059 (15)
C15	0.0353 (19)	0.070 (3)	0.0310 (17)	0.0007 (17)	0.0116 (14)	0.0010 (17)
C16	0.035 (2)	0.084 (3)	0.0359 (18)	0.0064 (19)	0.0094 (15)	0.0042 (18)
C17	0.0318 (18)	0.058 (2)	0.0298 (16)	0.0000 (15)	0.0078 (13)	0.0029 (15)
C18	0.0311 (18)	0.063 (2)	0.0272 (16)	0.0015 (16)	0.0076 (13)	0.0020 (15)
C19	0.035 (2)	0.081 (3)	0.0286 (19)	0.0021 (18)	0.0049 (15)	0.0021 (17)
C20	0.041 (2)	0.140 (5)	0.067 (3)	-0.006 (3)	-0.019 (2)	0.011 (3)
C21	0.047 (3)	0.131 (4)	0.054 (2)	0.021 (3)	0.014 (2)	-0.028 (3)
N1	0.0304 (15)	0.0493 (18)	0.0359 (15)	0.0019 (13)	0.0111 (11)	0.0052 (13)
N2	0.0316 (15)	0.0374 (16)	0.0335 (14)	-0.0010 (12)	0.0052 (11)	0.0032 (12)
N3	0.0300 (15)	0.062 (2)	0.0258 (13)	-0.0044 (13)	0.0061 (11)	-0.0009 (13)
N4	0.0345 (16)	0.067 (2)	0.0276 (14)	0.0020 (14)	0.0120 (12)	-0.0021 (13)
N5	0.043 (2)	0.136 (4)	0.0301 (17)	-0.014 (2)	0.0131 (15)	-0.0117 (19)
O1	0.0418 (15)	0.102 (2)	0.0273 (12)	0.0106 (14)	0.0117 (10)	-0.0078 (13)
O2	0.0513 (18)	0.146 (3)	0.0385 (16)	-0.0071 (17)	0.0009 (13)	0.0277 (17)
O3	0.0345 (15)	0.099 (2)	0.0440 (14)	-0.0096 (14)	-0.0039 (11)	0.0060 (14)
S1	0.0697 (8)	0.0732 (8)	0.0864 (8)	-0.0135 (6)	0.0235 (6)	0.0291 (7)
S2	0.0450 (6)	0.0578 (6)	0.0653 (6)	0.0135 (5)	0.0076 (5)	0.0124 (5)
S3	0.0585 (7)	0.0500 (6)	0.0549 (6)	0.0118 (5)	-0.0118 (5)	0.0046 (5)

supplementary materials

S4	0.0647 (7)	0.0398 (5)	0.0495 (5)	-0.0027 (5)	0.0023 (5)	-0.0028 (4)
S5	0.0347 (5)	0.1025 (9)	0.0359 (5)	0.0064 (5)	0.0150 (4)	0.0007 (5)
Ni1	0.0291 (2)	0.0506 (3)	0.0298 (2)	0.0005 (2)	0.00674 (16)	0.00160 (19)

Geometric parameters (Å, °)

C1—C2	1.382 (6)	C15—N3	1.315 (4)
C1—N1	1.392 (4)	C15—N5	1.340 (4)
C1—C6	1.398 (5)	C15—S5	1.730 (4)
C2—C3	1.380 (6)	C16—C17	1.339 (5)
C2—H2	0.9300	C16—S5	1.722 (4)
C3—C4	1.378 (8)	C16—H16	0.9300
C3—H3	0.9300	C17—N3	1.394 (4)
C4—C5	1.354 (7)	C17—C18	1.464 (5)
C4—H4	0.9300	C18—N4	1.281 (4)
C5—C6	1.394 (6)	C18—C19	1.514 (4)
C5—H5	0.9300	C19—O2	1.194 (4)
C6—S1	1.724 (5)	C19—O3	1.305 (4)
C7—N1	1.311 (4)	C20—O3	1.463 (4)
C7—S2	1.695 (4)	C20—H20A	0.9600
C7—S1	1.737 (4)	C20—H20B	0.9600
C8—C9	1.383 (5)	C20—H20C	0.9600
C8—C13	1.401 (4)	C21—O1	1.421 (4)
C8—S3	1.737 (4)	C21—H21A	0.9600
C9—C10	1.350 (6)	C21—H21B	0.9600
C9—H9	0.9300	C21—H21C	0.9600
C10—C11	1.377 (6)	N1—Ni1	2.103 (3)
C10—H10	0.9300	N2—Ni1	2.108 (2)
C11—C12	1.376 (5)	N3—Ni1	2.042 (3)
C11—H11	0.9300	N4—O1	1.389 (3)
C12—C13	1.385 (5)	N4—Ni1	2.153 (3)
C12—H12	0.9300	N5—H5A	0.848 (19)
C13—N2	1.390 (4)	N5—H5B	0.863 (19)
C14—N2	1.326 (4)	S2—Ni1	2.5410 (11)
C14—S4	1.697 (3)	S4—Ni1	2.5123 (10)
C14—S3	1.747 (3)		
C2—C1—N1	126.0 (4)	C17—C18—C19	121.1 (3)
C2—C1—C6	120.5 (4)	O2—C19—O3	125.2 (3)
N1—C1—C6	113.4 (4)	O2—C19—C18	122.5 (3)
C3—C2—C1	117.9 (5)	O3—C19—C18	112.2 (3)
C3—C2—H2	121.0	O3—C20—H20A	109.5
C1—C2—H2	121.0	O3—C20—H20B	109.5
C4—C3—C2	121.4 (5)	H20A—C20—H20B	109.5
C4—C3—H3	119.3	O3—C20—H20C	109.5
C2—C3—H3	119.3	H20A—C20—H20C	109.5
C5—C4—C3	121.3 (5)	H20B—C20—H20C	109.5
C5—C4—H4	119.3	O1—C21—H21A	109.5
C3—C4—H4	119.3	O1—C21—H21B	109.5
C4—C5—C6	118.7 (5)	H21A—C21—H21B	109.5

C4—C5—H5	120.7	O1—C21—H21C	109.5
C6—C5—H5	120.7	H21A—C21—H21C	109.5
C5—C6—C1	120.2 (4)	H21B—C21—H21C	109.5
C5—C6—S1	129.3 (4)	C7—N1—C1	111.8 (3)
C1—C6—S1	110.5 (3)	C7—N1—Ni1	98.7 (2)
N1—C7—S2	119.5 (3)	C1—N1—Ni1	148.0 (2)
N1—C7—S1	114.8 (3)	C14—N2—C13	112.0 (3)
S2—C7—S1	125.7 (2)	C14—N2—Ni1	96.95 (19)
C9—C8—C13	120.9 (4)	C13—N2—Ni1	148.1 (2)
C9—C8—S3	129.4 (3)	C15—N3—C17	110.3 (3)
C13—C8—S3	109.6 (2)	C15—N3—Ni1	133.5 (2)
C10—C9—C8	118.2 (4)	C17—N3—Ni1	115.8 (2)
C10—C9—H9	120.9	C18—N4—O1	114.2 (3)
C8—C9—H9	120.9	C18—N4—Ni1	115.5 (2)
C9—C10—C11	121.8 (4)	O1—N4—Ni1	128.5 (2)
C9—C10—H10	119.1	C15—N5—H5A	116 (3)
C11—C10—H10	119.1	C15—N5—H5B	118 (3)
C12—C11—C10	121.1 (4)	H5A—N5—H5B	125 (4)
C12—C11—H11	119.4	N4—O1—C21	110.6 (2)
C10—C11—H11	119.4	C19—O3—C20	116.0 (3)
C11—C12—C13	118.1 (3)	C6—S1—C7	89.47 (19)
C11—C12—H12	121.0	C7—S2—Ni1	74.11 (13)
C13—C12—H12	121.0	C8—S3—C14	90.12 (16)
C12—C13—N2	125.8 (3)	C14—S4—Ni1	74.26 (11)
C12—C13—C8	119.8 (3)	C16—S5—C15	89.61 (17)
N2—C13—C8	114.4 (3)	N3—Ni1—N1	160.83 (11)
N2—C14—S4	119.2 (2)	N3—Ni1—N2	100.11 (10)
N2—C14—S3	113.8 (2)	N1—Ni1—N2	85.53 (10)
S4—C14—S3	126.9 (2)	N3—Ni1—N4	76.10 (10)
N3—C15—N5	123.9 (3)	N1—Ni1—N4	101.37 (10)
N3—C15—S5	114.1 (2)	N2—Ni1—N4	169.15 (11)
N5—C15—S5	121.9 (3)	N3—Ni1—S4	100.16 (8)
C17—C16—S5	110.2 (3)	N1—Ni1—S4	98.95 (8)
C17—C16—H16	124.9	N2—Ni1—S4	68.27 (7)
S5—C16—H16	124.9	N4—Ni1—S4	102.12 (8)
C16—C17—N3	115.8 (3)	N3—Ni1—S2	93.47 (8)
C16—C17—C18	129.7 (3)	N1—Ni1—S2	67.44 (8)
N3—C17—C18	114.4 (3)	N2—Ni1—S2	100.15 (7)
N4—C18—C17	115.0 (3)	N4—Ni1—S2	90.28 (9)
N4—C18—C19	123.8 (3)	S4—Ni1—S2	163.39 (4)
N1—C1—C2—C3	178.5 (3)	C1—C6—S1—C7	0.4 (3)
C6—C1—C2—C3	-1.3 (5)	N1—C7—S1—C6	0.9 (3)
C1—C2—C3—C4	1.0 (6)	S2—C7—S1—C6	-176.7 (3)
C2—C3—C4—C5	-0.4 (7)	N1—C7—S2—Ni1	-5.0 (2)
C3—C4—C5—C6	0.0 (7)	S1—C7—S2—Ni1	172.4 (2)
C4—C5—C6—C1	-0.2 (6)	C9—C8—S3—C14	-178.4 (4)
C4—C5—C6—S1	-177.0 (4)	C13—C8—S3—C14	0.4 (3)
C2—C1—C6—C5	0.9 (6)	N2—C14—S3—C8	-0.3 (3)
N1—C1—C6—C5	-178.9 (3)	S4—C14—S3—C8	-178.2 (3)

supplementary materials

C2—C1—C6—S1	178.2 (3)	N2—C14—S4—Ni1	-10.3 (2)
N1—C1—C6—S1	-1.6 (4)	S3—C14—S4—Ni1	167.6 (3)
C13—C8—C9—C10	-0.1 (6)	C17—C16—S5—C15	-0.5 (3)
S3—C8—C9—C10	178.6 (3)	N3—C15—S5—C16	0.4 (3)
C8—C9—C10—C11	0.3 (6)	N5—C15—S5—C16	-177.5 (4)
C9—C10—C11—C12	0.4 (6)	C15—N3—Ni1—N1	100.0 (4)
C10—C11—C12—C13	-1.4 (6)	C17—N3—Ni1—N1	-71.8 (4)
C11—C12—C13—N2	-177.8 (3)	C15—N3—Ni1—N2	-5.8 (4)
C11—C12—C13—C8	1.6 (5)	C17—N3—Ni1—N2	-177.5 (2)
C9—C8—C13—C12	-0.9 (5)	C15—N3—Ni1—N4	-175.4 (4)
S3—C8—C13—C12	-179.8 (3)	C17—N3—Ni1—N4	12.9 (2)
C9—C8—C13—N2	178.5 (3)	C15—N3—Ni1—S4	-75.3 (3)
S3—C8—C13—N2	-0.4 (4)	C17—N3—Ni1—S4	113.0 (2)
S5—C16—C17—N3	0.5 (4)	C15—N3—Ni1—S2	95.2 (3)
S5—C16—C17—C18	-176.9 (3)	C17—N3—Ni1—S2	-76.6 (2)
C16—C17—C18—N4	171.8 (4)	C7—N1—Ni1—N3	-8.9 (4)
N3—C17—C18—N4	-5.7 (5)	C1—N1—Ni1—N3	-170.9 (4)
C16—C17—C18—C19	-11.0 (6)	C7—N1—Ni1—N2	99.2 (2)
N3—C17—C18—C19	171.5 (3)	C1—N1—Ni1—N2	-62.8 (4)
N4—C18—C19—O2	41.7 (6)	C7—N1—Ni1—N4	-89.2 (2)
C17—C18—C19—O2	-135.2 (4)	C1—N1—Ni1—N4	108.7 (4)
N4—C18—C19—O3	-141.9 (4)	C7—N1—Ni1—S4	166.38 (19)
C17—C18—C19—O3	41.1 (5)	C1—N1—Ni1—S4	4.3 (4)
S2—C7—N1—C1	175.8 (2)	C7—N1—Ni1—S2	-3.72 (18)
S1—C7—N1—C1	-1.9 (4)	C1—N1—Ni1—S2	-165.8 (4)
S2—C7—N1—Ni1	5.9 (3)	C14—N2—Ni1—N3	-104.5 (2)
S1—C7—N1—Ni1	-171.79 (17)	C13—N2—Ni1—N3	99.9 (4)
C2—C1—N1—C7	-177.5 (3)	C14—N2—Ni1—N1	94.0 (2)
C6—C1—N1—C7	2.3 (4)	C13—N2—Ni1—N1	-61.6 (4)
C2—C1—N1—Ni1	-16.7 (6)	C14—N2—Ni1—N4	-36.0 (6)
C6—C1—N1—Ni1	163.1 (3)	C13—N2—Ni1—N4	168.4 (5)
S4—C14—N2—C13	178.3 (2)	C14—N2—Ni1—S4	-7.52 (18)
S3—C14—N2—C13	0.1 (4)	C13—N2—Ni1—S4	-163.1 (4)
S4—C14—N2—Ni1	11.9 (3)	C14—N2—Ni1—S2	160.08 (19)
S3—C14—N2—Ni1	-166.23 (18)	C13—N2—Ni1—S2	4.5 (4)
C12—C13—N2—C14	179.5 (3)	C18—N4—Ni1—N3	-16.6 (3)
C8—C13—N2—C14	0.1 (4)	O1—N4—Ni1—N3	179.9 (3)
C12—C13—N2—Ni1	-26.7 (6)	C18—N4—Ni1—N1	144.0 (3)
C8—C13—N2—Ni1	153.9 (3)	O1—N4—Ni1—N1	-19.6 (3)
N5—C15—N3—C17	177.6 (4)	C18—N4—Ni1—N2	-87.2 (6)
S5—C15—N3—C17	-0.2 (4)	O1—N4—Ni1—N2	109.2 (6)
N5—C15—N3—Ni1	5.6 (6)	C18—N4—Ni1—S4	-114.2 (3)
S5—C15—N3—Ni1	-172.25 (18)	O1—N4—Ni1—S4	82.2 (3)
C16—C17—N3—C15	-0.2 (5)	C18—N4—Ni1—S2	76.9 (3)
C18—C17—N3—C15	177.6 (3)	O1—N4—Ni1—S2	-86.6 (3)
C16—C17—N3—Ni1	173.4 (3)	C14—S4—Ni1—N3	102.97 (14)
C18—C17—N3—Ni1	-8.8 (4)	C14—S4—Ni1—N1	-75.46 (14)
C17—C18—N4—O1	-177.3 (3)	C14—S4—Ni1—N2	6.05 (15)
C19—C18—N4—O1	5.6 (5)	C14—S4—Ni1—N4	-179.22 (15)

C17—C18—N4—Ni1	16.7 (4)	C14—S4—Ni1—S2	-41.67 (19)
C19—C18—N4—Ni1	-160.4 (3)	C7—S2—Ni1—N3	-178.75 (14)
C18—N4—O1—C21	159.1 (3)	C7—S2—Ni1—N1	2.95 (14)
Ni1—N4—O1—C21	-37.2 (4)	C7—S2—Ni1—N2	-77.83 (14)
O2—C19—O3—C20	0.5 (6)	C7—S2—Ni1—N4	105.16 (14)
C18—C19—O3—C20	-175.8 (4)	C7—S2—Ni1—S4	-33.55 (19)
C5—C6—S1—C7	177.4 (4)		

Hydrogen-bond geometry (Å, °)

Cg1 and Cg2 are the centroids of the C1—C6 and C8—C13 benzene rings, respectively.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N5—H5A...N2	0.85 (2)	2.25 (3)	3.036 (5)	154 (4)
N5—H5B...O2 ⁱ	0.86 (2)	2.24 (3)	3.025 (5)	150 (4)
N5—H5B...O1 ⁱ	0.86 (2)	2.38 (3)	3.036 (4)	133 (3)
C16—H16...O3	0.93	2.40	2.898 (4)	114
C21—H21B...N1	0.96	2.41	3.282 (5)	151
C4—H4...Cg2 ⁱⁱ	0.93	2.93	3.588 (6)	129
C9—H9...Cg1 ⁱ	0.93	2.99	3.636 (4)	128
C21—H21A...Cg2 ⁱⁱⁱ	0.96	2.76	3.556 (4)	141

Symmetry codes: (i) $x, -y+1/2, z-1/2$; (ii) $-x, -y, -z+2$; (iii) $x, -y+1/2, z+1/2$.

Fig. 1




Fig. 2




Fig. 3

