

(E)-4-Chloro-2-[(2-hydroxyphenyl)-iminomethyl]phenol

Naser Eltaher Eltayeb,^a‡ Siang Guan Teoh,^a Suchada Chantrapromma^b§ and Hoong-Kun Fun^{c*}¶

^aSchool of Chemical Sciences, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, ^bCrystal Materials Research Unit, Department of Chemistry, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand, and ^cX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia

Correspondence e-mail: hkfun@usm.my


Received 31 March 2010; accepted 1 April 2010

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.034; wR factor = 0.119; data-to-parameter ratio = 17.4.

The title compound, $\text{C}_{13}\text{H}_{10}\text{ClNO}_2$, exists in a *trans* configuration about the central $\text{C}=\text{N}$ bond. The two benzene rings are almost coplanar, making a dihedral angle of $2.48(10)^\circ$. An intramolecular $\text{O}-\text{H}\cdots\text{N}$ hydrogen bond generates an *S*(6) ring motif. In the crystal structure, $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into chains along [101]. Short $\text{C}\cdots\text{Cl}$ contacts [$3.584(2)$ – $3.646(2)\text{ \AA}$] are observed. A short intramolecular $\text{C}-\text{H}\cdots\text{O}$ contact occurs.

Related literature

For bond-length data, see: Allen *et al.* (1987). For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For background to Schiff bases and their applications, see: Dao *et al.* (2000); Eltayeb & Ahmed (2005a,b); Karthikeyan *et al.* (2006); Sriram *et al.* (2006); Wei & Atwood (1998). For related structures, see: Eltayeb *et al.* (2007a,b); Pu (2008). For the stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986).



‡ On study leave from Department of Chemistry, International University of Africa, Khartoum, Sudan, e-mail: naserataha90@hotmail.com.

§ Additional correspondence author, e-mail: suchada.c@psu.ac.th. Thomson Reuters ResearcherID: A-5085-2009.

¶ Thomson Reuters ResearcherID: A-3561-2009.

Experimental

Crystal data

$\text{C}_{13}\text{H}_{10}\text{ClNO}_2$	$V = 536.63(17)\text{ \AA}^3$
$M_r = 247.67$	$Z = 2$
Monoclinic, $P2_1$	Mo $K\alpha$ radiation
$a = 4.6681(9)\text{ \AA}$	$\mu = 0.34\text{ mm}^{-1}$
$b = 18.509(3)\text{ \AA}$	$T = 100\text{ K}$
$c = 6.2118(11)\text{ \AA}$	$0.55 \times 0.14 \times 0.07\text{ mm}$
$\beta = 90.980(4)^\circ$	

Data collection

Bruker APEX Duo CCD area detector diffractometer	4593 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2009)	2680 independent reflections
$T_{\min} = 0.834$, $T_{\max} = 0.976$	2569 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$	H-atom parameters constrained
$wR(F^2) = 0.119$	$\Delta\rho_{\text{max}} = 0.47\text{ e \AA}^{-3}$
$S = 1.20$	$\Delta\rho_{\text{min}} = -0.47\text{ e \AA}^{-3}$
2680 reflections	Absolute structure: Flack (1983),
154 parameters	1125 Friedel pairs
1 restraint	Flack parameter: $-0.03(7)$

H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.47\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.47\text{ e \AA}^{-3}$
 Absolute structure: Flack (1983),
 1125 Friedel pairs
 Flack parameter: $-0.03(7)$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots\text{A}$	$D-\text{H}$	$\text{H}\cdots\text{A}$	$D\cdots\text{A}$	$D-\text{H}\cdots\text{A}$
$\text{O}1-\text{H1O}1\cdots\text{O}2^{\dagger}$	0.82	1.74	2.553 (2)	169
$\text{O}2-\text{H1O}2\cdots\text{N}1$	0.82	1.86	2.602 (2)	149
$\text{C}7-\text{H7}\cdots\text{O}1$	0.93	2.16	2.789 (3)	124

Symmetry code: (i) $x + 1, y, z + 1$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

The authors thank the Malaysian Government, the Ministry of Science, Technology and Innovation (MOSTI) and Universiti Sains Malaysia for the RU research grants (PKIMIA/815002, and PKIMIA/811120). NEE would like to acknowledge Universiti Sains Malaysia for a post-doctoral fellowship and the International University of Africa (Sudan) for providing study leave. The authors also thank the Malaysian Government and Universiti Sains Malaysia for the Research University Golden Goose grant No. 1001/PFIZIK/811012.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2534).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2009). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.

- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
- Dao, V.-T., Gaspard, C., Mayer, M., Werner, G. H., Nguyen, S. N. & Michelot, R. J. (2000). *Eur. J. Med. Chem.* **35**, 805–813.
- Eltayeb, N. E. & Ahmed, T. A. (2005a). *J. Sci. Technol.* **6**, 51–59.
- Eltayeb, N. E. & Ahmed, T. A. (2005b). *Sudan J. Basic Sci.* **7**, 97–108.
- Eltayeb, N. E., Teoh, S. G., Chantrapromma, S., Fun, H.-K. & Ibrahim, K. (2007a). *Acta Cryst.* **E63**, o3094–o3095.
- Eltayeb, N. E., Teoh, S. G., Chantrapromma, S., Fun, H.-K. & Ibrahim, K. (2007b). *Acta Cryst.* **E63**, m2269–m2270.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Karthikeyan, M. S., Prasad, D. J., Poojary, B., Bhat, K. S., Holla, B. S. & Kumari, N. S. (2006). *Bioorg. Med. Chem.* **14**, 7482–7489.
- Pu, X.-H. (2008). *Acta Cryst.* **E64**, o1734.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Sriram, D., Yogeeswari, P., Myneedu, N. S. & Saraswat, V. (2006). *Bioorg. Med. Chem. Lett.* **16**, 2127–2129.
- Wei, P. & Atwood, D. A. (1998). *Inorg. Chem.* **37**, 4934–4938.

supplementary materials

Acta Cryst. (2010). E66, o1065-o1066 [doi:10.1107/S160053681001233X]

(E)-4-Chloro-2-[(2-hydroxyphenyl)iminomethyl]phenol

N. E. Eltayeb, S. G. Teoh, S. Chantrapromma and H.-K. Fun

Comment

We have been interested in synthesis of Schiff base ligands and their complexes (Eltayeb *et al.*, 2007*a,b*) due to their applications such as analytical reagents for the determination of trace elements (Eltayeb & Ahmed, 2005*a,b*), pharmacological activities, anticancer (Dao *et al.*, 2000), anti-HIV (Sriram *et al.*, 2006), antibacterial and antifungal (Karthikeyan *et al.*, 2006) activities. The title compound was used to synthesis the chelated borate catalyst (Wei & Atwood, 1998). Herein we report the crystal structure of the title Schiff base ligand (I).

The molecule of (I) (Fig. 1), C₁₃H₁₀ClNO₂, crystallizes in a *trans* configuration about the C=N bond [1.312 (3) Å] with a torsion angle C1–N1–C7–C8 = -179.2 (2)°. The molecule is almost planar with a dihedral angle between the two benzene rings being 2.48 (10)°. The chloro and two hydroxy groups lie on the same plane with their attached benzene rings with the r.m.s. of 0.0105 (2) Å for the seven non H atoms (C1, C2, C3, C4, C5, C6 and O1) and 0.0129 (2) Å for the eight non H atoms (C8, C9, C10, C11, C12, C13, O2 and Cl1). An intramolecular O—H···N hydrogen bond between the imine N atom and one hydroxy group generates an S(6) ring motif (Fig. 1 and Table 1) (Bernstein *et al.*, 1995). The bond distances are of normal values (Allen *et al.*, 1987) and are comparable with the related structure (Pu, 2008).

In the crystal packing (Fig. 2), O—H···O hydrogen bonds (Table 1) which formed between the two hydroxy O atoms link the molecules into chains along the [101] direction. The crystal is consolidated by O—H···O hydrogen bonds (Table 1) and C···Cl [3.584 (2)–3.646 (2) Å] short contacts.

Experimental

The title compound was synthesized by adding 5-chloro-2-hydroxybenzaldehyde (0.312 g, 2 mmol) to the solution of 2-aminophenol (0.218 g, 2 mmol) in ethanol (30 ml). The mixture was refluxed with stirring for half an hour. The resultant yellow-orange solution was filtered and the filtrate was evaporated to give a yellow solid product. Yellow needle-shaped single crystals of the title compound suitable for x-ray structure determination were obtained from ethanol by slow evaporation at room temperature after a few days.

Refinement

All H atoms were positioned geometrically and allowed to ride on their parent atoms, with $d(\text{O—H}) = 0.82 \text{ \AA}$ and $d(\text{C—H}) = 0.93 \text{ \AA}$. The $U_{\text{iso}}(\text{H})$ values were constrained to be $1.2U_{\text{eq}}$ of the carrier atoms. The highest residual electron density peak is located at 0.70 Å from C5 and the deepest hole is located at 0.05 Å from H1O2.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound, with 50% probability displacement ellipsoids and the atom-numbering scheme. The hydrogen bond is shown as a dashed line.




Fig. 2. The crystal packing of the title compound viewed down the a axis, showing chains running along the [101] direction. Hydrogen bonds are shown as dashed lines.

(E)-4-Chloro-2-[(2-hydroxyphenylimino)methyl]phenol

Crystal data

C ₁₃ H ₁₀ ClNO ₂	$F(000) = 256$
$M_r = 247.67$	$D_x = 1.533 \text{ Mg m}^{-3}$
Monoclinic, $P2_1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: P 2yb	Cell parameters from 2680 reflections
$a = 4.6681 (9) \text{ \AA}$	$\theta = 4.4\text{--}30.0^\circ$
$b = 18.509 (3) \text{ \AA}$	$\mu = 0.34 \text{ mm}^{-1}$
$c = 6.2118 (11) \text{ \AA}$	$T = 100 \text{ K}$
$\beta = 90.980 (4)^\circ$	Needle, yellow
$V = 536.63 (17) \text{ \AA}^3$	$0.55 \times 0.14 \times 0.07 \text{ mm}$
$Z = 2$	

Data collection

Bruker APEX Duo CCD area detector diffractometer	2680 independent reflections
Radiation source: sealed tube graphite	2569 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.023$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2009)	$\theta_{\text{max}} = 30.0^\circ, \theta_{\text{min}} = 4.4^\circ$
$T_{\text{min}} = 0.834, T_{\text{max}} = 0.976$	$h = -6 \rightarrow 6$
4593 measured reflections	$k = -24 \rightarrow 26$
	$l = -8 \rightarrow 8$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.034$	H-atom parameters constrained

$wR(F^2) = 0.119$	$w = 1/[\sigma^2(F_o^2) + (0.0756P)^2 + 0.0404P]$
	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.20$	$(\Delta/\sigma)_{\max} = 0.001$
2680 reflections	$\Delta\rho_{\max} = 0.47 \text{ e } \text{\AA}^{-3}$
154 parameters	$\Delta\rho_{\min} = -0.47 \text{ e } \text{\AA}^{-3}$
1 restraint	Absolute structure: Flack (1983), 1125 Friedel pairs
Primary atom site location: structure-invariant direct methods	Flack parameter: -0.03 (7)

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.02845 (11)	1.08595 (3)	0.36240 (8)	0.01689 (14)
O1	1.1428 (4)	0.82828 (9)	0.5209 (2)	0.0167 (3)
H1O1	1.2817	0.8379	0.5983	0.025*
O2	0.5318 (4)	0.86922 (9)	-0.2123 (2)	0.0151 (3)
H1O2	0.6550	0.8463	-0.1462	0.018*
N1	0.8674 (4)	0.83076 (10)	0.1062 (3)	0.0116 (3)
C1	1.0806 (4)	0.78021 (11)	0.1679 (3)	0.0113 (4)
C2	1.2187 (4)	0.77923 (11)	0.3724 (3)	0.0119 (4)
C3	1.4270 (5)	0.72625 (12)	0.4139 (3)	0.0144 (4)
H3	1.5136	0.7236	0.5496	0.017*
C4	1.5066 (5)	0.67719 (12)	0.2539 (4)	0.0157 (4)
H4	1.6481	0.6430	0.2828	0.019*
C5	1.3732 (5)	0.67964 (12)	0.0506 (4)	0.0147 (4)
H5	1.4255	0.6471	-0.0559	0.018*
C6	1.1629 (5)	0.73071 (12)	0.0089 (3)	0.0140 (4)
H6	1.0747	0.7323	-0.1264	0.017*
C7	0.7386 (5)	0.87972 (12)	0.2228 (3)	0.0122 (4)
H7	0.7886	0.8845	0.3677	0.015*
C8	0.5237 (5)	0.92579 (11)	0.1321 (3)	0.0114 (4)
C9	0.4240 (5)	0.91833 (11)	-0.0859 (3)	0.0115 (4)
C10	0.2019 (5)	0.96567 (12)	-0.1572 (3)	0.0132 (4)
H10	0.1328	0.9622	-0.2981	0.016*

supplementary materials

C11	0.0865 (5)	1.01690 (12)	-0.0213 (3)	0.0136 (4)
H11	-0.0572	1.0476	-0.0720	0.016*
C12	0.1857 (5)	1.02276 (11)	0.1934 (3)	0.0127 (4)
C13	0.4000 (5)	0.97834 (12)	0.2706 (3)	0.0124 (4)
H13	0.4641	0.9825	0.4126	0.015*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0209 (3)	0.0156 (2)	0.0141 (2)	0.00493 (18)	-0.00206 (15)	-0.0037 (2)
O1	0.0172 (8)	0.0217 (8)	0.0111 (7)	0.0046 (6)	-0.0046 (5)	-0.0049 (6)
O2	0.0166 (8)	0.0178 (8)	0.0108 (7)	0.0039 (5)	-0.0021 (5)	-0.0036 (6)
N1	0.0124 (8)	0.0118 (8)	0.0106 (8)	-0.0001 (6)	-0.0010 (5)	-0.0005 (6)
C1	0.0108 (9)	0.0113 (9)	0.0116 (9)	0.0006 (7)	-0.0013 (6)	0.0001 (7)
C2	0.0120 (9)	0.0139 (9)	0.0098 (9)	-0.0003 (6)	-0.0006 (6)	0.0006 (7)
C3	0.0148 (10)	0.0187 (11)	0.0097 (9)	0.0017 (7)	-0.0025 (6)	0.0017 (8)
C4	0.0143 (10)	0.0143 (10)	0.0183 (11)	0.0024 (7)	-0.0012 (7)	0.0015 (8)
C5	0.0157 (10)	0.0145 (10)	0.0140 (10)	-0.0002 (7)	0.0012 (7)	-0.0013 (8)
C6	0.0152 (10)	0.0154 (10)	0.0113 (10)	-0.0010 (7)	-0.0011 (7)	-0.0025 (7)
C7	0.0117 (9)	0.0138 (9)	0.0111 (9)	0.0001 (6)	0.0000 (6)	0.0000 (7)
C8	0.0120 (9)	0.0128 (9)	0.0092 (9)	0.0002 (6)	-0.0012 (6)	-0.0003 (7)
C9	0.0126 (10)	0.0130 (9)	0.0090 (9)	0.0001 (7)	-0.0002 (6)	0.0004 (7)
C10	0.0155 (10)	0.0157 (9)	0.0083 (9)	0.0009 (7)	-0.0011 (7)	0.0016 (7)
C11	0.0142 (10)	0.0134 (9)	0.0132 (10)	0.0006 (7)	-0.0008 (6)	0.0016 (8)
C12	0.0143 (10)	0.0121 (9)	0.0117 (9)	-0.0007 (7)	0.0017 (6)	-0.0012 (8)
C13	0.0138 (10)	0.0155 (10)	0.0077 (9)	-0.0011 (7)	-0.0001 (6)	-0.0005 (7)

Geometric parameters (\AA , $^\circ$)

Cl1—C12	1.742 (2)	C5—C6	1.384 (3)
O1—C2	1.346 (3)	C5—H5	0.9300
O1—H1O1	0.8200	C6—H6	0.9300
O2—C9	1.308 (3)	C7—C8	1.425 (3)
O2—H1O2	0.8200	C7—H7	0.9300
N1—C7	1.312 (3)	C8—C13	1.427 (3)
N1—C1	1.414 (3)	C8—C9	1.431 (3)
C1—C6	1.405 (3)	C9—C10	1.422 (3)
C1—C2	1.415 (3)	C10—C11	1.385 (3)
C2—C3	1.402 (3)	C10—H10	0.9300
C3—C4	1.401 (3)	C11—C12	1.408 (3)
C3—H3	0.9300	C11—H11	0.9300
C4—C5	1.399 (3)	C12—C13	1.375 (3)
C4—H4	0.9300	C13—H13	0.9300
C2—O1—H1O1	109.5	N1—C7—C8	121.39 (19)
C9—O2—H1O2	109.5	N1—C7—H7	119.3
C7—N1—C1	129.38 (18)	C8—C7—H7	119.3
C6—C1—N1	116.14 (18)	C7—C8—C13	117.29 (18)
C6—C1—C2	119.79 (18)	C7—C8—C9	122.13 (19)

N1—C1—C2	124.01 (19)	C13—C8—C9	120.52 (18)
O1—C2—C3	122.39 (18)	O2—C9—C10	121.83 (18)
O1—C2—C1	119.04 (18)	O2—C9—C8	120.86 (18)
C3—C2—C1	118.56 (19)	C10—C9—C8	117.30 (18)
C4—C3—C2	120.96 (19)	C11—C10—C9	121.41 (18)
C4—C3—H3	119.5	C11—C10—H10	119.3
C2—C3—H3	119.5	C9—C10—H10	119.3
C5—C4—C3	120.0 (2)	C10—C11—C12	120.33 (19)
C5—C4—H4	120.0	C10—C11—H11	119.8
C3—C4—H4	120.0	C12—C11—H11	119.8
C6—C5—C4	119.6 (2)	C13—C12—C11	120.7 (2)
C6—C5—H5	120.2	C13—C12—Cl1	120.17 (16)
C4—C5—H5	120.2	C11—C12—Cl1	119.16 (17)
C5—C6—C1	121.0 (2)	C12—C13—C8	119.77 (19)
C5—C6—H6	119.5	C12—C13—H13	120.1
C1—C6—H6	119.5	C8—C13—H13	120.1
C7—N1—C1—C6	175.5 (2)	N1—C7—C8—C9	4.0 (3)
C7—N1—C1—C2	-7.2 (4)	C7—C8—C9—O2	-1.3 (3)
C6—C1—C2—O1	178.20 (19)	C13—C8—C9—O2	-178.44 (19)
N1—C1—C2—O1	1.0 (3)	C7—C8—C9—C10	177.9 (2)
C6—C1—C2—C3	-2.8 (3)	C13—C8—C9—C10	0.8 (3)
N1—C1—C2—C3	180.0 (2)	O2—C9—C10—C11	179.1 (2)
O1—C2—C3—C4	-178.3 (2)	C8—C9—C10—C11	-0.1 (3)
C1—C2—C3—C4	2.8 (3)	C9—C10—C11—C12	-0.6 (3)
C2—C3—C4—C5	-1.4 (3)	C10—C11—C12—C13	0.6 (3)
C3—C4—C5—C6	0.0 (3)	C10—C11—C12—Cl1	-178.08 (17)
C4—C5—C6—C1	-0.1 (3)	C11—C12—C13—C8	0.1 (3)
N1—C1—C6—C5	178.9 (2)	C11—C12—C13—C8	178.79 (15)
C2—C1—C6—C5	1.5 (3)	C7—C8—C13—C12	-178.1 (2)
C1—N1—C7—C8	-179.2 (2)	C9—C8—C13—C12	-0.8 (3)
N1—C7—C8—C13	-178.77 (19)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1O1···O2 ⁱ	0.82	1.74	2.553 (2)	169
O2—H1O2···N1	0.82	1.86	2.602 (2)	149
C7—H7···O1	0.93	2.16	2.789 (3)	124

Symmetry codes: (i) $x+1, y, z+1$.

supplementary materials

Fig. 1




Fig. 2

