

8,9-Dimethoxy-5-phenylsulfonyl-5H-benzo[*b*]carbazole

T. Kavitha,^a M. Thenmozhi,^a V. Dhayalan,^b A. K. Mohanakrishnan^b and M. N. Ponnuswamy^{a*}

^aCentre of Advanced Study in Crystallography and Biophysics, University of Madras, Guindy Campus, Chennai 600 025, India, and ^bDepartment of Organic Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India

Correspondence e-mail: mnpsy2004@yahoo.com


Received 12 March 2010; accepted 2 April 2010

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.044; wR factor = 0.138; data-to-parameter ratio = 25.7.

In the title compound, $\text{C}_{24}\text{H}_{19}\text{NO}_4\text{S}$, the benzocarbazole ring system is planar (r.m.s. deviation = 0.016 Å) and forms a dihedral angle of 78.54 (4)° with the sulfonyl-bound phenyl ring. Intramolecular $\text{C}-\text{H}\cdots\text{O}$ interactions are observed. A $C(8)$ chain running along the b axis is formed *via* intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds. The chains are linked *via* weak $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For bond-length data, see: Allen *et al.* (1987). For the biological activity of carbazole derivatives, see: Itoigawa *et al.* (2000); Tachibana *et al.* (2001); Ramsewak *et al.* (1999). For related structures, see: Chakkaravarthi *et al.* (2008); Govindasamy *et al.* (1998).



Experimental

Crystal data

$\text{C}_{24}\text{H}_{19}\text{NO}_4\text{S}$	$\gamma = 105.883$ (1)°
$M_r = 417.46$	$V = 984.05$ (4) Å ³
Triclinic, $P\bar{1}$	$Z = 2$
$a = 7.8606$ (2) Å	Mo $K\alpha$ radiation
$b = 9.5892$ (2) Å	$\mu = 0.20$ mm ⁻¹
$c = 13.8846$ (4) Å	$T = 293$ K
$\alpha = 100.387$ (1)°	$0.30 \times 0.25 \times 0.20$ mm
$\beta = 93.168$ (2)°	

Data collection

Bruker Kappa APEXII area-detector diffractometer	26246 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 2001)	7025 independent reflections
$T_{\min} = 0.943$, $T_{\max} = 0.962$	5372 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$	273 parameters
$wR(F^2) = 0.138$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\text{max}} = 0.39$ e Å ⁻³
7025 reflections	$\Delta\rho_{\text{min}} = -0.25$ e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C4}-\text{H4}\cdots\text{O1}$	0.93	2.34	2.9238 (19)	120
$\text{C6}-\text{H6}\cdots\text{O2}$	0.93	2.37	2.9674 (15)	122
$\text{C2}-\text{H2}\cdots\text{O2}^{\text{i}}$	0.93	2.55	3.3361 (17)	143
$\text{C24}-\text{H24C}\cdots\text{Cg1}^{\text{ii}}$	0.96	2.84	3.727 (2)	154
$\text{C25}-\text{H25A}\cdots\text{Cg2}^{\text{iii}}$	0.96	2.90	3.628 (2)	134

Symmetry codes: (i) $x, y-1, z$; (ii) $-x+2, -y+1, -z$; (iii) $-x+2, -y, -z$. Cg1 and Cg2 are the centroids of the $\text{C7}-\text{C10}/\text{C22}/\text{C23}$ and $\text{C1}-\text{C4}/\text{C21}/\text{C20}$ rings, respectively.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997); software used to prepare material for publication: SHELXL97 and PLATON (Spek, 2009).

TK thanks the CSIR, India, for financial support in the form of a senior research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI5059).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chakkaravarthi, G., Dhayalan, V., Mohanakrishnan, A. K. & Manivannan, V. (2008). *Acta Cryst. E64*, o1667–o1668.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Govindasamy, L., Velmurugan, D., Ravikumar, K. & Mohanakrishnan, A. K. (1998). *Acta Cryst. C54*, 277–279.
- Itoigawa, M., Kashiwada, Y., Ito, C., Furukawa, H., Tachibana, Y., Bastow, K. F. & Lee, K. H. (2000). *J. Nat. Prod.* **63**, 893–897.
- Ramsewak, R. S., Nair, M. G., Strasburg, G. M., DeWitt, D. L. & Nitiss, J. L. (1999). *J. Agric. Food Chem.* **47**, 444–447.
- Sheldrick, G. M. (2001). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Tachibana, Y., Kikuzaki, H., Lajis, N. H. & Nakatani, N. (2001). *J. Agric. Food Chem.* **49**, 5589–5594.

supplementary materials

Acta Cryst. (2010). E66, o1071 [doi:10.1107/S160053681001247X]

8,9-Dimethoxy-5-phenylsulfonyl-5*H*-benzo[*b*]carbazole

T. Kavitha, M. Thenmozhi, V. Dhayalan, A. K. Mohanakrishnan and M. N. Ponnuswamy

Comment

Carbazole and its derivatives are considered as potential compounds owing to their applications in pharmacy and molecular electronics. They also possess various biological activities such as antitumor (Itoigawa *et al.*, 2000), antioxidative (Tachibana *et al.*, 2001), anti-inflammatory and antimutagenic (Ramsewak *et al.*, 1999) properties.

The benzocarbazole ring system is planar (r.m.s. deviation 0.016 Å) and it forms a dihedral angle of 78.54 (4)° with the sulfonyl-bound phenyl ring (Fig.1). The N—C bond lengths, namely N5—C19 and N5—C21 [1.432 (1) & 1.436 (2) Å] deviate slightly from the mean value reported in the literature (1.370 (12) Å; Allen *et al.*, 1987). This may be due to the electron-withdrawing character of the phenylsulfonyl group (Govindasamy *et al.*, 1998) substituted at N atom of the carbazole group. The S atom exhibits a distorted tetrahedral geometry. The widening of the O1—S1—O2 [119.60 (6)°] angle may be due to repulsive interactions between the two S=O bonds (Chakkaravarthi *et al.*, 2008). The sum of the bond angles around N1 [350.6°] indicate the sp² hybridization. The methoxy groups substituted at C8 and C9 lie almost in the plane of the attached benzene ring [C7—C8—O3—C24 = -3.8 (2)° and C10—C9—O4—C25 = 6.8 (2)°].

The intermolecular C—H···O hydrogen bonds link the molecules into a C(8) chain running along the *b* axis (Fig.2). The packing of the molecules is further influenced by C—H···π interactions.

Experimental

To a solution of diethyl-2-[(2-bromomethyl-1-phenylsulfonyl-1*H*-indol-3-yl)methylene]malonate (0.41 g, 0.78 mmol) in dry 1,2-dichloroethane (15 ml), anhydrous ZnBr₂ (0.35 g, 1.55 mmol) and veratrole (0.12 ml, 0.94 mmol) were added. The mixture was refluxed for 5 h under N₂ atmosphere. The solvent was removed and the reaction mixture was quenched with ice-water (50 ml) containing 1 ml of conc. HCl, extracted with chloroform (2 × 10 ml) and dried (Na₂SO₄). Removal of the solvent followed by flash column chromatographic purification (n-hexane-ethyl acetate 99:1) led to the isolation of the title compound as a colourless solid. The compound was recrystallized from CDCl₃.

Refinement

H atoms were positioned geometrically (C—H = 0.93–0.96 Å) and allowed to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$ and $1.2U_{\text{eq}}(\text{C})$. A rotating group model was used for the methyl groups.

Figures




Fig. 1. The molecular structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level. H atoms have been omitted for clarity.




Fig. 2. Crystal packing of the title compound. H atoms not involved in hydrogen bonding (dashed lines) have been omitted for clarity.

8,9-Dimethoxy-5-phenylsulfonyl-5H-benzo[b]carbazole

Crystal data

$C_{24}H_{19}NO_4S$

$M_r = 417.46$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 7.8606$ (2) Å

$b = 9.5892$ (2) Å

$c = 13.8846$ (4) Å

$\alpha = 100.387$ (1)°

$\beta = 93.168$ (2)°

$\gamma = 105.883$ (1)°

$V = 984.05$ (4) Å³

$Z = 2$

$F(000) = 436$

$D_x = 1.409$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 7025 reflections

$\theta = 1.5$ – 33.3 °

$\mu = 0.20$ mm⁻¹

$T = 293$ K

Block, colourless

$0.30 \times 0.25 \times 0.20$ mm

Data collection

Bruker Kappa APEXII area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

ω and ϕ scans

Absorption correction: multi-scan (SADABS; Sheldrick, 2001)

$T_{\min} = 0.943$, $T_{\max} = 0.962$

26246 measured reflections

7025 independent reflections

5372 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.024$

$\theta_{\max} = 33.3$ °, $\theta_{\min} = 1.5$ °

$h = -12 \rightarrow 11$

$k = -14 \rightarrow 13$

$l = -21 \rightarrow 21$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.044$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.138$	H-atom parameters constrained
$S = 1.04$	$w = 1/[\sigma^2(F_o^2) + (0.0749P)^2 + 0.1333P]$
7025 reflections	where $P = (F_o^2 + 2F_c^2)/3$
273 parameters	$(\Delta/\sigma)_{\max} = 0.001$
0 restraints	$\Delta\rho_{\max} = 0.39 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.62768 (4)	0.24308 (3)	0.35159 (2)	0.03860 (9)
O1	0.54424 (14)	0.18104 (12)	0.42890 (8)	0.0543 (3)
O2	0.57390 (13)	0.35798 (11)	0.31820 (7)	0.0483 (2)
O3	0.84578 (17)	0.47850 (12)	-0.16011 (8)	0.0580 (3)
O4	0.89925 (15)	0.24214 (13)	-0.25315 (7)	0.0566 (3)
C1	0.6666 (2)	-0.24922 (15)	0.16579 (12)	0.0538 (3)
H1	0.7018	-0.2941	0.1083	0.065*
C2	0.6256 (2)	-0.32426 (16)	0.24138 (14)	0.0600 (4)
H2	0.6350	-0.4200	0.2351	0.072*
C3	0.5710 (2)	-0.25904 (16)	0.32592 (13)	0.0543 (4)
H3	0.5440	-0.3121	0.3757	0.065*
C4	0.55495 (18)	-0.11624 (15)	0.33915 (11)	0.0484 (3)
H4	0.5166	-0.0732	0.3962	0.058*
N5	0.59104 (14)	0.10465 (11)	0.25487 (8)	0.0387 (2)
C6	0.67353 (17)	0.25195 (13)	0.12110 (9)	0.0386 (2)
H6	0.6486	0.3362	0.1542	0.046*
C7	0.76027 (18)	0.37119 (14)	-0.01873 (9)	0.0421 (3)
H7	0.7390	0.4572	0.0142	0.051*

supplementary materials

C8	0.81631 (18)	0.36643 (15)	-0.11016 (9)	0.0430 (3)
C9	0.84830 (17)	0.23465 (15)	-0.16184 (9)	0.0432 (3)
C10	0.82608 (18)	0.11485 (15)	-0.11883 (9)	0.0442 (3)
H10	0.8490	0.0300	-0.1525	0.053*
C11	0.74574 (18)	-0.00587 (13)	0.02134 (10)	0.0424 (3)
H11	0.7687	-0.0913	-0.0112	0.051*
C12	0.85883 (16)	0.30130 (14)	0.38223 (8)	0.0386 (2)
C13	0.9589 (2)	0.42875 (18)	0.35613 (11)	0.0530 (3)
H13	0.9045	0.4838	0.3228	0.064*
C14	1.1410 (2)	0.4730 (2)	0.38032 (12)	0.0677 (5)
H14	1.2102	0.5592	0.3641	0.081*
C15	1.2202 (2)	0.3895 (2)	0.42836 (13)	0.0687 (5)
H15	1.3429	0.4197	0.4445	0.082*
C16	1.1192 (3)	0.2617 (2)	0.45275 (15)	0.0726 (5)
H16	1.1743	0.2055	0.4845	0.087*
C17	0.9365 (2)	0.21628 (18)	0.43036 (12)	0.0575 (4)
H17	0.8674	0.1306	0.4473	0.069*
C18	0.68934 (16)	0.00057 (12)	0.11331 (9)	0.0379 (2)
C19	0.65257 (15)	0.12981 (12)	0.16256 (8)	0.0348 (2)
C20	0.65429 (16)	-0.10571 (13)	0.17721 (10)	0.0406 (2)
C21	0.59885 (15)	-0.04054 (13)	0.26328 (10)	0.0396 (2)
C22	0.76869 (16)	0.11715 (13)	-0.02368 (9)	0.0382 (2)
C23	0.73374 (16)	0.24697 (12)	0.02715 (8)	0.0368 (2)
C24	0.8057 (3)	0.60947 (18)	-0.11663 (13)	0.0630 (4)
H24A	0.6826	0.5860	-0.1055	0.094*
H24B	0.8287	0.6784	-0.1600	0.094*
H24C	0.8788	0.6528	-0.0550	0.094*
C25	0.9127 (2)	0.1091 (2)	-0.31191 (12)	0.0639 (4)
H25A	1.0028	0.0769	-0.2805	0.096*
H25B	0.9438	0.1264	-0.3755	0.096*
H25C	0.8005	0.0341	-0.3195	0.096*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.04190 (16)	0.04201 (16)	0.03897 (15)	0.01777 (12)	0.01525 (11)	0.01425 (11)
O1	0.0600 (6)	0.0618 (6)	0.0504 (5)	0.0199 (5)	0.0287 (5)	0.0245 (5)
O2	0.0553 (5)	0.0502 (5)	0.0523 (5)	0.0299 (4)	0.0170 (4)	0.0168 (4)
O3	0.0840 (8)	0.0512 (6)	0.0463 (5)	0.0219 (5)	0.0177 (5)	0.0221 (4)
O4	0.0694 (7)	0.0673 (7)	0.0392 (5)	0.0250 (5)	0.0174 (5)	0.0151 (4)
C1	0.0658 (9)	0.0351 (6)	0.0620 (9)	0.0165 (6)	0.0024 (7)	0.0124 (6)
C2	0.0685 (9)	0.0359 (6)	0.0768 (11)	0.0123 (6)	-0.0037 (8)	0.0224 (7)
C3	0.0503 (7)	0.0453 (7)	0.0676 (9)	0.0041 (6)	-0.0019 (6)	0.0296 (7)
C4	0.0456 (6)	0.0460 (7)	0.0551 (7)	0.0068 (5)	0.0065 (6)	0.0241 (6)
N5	0.0430 (5)	0.0359 (5)	0.0405 (5)	0.0118 (4)	0.0081 (4)	0.0140 (4)
C6	0.0494 (6)	0.0326 (5)	0.0367 (5)	0.0152 (4)	0.0084 (5)	0.0084 (4)
C7	0.0546 (7)	0.0358 (5)	0.0382 (6)	0.0146 (5)	0.0078 (5)	0.0103 (4)
C8	0.0482 (6)	0.0444 (6)	0.0377 (6)	0.0121 (5)	0.0051 (5)	0.0134 (5)

C9	0.0442 (6)	0.0520 (7)	0.0340 (5)	0.0147 (5)	0.0057 (5)	0.0088 (5)
C10	0.0513 (7)	0.0448 (6)	0.0381 (6)	0.0186 (5)	0.0072 (5)	0.0044 (5)
C11	0.0524 (7)	0.0340 (5)	0.0428 (6)	0.0173 (5)	0.0050 (5)	0.0054 (4)
C12	0.0426 (6)	0.0437 (6)	0.0313 (5)	0.0159 (5)	0.0085 (4)	0.0052 (4)
C13	0.0522 (7)	0.0612 (8)	0.0453 (7)	0.0092 (6)	0.0095 (6)	0.0198 (6)
C14	0.0506 (8)	0.0878 (12)	0.0531 (8)	-0.0009 (8)	0.0129 (7)	0.0152 (8)
C15	0.0437 (7)	0.0943 (13)	0.0579 (9)	0.0186 (8)	0.0040 (7)	-0.0083 (9)
C16	0.0676 (10)	0.0737 (11)	0.0762 (12)	0.0361 (9)	-0.0170 (9)	-0.0025 (9)
C17	0.0611 (8)	0.0498 (8)	0.0610 (9)	0.0184 (6)	-0.0082 (7)	0.0106 (6)
C18	0.0411 (6)	0.0313 (5)	0.0414 (6)	0.0105 (4)	0.0015 (4)	0.0084 (4)
C19	0.0370 (5)	0.0329 (5)	0.0356 (5)	0.0105 (4)	0.0036 (4)	0.0089 (4)
C20	0.0416 (6)	0.0327 (5)	0.0477 (6)	0.0089 (4)	-0.0004 (5)	0.0127 (5)
C21	0.0359 (5)	0.0349 (5)	0.0482 (6)	0.0065 (4)	0.0014 (5)	0.0157 (5)
C22	0.0417 (6)	0.0369 (5)	0.0362 (5)	0.0129 (4)	0.0036 (4)	0.0060 (4)
C23	0.0425 (6)	0.0337 (5)	0.0344 (5)	0.0112 (4)	0.0042 (4)	0.0074 (4)
C24	0.0872 (12)	0.0493 (8)	0.0610 (9)	0.0243 (8)	0.0164 (8)	0.0236 (7)
C25	0.0695 (10)	0.0806 (11)	0.0418 (7)	0.0277 (8)	0.0139 (7)	0.0017 (7)

Geometric parameters (Å, °)

S1—O2	1.4231 (10)	C9—C10	1.3623 (19)
S1—O2	1.4231 (10)	C10—C22	1.4177 (17)
S1—O1	1.4251 (9)	C10—H10	0.93
S1—O1	1.4251 (9)	C11—C18	1.3712 (17)
S1—N5	1.6594 (11)	C11—C22	1.4060 (17)
S1—C12	1.7515 (13)	C11—H11	0.93
O3—C8	1.3564 (15)	C12—C17	1.3811 (19)
O3—C24	1.416 (2)	C12—C13	1.3816 (19)
O4—C9	1.3587 (15)	C13—C14	1.380 (2)
O4—C25	1.4183 (19)	C13—H13	0.93
C1—C2	1.381 (2)	C14—C15	1.376 (3)
C1—C20	1.3872 (18)	C14—H14	0.93
C1—H1	0.93	C15—C16	1.376 (3)
C2—C3	1.375 (3)	C15—H15	0.93
C2—H2	0.93	C16—C17	1.381 (2)
C3—C4	1.390 (2)	C16—H16	0.93
C3—H3	0.93	C17—H17	0.93
C4—C21	1.3909 (17)	C18—C19	1.4139 (16)
C4—H4	0.93	C18—C20	1.4494 (16)
N5—C19	1.4321 (14)	C20—C21	1.3950 (19)
N5—C21	1.4355 (15)	C22—C23	1.4196 (17)
C6—C19	1.3705 (15)	C24—H24A	0.96
C6—C23	1.4093 (16)	C24—H24B	0.96
C6—H6	0.93	C24—H24C	0.96
C7—C8	1.3637 (17)	C25—H25A	0.96
C7—C23	1.4215 (16)	C25—H25B	0.96
C7—H7	0.93	C25—H25C	0.96
C8—C9	1.4289 (19)		
O2—S1—O1	119.60 (6)	C14—C13—C12	118.85 (16)

supplementary materials

O2—S1—N5	106.42 (6)	C14—C13—H13	120.6
O1—S1—N5	106.38 (6)	C12—C13—H13	120.6
O2—S1—C12	109.08 (6)	C15—C14—C13	120.03 (17)
O2—S1—C12	109.08 (6)	C15—C14—H14	120.0
O1—S1—C12	109.33 (6)	C13—C14—H14	120.0
O1—S1—C12	109.33 (6)	C14—C15—C16	120.53 (15)
N5—S1—C12	105.02 (5)	C14—C15—H15	119.7
C8—O3—C24	117.66 (12)	C16—C15—H15	119.7
C9—O4—C25	116.75 (12)	C15—C16—C17	120.44 (17)
C2—C1—C20	118.78 (15)	C15—C16—H16	119.8
C2—C1—H1	120.6	C17—C16—H16	119.8
C20—C1—H1	120.6	C12—C17—C16	118.38 (16)
C3—C2—C1	120.75 (14)	C12—C17—H17	120.8
C3—C2—H2	119.6	C16—C17—H17	120.8
C1—C2—H2	119.6	C11—C18—C19	120.27 (11)
C2—C3—C4	121.93 (14)	C11—C18—C20	132.36 (11)
C2—C3—H3	119.0	C19—C18—C20	107.37 (11)
C4—C3—H3	119.0	C6—C19—C18	121.62 (11)
C3—C4—C21	117.01 (15)	C6—C19—N5	129.82 (10)
C3—C4—H4	121.5	C18—C19—N5	108.55 (9)
C21—C4—H4	121.5	C1—C20—C21	120.01 (12)
C19—N5—C21	107.00 (9)	C1—C20—C18	131.88 (13)
C19—N5—S1	120.99 (8)	C21—C20—C18	108.10 (11)
C21—N5—S1	122.60 (8)	C4—C21—C20	121.51 (12)
C19—C6—C23	118.47 (11)	C4—C21—N5	129.56 (13)
C19—C6—H6	120.8	C20—C21—N5	108.88 (10)
C23—C6—H6	120.8	C11—C22—C10	121.58 (11)
C8—C7—C23	121.15 (12)	C11—C22—C23	119.36 (11)
C8—C7—H7	119.4	C10—C22—C23	119.06 (11)
C23—C7—H7	119.4	C6—C23—C22	120.47 (11)
O3—C8—C7	125.72 (13)	C6—C23—C7	120.86 (11)
O3—C8—C9	114.26 (11)	C22—C23—C7	118.67 (11)
C7—C8—C9	120.02 (12)	O3—C24—H24A	109.5
O4—C9—C10	125.54 (12)	O3—C24—H24B	109.5
O4—C9—C8	114.56 (12)	H24A—C24—H24B	109.5
C10—C9—C8	119.89 (12)	O3—C24—H24C	109.5
C9—C10—C22	121.20 (12)	H24A—C24—H24C	109.5
C9—C10—H10	119.4	H24B—C24—H24C	109.5
C22—C10—H10	119.4	O4—C25—H25A	109.5
C18—C11—C22	119.80 (11)	O4—C25—H25B	109.5
C18—C11—H11	120.1	H25A—C25—H25B	109.5
C22—C11—H11	120.1	O4—C25—H25C	109.5
C17—C12—C13	121.76 (13)	H25A—C25—H25C	109.5
C17—C12—S1	118.76 (11)	H25B—C25—H25C	109.5
C13—C12—S1	119.47 (11)		
O2—S1—O1—O1	0.00 (16)	C13—C14—C15—C16	0.0 (3)
O2—S1—O1—O1	0.00 (16)	C14—C15—C16—C17	-0.8 (3)
N5—S1—O1—O1	0.00 (14)	C13—C12—C17—C16	0.2 (2)
C12—S1—O1—O1	0.00 (17)	S1—C12—C17—C16	178.91 (12)

O1—S1—O2—O2	0.00 (14)	C15—C16—C17—C12	0.7 (3)
O1—S1—O2—O2	0.00 (14)	C22—C11—C18—C19	0.61 (19)
N5—S1—O2—O2	0.00 (15)	C22—C11—C18—C20	-179.99 (12)
C12—S1—O2—O2	0.00 (16)	C23—C6—C19—C18	-0.16 (18)
C20—C1—C2—C3	0.9 (2)	C23—C6—C19—N5	-178.68 (11)
C1—C2—C3—C4	-0.1 (2)	C11—C18—C19—C6	-0.69 (18)
C2—C3—C4—C21	-0.8 (2)	C20—C18—C19—C6	179.78 (11)
O2—S1—N5—C19	49.36 (10)	C11—C18—C19—N5	178.11 (11)
O2—S1—N5—C19	49.36 (10)	C20—C18—C19—N5	-1.42 (13)
O1—S1—N5—C19	177.92 (9)	C21—N5—C19—C6	-178.66 (12)
O1—S1—N5—C19	177.92 (9)	S1—N5—C19—C6	-31.42 (17)
C12—S1—N5—C19	-66.23 (10)	C21—N5—C19—C18	2.67 (12)
O2—S1—N5—C21	-168.55 (9)	S1—N5—C19—C18	149.90 (9)
O2—S1—N5—C21	-168.55 (9)	C2—C1—C20—C21	-0.9 (2)
O1—S1—N5—C21	-39.99 (11)	C2—C1—C20—C18	179.24 (14)
O1—S1—N5—C21	-39.99 (11)	C11—C18—C20—C1	0.0 (2)
C12—S1—N5—C21	75.86 (10)	C19—C18—C20—C1	179.45 (14)
C24—O3—C8—C7	-3.8 (2)	C11—C18—C20—C21	-179.87 (13)
C24—O3—C8—C9	175.91 (13)	C19—C18—C20—C21	-0.41 (13)
C23—C7—C8—O3	-179.89 (12)	C3—C4—C21—C20	0.81 (19)
C23—C7—C8—C9	0.4 (2)	C3—C4—C21—N5	178.13 (12)
C25—O4—C9—C10	6.8 (2)	C1—C20—C21—C4	0.02 (19)
C25—O4—C9—C8	-172.91 (13)	C18—C20—C21—C4	179.91 (11)
O3—C8—C9—O4	-1.32 (17)	C1—C20—C21—N5	-177.80 (12)
C7—C8—C9—O4	178.41 (12)	C18—C20—C21—N5	2.09 (13)
O3—C8—C9—C10	179.00 (12)	C19—N5—C21—C4	179.47 (12)
C7—C8—C9—C10	-1.3 (2)	S1—N5—C21—C4	32.88 (17)
O4—C9—C10—C22	-178.64 (12)	C19—N5—C21—C20	-2.94 (13)
C8—C9—C10—C22	1.0 (2)	S1—N5—C21—C20	-149.53 (9)
O2—S1—C12—C17	170.30 (11)	C18—C11—C22—C10	-179.74 (11)
O2—S1—C12—C17	170.30 (11)	C18—C11—C22—C23	0.28 (19)
O1—S1—C12—C17	37.83 (12)	C9—C10—C22—C11	-179.87 (12)
O1—S1—C12—C17	37.83 (12)	C9—C10—C22—C23	0.11 (19)
N5—S1—C12—C17	-75.97 (12)	C19—C6—C23—C22	1.07 (18)
O2—S1—C12—C13	-10.97 (13)	C19—C6—C23—C7	-179.10 (12)
O2—S1—C12—C13	-10.97 (13)	C11—C22—C23—C6	-1.14 (18)
O1—S1—C12—C13	-143.44 (11)	C10—C22—C23—C6	178.88 (11)
O1—S1—C12—C13	-143.44 (11)	C11—C22—C23—C7	179.02 (12)
N5—S1—C12—C13	102.76 (11)	C10—C22—C23—C7	-0.95 (18)
C17—C12—C13—C14	-1.0 (2)	C8—C7—C23—C6	-179.15 (12)
S1—C12—C13—C14	-179.69 (12)	C8—C7—C23—C22	0.7 (2)
C12—C13—C14—C15	0.9 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C4—H4 \cdots O1	0.93	2.34	2.9238 (19)	120
C6—H6 \cdots O2	0.93	2.37	2.9674 (15)	122
C2—H2 \cdots O2 ⁱ	0.93	2.55	3.3361 (17)	143

supplementary materials

C24—H24C...Cg1 ⁱⁱ	0.96	2.84	3.7272 (21)	154
C25—H25A...Cg2 ⁱⁱⁱ	0.96	2.90	3.6276 (21)	134

Symmetry codes: (i) $x, y-1, z$; (ii) $-x+2, -y+1, -z$; (iii) $-x+2, -y, -z$.

Fig. 1




Fig. 2

