

4-Acetamido-N-(λ^5 -triphenylphosphoranylidene)benzenesulfonamideBiserka Prugovečki,^{a*} Marina Marinković,^b Mladen Vinković^{b‡} and Miljenko Dumić^c^aLaboratory of General and Inorganic Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, HR-10000 Zagreb, Croatia, ^bPLIVA Croatia Ltd, TAPI Research and Development, Prilaz baruna Filipovića 29, HR-10000 Zagreb, Croatia, and ^cDepartment of Biotechnology, University of Rijeka, S. Krautzeka bb, HR-51000 Rijeka, Croatia

Correspondence e-mail: biserka@chem.pmf.hr


Received 8 March 2010; accepted 13 April 2010

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.041; wR factor = 0.106; data-to-parameter ratio = 14.2.

There are two independent molecules per asymmetric unit of the title compound, $\text{C}_{26}\text{H}_{23}\text{N}_2\text{O}_3\text{PS}$. Their superposition shows that they differ in the conformation of the $\text{CH}_3\text{CO}-$ group and the benzene rings from the triphenylphosphorane group. In the crystal structure, independent molecules are interconnected by strong $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, forming infinite chains along the a axis.

Related literature

For related structures, see: Andersen *et al.* (1999, 2001, 2004); Matano *et al.* (2002); Monkowius *et al.* (2004); Zhu *et al.* (1997). For the synthesis, see: Ashley *et al.* (1947); Khmel'nitzkaya & Mikhel's (1934). For structural and synthetic studies of azirine antihyperglycaemics, see: Dumić *et al.* (1993, 1995); Filić *et al.* (1996); Orešić *et al.* (2001); Prugovečki *et al.* (2005, 2006); Vinković *et al.* (1993); Žegarac *et al.* (2010).



Experimental

Crystal data

 $\text{C}_{26}\text{H}_{23}\text{N}_2\text{O}_3\text{PS}$
 $M_r = 474.49$ Monoclinic, $P2_1/n$
 $a = 15.0419$ (10) Å[‡] Present address: Astex Therapeutics, 436 Cambridge Science Park, Cambridge CB4 0QA, England. $b = 18.6355$ (10) Å
 $c = 18.5917$ (18) Å
 $\beta = 113.413$ (10)°
 $V = 4782.4$ (6) Å³
 $Z = 8$ Mo $K\alpha$ radiation
 $\mu = 0.23$ mm⁻¹
 $T = 295$ K
 $0.56 \times 0.30 \times 0.15$ mm

Data collection

Oxford Diffraction Xcalibur CCD diffractometer
32329 measured reflections8441 independent reflections
6309 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.106$
 $S = 1.04$
8441 reflections595 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.25$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.28$ e Å⁻³**Table 1**
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2}-\text{H2}\cdots\text{O2}^{\text{ii}}$	0.86	2.11	2.966 (2)	173
$\text{N2}'-\text{H2}'\cdots\text{O2}^{\text{ii}}$	0.86	2.11	2.961 (2)	174

Symmetry codes: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2003); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2003); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

The authors thank the Ministry of Science and Technology of the Republic of Croatia for partial financial support of this work (grant Nos. 0006543 and 0119632).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BG2337).

References

- Andersen, N. G., Parvez, M., McDonald, R. & Keay, B. A. (2004). *Can. J. Chem.* **82**, 145–17161.
- Andersen, N. G., Ramsden, P. D., Che, D., Parvez, M. & Keay, B. A. (1999). *Org. Lett.* **1**, 2009–2011.
- Andersen, N. G., Ramsden, P. D., Che, D., Parvez, M. & Keay, B. A. (2001). *J. Org. Chem.* **66**, 7478–7486.
- Ashley, J. N., Buchanan, G. L. & Easson, A. P. T. (1947). *J. Chem. Soc.* pp. 60–67.
- Dumić, M., Filić, D., Vinković, M., Jamnicki, B. & Kamenar, B. (1993). *Tetrahedron Lett.* **34**, 3639–3642.
- Dumić, M., Vinković, M., Filić, D., Jamnicki, B., Eškinja, M. & Kamenar, B. (1995). *J. Med. Chem.* **38**, 3034–3042.
- Filić, D., Vinković, M., Jamnicki, B. & Dumić, M. (1996). *Croat. Chem. Acta*, **69**, 631–641.
- Khmel'nitzkaya, I. & Mikhel's, V. (1934). *Zh. Obshch. Khim.* pp. 452–455.
- Matano, Y., Nomura, H. & Suzuki, H. (2002). *Inorg. Chem.* **41**, 1940–1948.
- Monkowius, U. V., Nogai, S. & Schmidbaur, H. (2004). *Dalton Trans.* pp. 1610–1617.
- Orešić, M., Filić, D., Prugovečki, B., Vinković, M. & Dumić, M. (2001). *Croat. Chem. Acta*, **74**, 667–682.
- Oxford Diffraction (2003). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, England.
- Prugovečki, B., Marinković, M., Vinković, M. & Dumić, M. (2005). *Acta Cryst. E* **61**, o2844–o2846.
- Prugovečki, B., Marinković, M., Vinković, M. & Dumić, M. (2006). *Croat. Chem. Acta*, **79**, 219–226.

Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

Vinković, M., Dumić, M. & Kamenar, B. (1993). *Acta Cryst.* **C49**, 1661–1663.

Žegarac, M., Marinković, M., Prugovečki, B., Vinković, M. & Dumić, M. (2010). In preparation.

Zhu, S., Xu, B., Qin, C. & Xu, G. (1997). *Inorg. Chem.* **36**, 4909–4911.

supplementary materials

Acta Cryst. (2010). E66, o1098-o1099 [doi:10.1107/S1600536810013620]

4-Acetamido-*N*-(λ^5 -triphenylphosphoranylidene)benzenesulfonamide

B. Prugovecki, M. Marinkovic, M. Vinkovic and M. Dumic

Comment

As a part of our ongoing research on the synthetic and structural studies of 1-sulfonyl-1a,2,6,6a-tetrahydro-1H,4H-[1,3]dioxepino[5,6-b]azirine antihyperglycaemics (Dumić *et al.* 1993 ; 1995, Filić *et al.* 1996, Vinković *et al.* 1993, Orešić *et al.* 2001 and Prugovečki *et al.* 2005 ; 2006), we required suitable synthons carrying 4-acetylaminobenzenesulfanyl and 4-acetylaminobenzenesulfonyl pattern. Thus, the 4-acetylaminobenzenesulfonylimino-triphenylphosphorane (Title compound, I) and bis(4-acetylaminophenyl) disulfide compound (II) were chosen for this study. We prepared both of them in the same reaction, i.e. by treatment of 4-acetylaminobenzenesulfonylazide with triphenylphosphine in acetonitrile at room temperature (Scheme 1). 4-Acetylaminobenzenesulfonylimino-triphenylphosphorane (Title compound, I) was obtained as colorless prisms (m.p. 495-497 K). Bis(4-acetylaminophenyl) disulfide (compound II) was obtained as a yellow solid (m.p. 485-488 K), i.e. in one of its three known forms; m.ps. 488 K, 454-455 K and 395 K respectively (Khmel'nitzkaya, *et al.* 1934). Its structure and solid state behaviour will be published elsewhere (Žegarac *et al.* 2010).

In the title compound, C₂₆H₂₃N₂O₃PS, (I), there are two independent molecules per asymmetric unit. Their superposition shows that they are different in the conformation of the CH₃CO group and the benzene rings from the triphenylphosphorane group. In the crystal structure independent molecules are interconnected by strong N—H···O hydrogen bonds forming infinite one-dimensional chains along the *a* axis.

Experimental

Triphenylphosphine (28.18 g, 0.107 mol) was added in small portions to a stirred solution of 4-acetylaminobenzenesulfonyl azide (13.0 g, 50.4 mmol) [prepared according Ashley *et al.* (1947)] in acetonitrile (211 ml) at 273 K. After being stirred for 3 hrs at room temperature, the mixture was concentrated under reduced pressure to dryness. The residue was purified by silicagel column chromatography (dichloromethane-methanol-25 % ammonia, 10:1:0.3 v/v) to afford the title compound (I) as a colorless solid [5.9 g, 22.7 %; m.p. 491-495 K; R_f = 0.32 (dichloromethane-methanol- 25 % ammonia, 10:1:0.3 v/v)]. Single crystals suitable for X-ray diffraction were prepared by recrystallization from ethyl acetate-methanol, 1:1 v/v). M.p. 495-497 K. Spectroscopic analysis: IR (KBr) $\nu_{\max}/\text{cm}^{-1}$: 3305, 3269, 3190, 3116, 3057, 1691, 1594, 1536, 1485, 1437, 1400, 1372, 1315, 1252, 1194, 1170, 1131, 1085, 1031, 1015, 998, 954, 851, 801, 790, 753, 723, 692, 638, 620. ¹H NMR (DMSO-d₆) δ/ppm : 10.12 (s, 1H, NH), 7.52 i 7.40 (d.d., 4H, J=4.3, H-arom.), 7.73-7.68 and 7.59-7.57 (2 m, 15H, H-arom.), 2.06 (s, 3H, CH₃). ¹³C NMR (DMSO-d₆) δ/ppm : 141.10 (s), 140.32 (s), 125.93 (d), 117.96 (d) (C arom.), 133.05 (d), 132.60 (d), 128.90 (d), 126.94 (d, J(C—P)=102.9) (C-arom), 24.04 (q, CH₃).

Evaporation of other selected fractions to dryness afford bis(4-acetylaminophenyl) disulfide (II) as a TLC pure yellow solid [5.7 g, 31.5 % m.p. 485-488 K; R_f = 0.26 (dichloromethane-methanol-25 % ammonia, 10:1:0.3 v/v)]. Spectroscopic analysis: IR (KBr) $\nu_{\max}/\text{cm}^{-1}$: 3291, 3246, 3178, 3105, 3058, 1681, 1658, 1608, 1593, 1538, 1490, 1397, 1367, 1317, 1292,

supplementary materials

1263, 1175, 1121, 1014, 967, 838, 825, 816, 758, 703, 604. ^1H NMR (DMSO- d_6) δ /ppm: 10.71 (s, 2H, NH), 7.59 and 7.42 (dd, 8H, H arom, $J=8.7$), 2.04 (s, 6H, CH_3). ^{13}C NMR (DMSO- d_6) δ /ppm: 168.50 (s, CO), 139.50 (s), 130.10 (d), 129.10 (s), 119.50 (d) (C arom), 24.20 (q, CH_3).

Refinement

H atoms were positioned geometrically, C-H: 0.93-0.96Å, N-H: 0.86Å, and allowed to ride, with $U(\text{H})=1.2/1.5 \times U_{\text{eq}}(\text{host})$. In order to avoid beamstop shadowing effects $\theta(\text{min})$ was set to 3.24° , with what 20 reflections below this value were left aside the data set.

Figures




Fig. 1. View of the molecule I with the atom labelling scheme. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. Overlapped structures of independent molecules of compound I, showing structural differences.




Fig. 3. Packing of the molecules in the unit cell. Hydrogen bonds ($d(\text{N2}-\text{H2}\cdots\text{O2}'(-x + 1/2, y - 1/2, -z+1/2))=2.966 (2) \text{ \AA}$ and $\text{N2}'-\text{H2}'\cdots\text{O2}(-x + 3/2, y + 1/2, -z+1/2))=2.961 (2) \text{ \AA}$) are shown as dotted lines and hydrogen atoms that are not involved in hydrogen bonding are omitted for clarity.




Fig. 4. Synthetic route to the molecule I.

4-Acetamido-*N*-(λ^5 -triphenylphosphoranylidene)benzenesulfonamide

Crystal data

$\text{C}_{26}\text{H}_{23}\text{N}_2\text{O}_3\text{PS}$

$M_r = 474.49$

Monoclinic, $P2_1/n$

Hall symbol: $-P 2_1n$

$a = 15.0419 (10) \text{ \AA}$

$b = 18.6355 (10) \text{ \AA}$

$c = 18.5917 (18) \text{ \AA}$

$\beta = 113.413 (10)^\circ$

$F(000) = 1984$

$D_x = 1.318 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1548 reflections

$\theta = 15\text{--}25^\circ$

$\mu = 0.23 \text{ mm}^{-1}$

$T = 295 \text{ K}$

Plate, colourless

$V = 4782.4 (6) \text{ \AA}^3$
 $Z = 8$

$0.56 \times 0.30 \times 0.15 \text{ mm}$

Data collection

Oxford Diffraction Xcalibur CCD diffractometer	6309 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube graphite	$R_{\text{int}} = 0.029$ $\theta_{\text{max}} = 25.1^\circ$, $\theta_{\text{min}} = 3.2^\circ$
CCD scans	$h = -17 \rightarrow 17$
32329 measured reflections	$k = -22 \rightarrow 22$
8441 independent reflections	$l = -22 \rightarrow 21$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.041$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.106$	H-atom parameters constrained
$S = 1.04$	$w = 1/[\sigma^2(F_o^2) + (0.0611P)^2 + 0.0767P]$ where $P = (F_o^2 + 2F_c^2)/3$
8441 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
595 parameters	$\Delta\rho_{\text{max}} = 0.25 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.28 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.60435 (4)	0.26549 (3)	0.06832 (3)	0.03823 (15)
P1	0.73384 (4)	0.22036 (3)	0.22228 (3)	0.03282 (14)
O1	0.63908 (11)	0.33781 (8)	0.08560 (10)	0.0540 (4)
O2	0.59940 (12)	0.23771 (9)	-0.00544 (8)	0.0545 (4)
O3	0.15446 (12)	0.38553 (10)	0.01877 (11)	0.0640 (5)
N1	0.66148 (12)	0.20983 (9)	0.13335 (10)	0.0381 (4)
N2	0.19539 (13)	0.26777 (10)	0.03546 (10)	0.0439 (5)
H2	0.1725	0.2259	0.0379	0.053*
C5	0.13239 (17)	0.32332 (14)	0.02196 (13)	0.0492 (6)
C6	0.0322 (2)	0.30213 (17)	0.0123 (2)	0.0857 (10)
H6A	0.0295	0.2985	0.0629	0.128*
H6B	0.0159	0.2566	-0.0139	0.128*
H6C	-0.0130	0.3378	-0.0183	0.128*
C11	0.74745 (15)	0.13304 (11)	0.26552 (12)	0.0364 (5)
C12	0.72650 (19)	0.07279 (12)	0.21875 (15)	0.0559 (7)
H12	0.7056	0.0773	0.1647	0.067*
C13	0.7369 (2)	0.00600 (13)	0.25234 (19)	0.0715 (8)
H13	0.7216	-0.0346	0.2207	0.086*

supplementary materials

C14	0.76928 (19)	-0.00145 (13)	0.33235 (17)	0.0595 (7)
H14	0.7773	-0.0469	0.3546	0.071*
C15	0.78979 (19)	0.05756 (13)	0.37871 (15)	0.0576 (7)
H15	0.8114	0.0527	0.4328	0.069*
C16	0.77845 (18)	0.12463 (12)	0.34556 (13)	0.0492 (6)
H16	0.7919	0.1650	0.3775	0.059*
C21	0.85231 (15)	0.25325 (11)	0.23595 (12)	0.0372 (5)
C22	0.93602 (18)	0.22040 (15)	0.28608 (15)	0.0589 (7)
H22	0.9322	0.1797	0.3136	0.071*
C23	1.02521 (19)	0.24721 (17)	0.29590 (17)	0.0716 (8)
H23	1.0811	0.2241	0.3295	0.086*
C24	1.03252 (19)	0.30698 (16)	0.25716 (17)	0.0641 (7)
H24	1.0930	0.3255	0.2647	0.077*
C25	0.9506 (2)	0.33945 (15)	0.20712 (18)	0.0690 (8)
H25	0.9553	0.3802	0.1800	0.083*
C26	0.86062 (18)	0.31328 (13)	0.19594 (16)	0.0562 (7)
H26	0.8052	0.3361	0.1611	0.067*
C31	0.68889 (15)	0.27629 (11)	0.27907 (12)	0.0352 (5)
C32	0.59488 (17)	0.26534 (13)	0.27220 (14)	0.0498 (6)
H32	0.5557	0.2321	0.2363	0.060*
C33	0.5591 (2)	0.30353 (15)	0.31856 (16)	0.0621 (7)
H33	0.4961	0.2957	0.3142	0.074*
C34	0.6167 (2)	0.35311 (14)	0.37095 (15)	0.0592 (7)
H34	0.5927	0.3789	0.4021	0.071*
C35	0.7090 (2)	0.36480 (13)	0.37737 (14)	0.0578 (7)
H35	0.7473	0.3988	0.4127	0.069*
C36	0.74610 (18)	0.32633 (12)	0.33204 (13)	0.0462 (6)
H36	0.8094	0.3342	0.3371	0.055*
C41	0.48472 (15)	0.26548 (11)	0.06320 (11)	0.0345 (5)
C42	0.43550 (17)	0.20264 (11)	0.06204 (13)	0.0442 (6)
H42	0.4667	0.1587	0.0668	0.053*
C43	0.34066 (17)	0.20470 (11)	0.05396 (13)	0.0435 (5)
H43	0.3082	0.1622	0.0538	0.052*
C44	0.29302 (15)	0.26972 (11)	0.04594 (11)	0.0368 (5)
C45	0.34292 (16)	0.33258 (12)	0.04923 (13)	0.0438 (5)
H45	0.3123	0.3766	0.0457	0.053*
C46	0.43846 (16)	0.32999 (11)	0.05761 (13)	0.0419 (5)
H46	0.4718	0.3725	0.0596	0.050*
S1'	0.39565 (4)	0.60029 (3)	0.37549 (3)	0.03796 (15)
P1'	0.29976 (4)	0.48985 (3)	0.27043 (3)	0.03572 (15)
O1'	0.36387 (11)	0.64762 (8)	0.30907 (9)	0.0540 (4)
O2'	0.37677 (11)	0.62497 (8)	0.44208 (9)	0.0499 (4)
O3'	0.86803 (13)	0.47540 (10)	0.53859 (12)	0.0703 (5)
N1'	0.35642 (13)	0.52178 (9)	0.35569 (10)	0.0399 (4)
N2'	0.82514 (13)	0.59181 (10)	0.50987 (11)	0.0456 (5)
H2'	0.8495	0.6340	0.5122	0.055*
C03'	0.0174 (2)	0.52015 (17)	0.12570 (19)	0.0807 (10)
H23'	-0.0241	0.5092	0.0745	0.097*
C5'	0.89003 (17)	0.53794 (15)	0.54146 (13)	0.0493 (6)

C6'	0.99288 (18)	0.56331 (16)	0.58155 (16)	0.0698 (8)
H6'1	1.0212	0.5429	0.6332	0.105*
H6'2	0.9940	0.6147	0.5853	0.105*
H6'3	1.0292	0.5484	0.5518	0.105*
C11'	0.30394 (14)	0.39466 (11)	0.28443 (12)	0.0348 (5)
C12'	0.28480 (17)	0.34757 (12)	0.22329 (13)	0.0475 (6)
H12'	0.2732	0.3650	0.1735	0.057*
C13'	0.28260 (19)	0.27466 (13)	0.23523 (14)	0.0560 (7)
H13'	0.2698	0.2431	0.1936	0.067*
C14'	0.29939 (18)	0.24874 (13)	0.30821 (14)	0.0531 (6)
H14'	0.2978	0.1996	0.3161	0.064*
C15'	0.31851 (19)	0.29467 (13)	0.36925 (14)	0.0554 (7)
H15'	0.3294	0.2767	0.4187	0.066*
C16'	0.32181 (18)	0.36738 (12)	0.35837 (13)	0.0492 (6)
H16'	0.3360	0.3984	0.4006	0.059*
C21'	0.17467 (16)	0.51602 (12)	0.23034 (13)	0.0429 (5)
C22'	0.11238 (18)	0.49844 (14)	0.15399 (16)	0.0615 (7)
H22'	0.1352	0.4722	0.1223	0.074*
C24'	-0.0166 (2)	0.55839 (19)	0.1731 (2)	0.0878 (10)
H24'	-0.0808	0.5734	0.1535	0.105*
C25'	0.0431 (2)	0.57394 (18)	0.2480 (2)	0.0856 (10)
H25'	0.0192	0.5986	0.2800	0.103*
C26'	0.13908 (18)	0.55348 (14)	0.27692 (16)	0.0609 (7)
H26'	0.1800	0.5651	0.3281	0.073*
C31'	0.35067 (17)	0.51011 (12)	0.19990 (13)	0.0437 (5)
C32'	0.43991 (19)	0.48131 (15)	0.21220 (17)	0.0645 (7)
H32'	0.4698	0.4499	0.2538	0.077*
C33'	0.4851 (2)	0.4989 (2)	0.1627 (2)	0.0864 (10)
H33'	0.5454	0.4794	0.1714	0.104*
C34'	0.4426 (3)	0.5441 (2)	0.1020 (2)	0.0937 (12)
H34'	0.4735	0.5553	0.0689	0.112*
C35'	0.3549 (3)	0.57337 (19)	0.08893 (19)	0.0904 (11)
H35'	0.3259	0.6045	0.0470	0.108*
C36'	0.3084 (2)	0.55685 (15)	0.13822 (16)	0.0690 (8)
H36'	0.2486	0.5773	0.1295	0.083*
C41'	0.52275 (15)	0.59244 (11)	0.40911 (12)	0.0336 (5)
C42'	0.57060 (15)	0.53071 (11)	0.44354 (12)	0.0355 (5)
H42'	0.5352	0.4904	0.4455	0.043*
C43'	0.67037 (16)	0.52770 (11)	0.47535 (12)	0.0374 (5)
H43'	0.7019	0.4853	0.4977	0.045*
C44'	0.72346 (15)	0.58761 (11)	0.47384 (12)	0.0361 (5)
C45'	0.67459 (16)	0.64967 (12)	0.43796 (13)	0.0430 (5)
H45'	0.7098	0.6900	0.4358	0.052*
C46'	0.57519 (16)	0.65226 (12)	0.40549 (12)	0.0408 (5)
H46'	0.5433	0.6940	0.3813	0.049*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0389 (3)	0.0373 (3)	0.0383 (3)	0.0041 (2)	0.0152 (2)	0.0027 (2)
P1	0.0348 (3)	0.0288 (3)	0.0341 (3)	0.0015 (2)	0.0129 (2)	-0.0013 (2)
O1	0.0462 (10)	0.0352 (9)	0.0782 (12)	-0.0018 (7)	0.0221 (8)	0.0049 (8)
O2	0.0603 (11)	0.0717 (11)	0.0345 (8)	0.0177 (9)	0.0221 (8)	0.0050 (8)
O3	0.0454 (11)	0.0511 (11)	0.0861 (13)	0.0109 (9)	0.0160 (9)	0.0114 (10)
N1	0.0396 (11)	0.0365 (10)	0.0349 (10)	0.0054 (8)	0.0113 (8)	-0.0015 (8)
N2	0.0400 (11)	0.0431 (11)	0.0482 (11)	-0.0026 (9)	0.0170 (9)	0.0012 (9)
C5	0.0395 (14)	0.0572 (16)	0.0450 (14)	0.0049 (13)	0.0104 (11)	0.0061 (12)
C6	0.0487 (18)	0.086 (2)	0.122 (3)	0.0049 (16)	0.0332 (18)	0.009 (2)
C11	0.0347 (12)	0.0310 (11)	0.0430 (12)	0.0026 (9)	0.0149 (10)	-0.0007 (10)
C12	0.0745 (19)	0.0366 (13)	0.0501 (15)	0.0027 (13)	0.0178 (13)	-0.0059 (11)
C13	0.090 (2)	0.0291 (14)	0.085 (2)	0.0007 (14)	0.0236 (18)	-0.0074 (14)
C14	0.0590 (17)	0.0377 (14)	0.077 (2)	0.0026 (12)	0.0215 (15)	0.0157 (13)
C15	0.0685 (18)	0.0488 (16)	0.0534 (15)	0.0060 (13)	0.0220 (13)	0.0142 (13)
C16	0.0641 (17)	0.0369 (13)	0.0426 (13)	0.0002 (12)	0.0169 (12)	0.0032 (11)
C21	0.0369 (13)	0.0368 (12)	0.0389 (12)	0.0020 (10)	0.0162 (10)	-0.0053 (10)
C22	0.0427 (15)	0.0688 (18)	0.0598 (16)	-0.0004 (13)	0.0146 (13)	0.0184 (14)
C23	0.0377 (16)	0.095 (2)	0.0728 (19)	0.0017 (15)	0.0120 (14)	0.0172 (17)
C24	0.0409 (16)	0.080 (2)	0.0736 (19)	-0.0127 (14)	0.0248 (14)	-0.0107 (16)
C25	0.0576 (19)	0.0526 (16)	0.100 (2)	-0.0094 (14)	0.0350 (17)	0.0108 (16)
C26	0.0421 (15)	0.0423 (14)	0.0804 (18)	0.0015 (11)	0.0203 (13)	0.0119 (13)
C31	0.0427 (13)	0.0300 (11)	0.0331 (11)	0.0034 (10)	0.0151 (10)	0.0008 (9)
C32	0.0465 (15)	0.0543 (15)	0.0506 (14)	-0.0021 (12)	0.0214 (12)	-0.0123 (12)
C33	0.0537 (17)	0.0761 (19)	0.0657 (17)	0.0099 (14)	0.0336 (14)	-0.0046 (15)
C34	0.084 (2)	0.0533 (16)	0.0495 (15)	0.0160 (15)	0.0357 (15)	-0.0042 (12)
C35	0.082 (2)	0.0431 (14)	0.0485 (15)	-0.0036 (13)	0.0263 (14)	-0.0140 (12)
C36	0.0547 (15)	0.0374 (13)	0.0479 (14)	-0.0033 (11)	0.0218 (12)	-0.0064 (11)
C41	0.0376 (12)	0.0326 (12)	0.0310 (11)	0.0018 (10)	0.0110 (9)	-0.0007 (9)
C42	0.0477 (15)	0.0297 (12)	0.0518 (14)	0.0052 (10)	0.0162 (11)	-0.0008 (10)
C43	0.0452 (14)	0.0321 (12)	0.0518 (14)	-0.0049 (10)	0.0176 (11)	0.0000 (10)
C44	0.0356 (12)	0.0425 (13)	0.0293 (11)	0.0003 (10)	0.0097 (9)	0.0001 (10)
C45	0.0480 (14)	0.0321 (12)	0.0547 (14)	0.0063 (11)	0.0240 (12)	0.0057 (10)
C46	0.0449 (14)	0.0313 (12)	0.0513 (14)	-0.0017 (10)	0.0210 (11)	0.0020 (10)
S1'	0.0363 (3)	0.0317 (3)	0.0452 (3)	0.0027 (2)	0.0155 (2)	-0.0002 (2)
P1'	0.0336 (3)	0.0361 (3)	0.0358 (3)	0.0005 (2)	0.0120 (2)	0.0008 (2)
O1'	0.0474 (10)	0.0444 (10)	0.0613 (11)	0.0077 (8)	0.0123 (8)	0.0164 (8)
O2'	0.0499 (10)	0.0455 (9)	0.0630 (10)	-0.0013 (8)	0.0318 (8)	-0.0155 (8)
O3'	0.0495 (11)	0.0474 (11)	0.1054 (16)	0.0086 (9)	0.0216 (10)	0.0047 (10)
N1'	0.0435 (11)	0.0343 (10)	0.0388 (10)	-0.0022 (8)	0.0131 (8)	-0.0006 (8)
N2'	0.0366 (11)	0.0444 (11)	0.0549 (12)	-0.0016 (9)	0.0174 (9)	0.0056 (9)
C03'	0.0468 (18)	0.084 (2)	0.080 (2)	0.0057 (16)	-0.0078 (16)	0.0192 (18)
C5'	0.0418 (15)	0.0585 (17)	0.0466 (14)	0.0058 (13)	0.0166 (11)	-0.0009 (12)
C6'	0.0415 (16)	0.080 (2)	0.078 (2)	0.0046 (15)	0.0131 (14)	-0.0015 (16)
C11'	0.0310 (11)	0.0374 (12)	0.0349 (11)	-0.0026 (9)	0.0119 (9)	-0.0030 (9)

C12'	0.0583 (16)	0.0457 (14)	0.0362 (12)	-0.0105 (12)	0.0164 (11)	-0.0010 (10)
C13'	0.0768 (19)	0.0414 (14)	0.0445 (14)	-0.0163 (13)	0.0185 (13)	-0.0119 (11)
C14'	0.0650 (17)	0.0378 (13)	0.0519 (15)	-0.0100 (12)	0.0183 (13)	-0.0007 (12)
C15'	0.0785 (19)	0.0460 (15)	0.0393 (13)	-0.0033 (13)	0.0210 (13)	0.0074 (11)
C16'	0.0684 (17)	0.0405 (13)	0.0370 (13)	-0.0005 (12)	0.0192 (12)	-0.0059 (10)
C21'	0.0363 (13)	0.0423 (13)	0.0479 (14)	0.0040 (10)	0.0145 (11)	0.0066 (11)
C22'	0.0469 (16)	0.0644 (18)	0.0604 (17)	0.0041 (13)	0.0078 (13)	-0.0002 (13)
C24'	0.0404 (17)	0.095 (3)	0.120 (3)	0.0178 (17)	0.0229 (19)	0.020 (2)
C25'	0.058 (2)	0.102 (3)	0.105 (3)	0.0288 (19)	0.041 (2)	0.011 (2)
C26'	0.0495 (16)	0.0701 (18)	0.0644 (17)	0.0122 (14)	0.0240 (13)	0.0032 (14)
C31'	0.0496 (14)	0.0414 (13)	0.0436 (13)	-0.0076 (11)	0.0224 (11)	-0.0033 (10)
C32'	0.0516 (17)	0.0740 (19)	0.0761 (19)	-0.0051 (14)	0.0341 (15)	-0.0010 (15)
C33'	0.070 (2)	0.107 (3)	0.106 (3)	-0.021 (2)	0.061 (2)	-0.021 (2)
C34'	0.124 (3)	0.101 (3)	0.090 (3)	-0.052 (3)	0.078 (3)	-0.025 (2)
C35'	0.131 (3)	0.089 (2)	0.068 (2)	-0.014 (2)	0.058 (2)	0.0175 (18)
C36'	0.087 (2)	0.0662 (18)	0.0634 (18)	0.0045 (16)	0.0396 (16)	0.0163 (15)
C41'	0.0374 (12)	0.0333 (11)	0.0335 (11)	0.0008 (9)	0.0178 (9)	-0.0021 (9)
C42'	0.0404 (13)	0.0269 (11)	0.0416 (12)	-0.0018 (9)	0.0190 (10)	-0.0027 (9)
C43'	0.0420 (13)	0.0318 (12)	0.0410 (12)	0.0058 (10)	0.0193 (10)	0.0011 (9)
C44'	0.0359 (12)	0.0413 (13)	0.0348 (11)	0.0009 (10)	0.0178 (10)	0.0019 (9)
C45'	0.0401 (13)	0.0415 (13)	0.0527 (14)	-0.0026 (11)	0.0239 (11)	0.0093 (11)
C46'	0.0451 (14)	0.0353 (12)	0.0450 (13)	0.0052 (10)	0.0210 (11)	0.0106 (10)

Geometric parameters (Å, °)

S1—O1	1.4348 (16)	S1'—O1'	1.4361 (15)
S1—O2	1.4398 (15)	S1'—O2'	1.4506 (15)
S1—N1	1.5655 (17)	S1'—N1'	1.5658 (18)
S1—C41	1.764 (2)	S1'—C41'	1.766 (2)
P1—N1	1.5896 (18)	P1'—N1'	1.5872 (18)
P1—C11	1.790 (2)	P1'—C11'	1.790 (2)
P1—C31	1.796 (2)	P1'—C21'	1.794 (2)
P1—C21	1.804 (2)	P1'—C31'	1.802 (2)
O3—C5	1.214 (3)	O3'—C5'	1.207 (3)
N2—C5	1.358 (3)	N2'—C5'	1.360 (3)
N2—C44	1.403 (3)	N2'—C44'	1.407 (3)
N2—H2	0.8605	N2'—H2'	0.8606
C5—C6	1.498 (3)	C03'—C22'	1.373 (4)
C6—H6A	0.9600	C03'—C24'	1.379 (5)
C6—H6B	0.9600	C03'—H23'	0.9300
C6—H6C	0.9600	C5'—C6'	1.502 (3)
C11—C16	1.380 (3)	C6'—H6'1	0.9600
C11—C12	1.378 (3)	C6'—H6'2	0.9600
C12—C13	1.373 (3)	C6'—H6'3	0.9600
C12—H12	0.9300	C11'—C12'	1.373 (3)
C13—C14	1.376 (4)	C11'—C16'	1.388 (3)
C13—H13	0.9300	C12'—C13'	1.379 (3)
C14—C15	1.355 (3)	C12'—H12'	0.9300
C14—H14	0.9300	C13'—C14'	1.366 (3)

supplementary materials

C15—C16	1.374 (3)	C13'—H13'	0.9300
C15—H15	0.9300	C14'—C15'	1.358 (3)
C16—H16	0.9300	C14'—H14'	0.9300
C21—C22	1.377 (3)	C15'—C16'	1.374 (3)
C21—C26	1.376 (3)	C15'—H15'	0.9300
C22—C23	1.374 (3)	C16'—H16'	0.9300
C22—H22	0.9300	C21'—C26'	1.376 (3)
C23—C24	1.354 (4)	C21'—C22'	1.394 (3)
C23—H23	0.9300	C22'—H22'	0.9300
C24—C25	1.357 (4)	C24'—C25'	1.355 (5)
C24—H24	0.9300	C24'—H24'	0.9300
C25—C26	1.374 (3)	C25'—C26'	1.379 (4)
C25—H25	0.9300	C25'—H25'	0.9300
C26—H26	0.9300	C26'—H26'	0.9300
C31—C32	1.384 (3)	C31'—C32'	1.378 (3)
C31—C36	1.381 (3)	C31'—C36'	1.378 (3)
C32—C33	1.381 (3)	C32'—C33'	1.383 (4)
C32—H32	0.9300	C32'—H32'	0.9300
C33—C34	1.372 (4)	C33'—C34'	1.349 (5)
C33—H33	0.9300	C33'—H33'	0.9300
C34—C35	1.362 (4)	C34'—C35'	1.357 (5)
C34—H34	0.9300	C34'—H34'	0.9300
C35—C36	1.382 (3)	C35'—C36'	1.390 (4)
C35—H35	0.9300	C35'—H35'	0.9300
C36—H36	0.9300	C36'—H36'	0.9300
C41—C46	1.372 (3)	C41'—C42'	1.373 (3)
C41—C42	1.381 (3)	C41'—C46'	1.383 (3)
C42—C43	1.374 (3)	C42'—C43'	1.378 (3)
C42—H42	0.9300	C42'—H42'	0.9300
C43—C44	1.385 (3)	C43'—C44'	1.379 (3)
C43—H43	0.9300	C43'—H43'	0.9300
C44—C45	1.379 (3)	C44'—C45'	1.390 (3)
C45—C46	1.383 (3)	C45'—C46'	1.373 (3)
C45—H45	0.9300	C45'—H45'	0.9300
C46—H46	0.9300	C46'—H46'	0.9300
O1—S1—O2	115.31 (10)	O1'—S1'—O2'	115.34 (10)
O1—S1—N1	114.21 (10)	O1'—S1'—N1'	113.96 (10)
O2—S1—N1	107.58 (9)	O2'—S1'—N1'	108.28 (9)
O1—S1—C41	106.56 (10)	O1'—S1'—C41'	107.39 (10)
O2—S1—C41	106.44 (10)	O2'—S1'—C41'	106.02 (9)
N1—S1—C41	106.06 (10)	N1'—S1'—C41'	105.05 (10)
N1—P1—C11	105.29 (9)	N1'—P1'—C11'	104.62 (9)
N1—P1—C31	114.90 (9)	N1'—P1'—C21'	111.80 (10)
C11—P1—C31	105.43 (9)	C11'—P1'—C21'	107.47 (10)
N1—P1—C21	114.72 (9)	N1'—P1'—C31'	115.68 (10)
C11—P1—C21	108.37 (10)	C11'—P1'—C31'	108.23 (10)
C31—P1—C21	107.54 (10)	C21'—P1'—C31'	108.63 (11)
S1—N1—P1	131.33 (11)	S1'—N1'—P1'	126.00 (11)
C5—N2—C44	128.4 (2)	C5'—N2'—C44'	128.5 (2)

C5—N2—H2	115.8	C5'—N2'—H2'	115.7
C44—N2—H2	115.8	C44'—N2'—H2'	115.8
O3—C5—N2	123.6 (2)	C22'—C03'—C24'	120.1 (3)
O3—C5—C6	121.8 (2)	C22'—C03'—H23'	120.0
N2—C5—C6	114.6 (2)	C24'—C03'—H23'	119.9
C5—C6—H6A	109.5	O3'—C5'—N2'	123.8 (2)
C5—C6—H6B	109.6	O3'—C5'—C6'	122.4 (2)
H6A—C6—H6B	109.5	N2'—C5'—C6'	113.8 (2)
C5—C6—H6C	109.3	C5'—C6'—H6'1	109.4
H6A—C6—H6C	109.5	C5'—C6'—H6'2	109.6
H6B—C6—H6C	109.5	H6'1—C6'—H6'2	109.5
C16—C11—C12	118.8 (2)	C5'—C6'—H6'3	109.5
C16—C11—P1	121.12 (17)	H6'1—C6'—H6'3	109.5
C12—C11—P1	120.04 (17)	H6'2—C6'—H6'3	109.5
C13—C12—C11	119.7 (2)	C12'—C11'—C16'	118.7 (2)
C13—C12—H12	120.1	C12'—C11'—P1'	121.96 (16)
C11—C12—H12	120.2	C16'—C11'—P1'	119.25 (16)
C12—C13—C14	120.7 (3)	C11'—C12'—C13'	120.5 (2)
C12—C13—H13	119.7	C11'—C12'—H12'	119.7
C14—C13—H13	119.6	C13'—C12'—H12'	119.8
C15—C14—C13	119.9 (2)	C14'—C13'—C12'	120.1 (2)
C15—C14—H14	120.0	C14'—C13'—H13'	119.9
C13—C14—H14	120.0	C12'—C13'—H13'	120.0
C14—C15—C16	119.8 (2)	C15'—C14'—C13'	120.1 (2)
C14—C15—H15	120.2	C15'—C14'—H14'	120.0
C16—C15—H15	120.0	C13'—C14'—H14'	119.9
C15—C16—C11	121.0 (2)	C14'—C15'—C16'	120.4 (2)
C15—C16—H16	119.5	C14'—C15'—H15'	119.7
C11—C16—H16	119.4	C16'—C15'—H15'	119.9
C22—C21—C26	118.2 (2)	C15'—C16'—C11'	120.2 (2)
C22—C21—P1	122.05 (18)	C15'—C16'—H16'	119.9
C26—C21—P1	119.78 (17)	C11'—C16'—H16'	119.9
C23—C22—C21	120.7 (2)	C26'—C21'—C22'	119.2 (2)
C23—C22—H22	119.6	C26'—C21'—P1'	118.85 (18)
C21—C22—H22	119.7	C22'—C21'—P1'	121.92 (19)
C24—C23—C22	120.6 (3)	C03'—C22'—C21'	119.8 (3)
C24—C23—H23	119.7	C03'—C22'—H22'	120.1
C22—C23—H23	119.7	C21'—C22'—H22'	120.1
C23—C24—C25	119.2 (3)	C25'—C24'—C03'	120.3 (3)
C23—C24—H24	120.4	C25'—C24'—H24'	119.8
C25—C24—H24	120.4	C03'—C24'—H24'	119.9
C24—C25—C26	121.2 (3)	C24'—C25'—C26'	120.3 (3)
C24—C25—H25	119.4	C24'—C25'—H25'	119.8
C26—C25—H25	119.4	C26'—C25'—H25'	119.8
C25—C26—C21	120.1 (2)	C21'—C26'—C25'	120.2 (3)
C25—C26—H26	119.9	C21'—C26'—H26'	119.8
C21—C26—H26	119.9	C25'—C26'—H26'	119.9
C32—C31—C36	119.3 (2)	C32'—C31'—C36'	118.8 (2)
C32—C31—P1	118.03 (16)	C32'—C31'—P1'	117.74 (19)

supplementary materials

C36—C31—P1	122.53 (17)	C36'—C31'—P1'	123.3 (2)
C33—C32—C31	120.2 (2)	C31'—C32'—C33'	120.1 (3)
C33—C32—H32	119.9	C31'—C32'—H32'	120.0
C31—C32—H32	119.8	C33'—C32'—H32'	119.9
C34—C33—C32	119.8 (2)	C34'—C33'—C32'	120.7 (3)
C34—C33—H33	120.1	C34'—C33'—H33'	119.6
C32—C33—H33	120.1	C32'—C33'—H33'	119.7
C35—C34—C33	120.3 (2)	C33'—C34'—C35'	120.3 (3)
C35—C34—H34	119.9	C33'—C34'—H34'	119.9
C33—C34—H34	119.9	C35'—C34'—H34'	119.8
C34—C35—C36	120.5 (2)	C34'—C35'—C36'	120.1 (3)
C34—C35—H35	119.7	C34'—C35'—H35'	119.9
C36—C35—H35	119.8	C36'—C35'—H35'	120.0
C31—C36—C35	119.8 (2)	C31'—C36'—C35'	120.1 (3)
C31—C36—H36	120.1	C31'—C36'—H36'	120.0
C35—C36—H36	120.1	C35'—C36'—H36'	119.9
C46—C41—C42	119.3 (2)	C42'—C41'—C46'	119.7 (2)
C46—C41—S1	118.67 (16)	C42'—C41'—S1'	121.85 (16)
C42—C41—S1	122.01 (16)	C46'—C41'—S1'	118.32 (16)
C43—C42—C41	120.3 (2)	C41'—C42'—C43'	120.91 (19)
C43—C42—H42	119.8	C41'—C42'—H42'	119.5
C41—C42—H42	119.8	C43'—C42'—H42'	119.5
C42—C43—C44	120.4 (2)	C44'—C43'—C42'	119.9 (2)
C42—C43—H43	119.8	C44'—C43'—H43'	120.0
C44—C43—H43	119.8	C42'—C43'—H43'	120.1
C45—C44—C43	119.3 (2)	C43'—C44'—C45'	118.9 (2)
C45—C44—N2	123.3 (2)	C43'—C44'—N2'	124.32 (19)
C43—C44—N2	117.41 (19)	C45'—C44'—N2'	116.72 (19)
C44—C45—C46	119.9 (2)	C46'—C45'—C44'	121.1 (2)
C44—C45—H45	120.0	C46'—C45'—H45'	119.5
C46—C45—H45	120.1	C44'—C45'—H45'	119.4
C41—C46—C45	120.8 (2)	C45'—C46'—C41'	119.5 (2)
C41—C46—H46	119.6	C45'—C46'—H46'	120.3
C45—C46—H46	119.6	C41'—C46'—H46'	120.2

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N2-H2\cdots O2^{ii}$	0.86	2.11	2.966 (2)	173.
$N2'-H2'\cdots O2^{ii}$	0.86	2.11	2.961 (2)	174.

Symmetry codes: (i) $-x+1/2, y-1/2, -z+1/2$; (ii) $-x+3/2, y+1/2, -z+1/2$.

Fig. 1




Fig. 2




Fig. 3




Fig. 4

