

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4,4'-Bipyridine–2,2'-(1,2-phenylene-dioxy)diacetic acid–water (1/1/1)

Feng-Quan Bu,^a Ying Wu,^b Ling Ye^{b*} and Peng Yan^c

^aSchool of Pharmaceutical Science, Jilin University, Changchun 130012, People's Republic of China, ^bState Key Laboratory for Supramolecular Structure and Materials, Jilin University, Changchun 130012, People's Republic of China, and ^cDepartment of Chemistry and Chemical Engineering, Jinzhong College, Jinzhong 030600, People's Republic of China

Correspondence e-mail: yeling@jlu.edu.cn


Received 7 April 2010; accepted 15 April 2010

Key indicators: single-crystal X-ray study; $T = 291$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.039; wR factor = 0.113; data-to-parameter ratio = 16.4.

In the title 1:1:1 adduct, $\text{C}_{10}\text{H}_8\text{N}_2 \cdot \text{C}_{10}\text{H}_{10}\text{O}_6 \cdot \text{H}_2\text{O}$, the dihedral angle between the rings of the 4,4'-bipyridine molecule is $10.981(8)^\circ$. In the crystal, $\text{O}-\text{H} \cdots \text{O}$ and $\text{O}-\text{H} \cdots \text{N}$ hydrogen bonds link the molecules into a zigzag chain structure.

Related literature

For the synthesis of 1,2-phenylenedi(oxyacetic acid), see: Mirci (1990). For related structures, see: Soleimannejad *et al.* (2009); Yu *et al.* (2006). For hydrogen-bonding motifs, see: Etter *et al.* (1990); Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{10}\text{H}_8\text{N}_2 \cdot \text{C}_{10}\text{H}_{10}\text{O}_6 \cdot \text{H}_2\text{O}$
 $M_r = 400.38$
 Monoclinic, $P2_1/c$
 $a = 11.566(5)$ Å

$b = 9.712(5)$ Å
 $c = 16.810(7)$ Å
 $\beta = 90.81(2)^\circ$
 $V = 1888.2(15)$ Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.11$ mm⁻¹

$T = 291$ K
 $0.20 \times 0.18 \times 0.12$ mm

Data collection

Rigaku R-Axis RAPID
 diffractometer
 Absorption correction: multi-scan
 (ABSCOR; Higashi, 1995)
 $T_{\min} = 0.979$, $T_{\max} = 0.987$

18105 measured reflections
 4306 independent reflections
 2654 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.113$
 $S = 0.97$
 4306 reflections

262 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.17$ e Å⁻³
 $\Delta\rho_{\min} = -0.15$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{O3}-\text{H23} \cdots \text{N2}^i$	0.85	1.76	2.6051 (19)	173
$\text{O6}-\text{H24} \cdots \text{N1}$	0.85	1.74	2.5871 (19)	176
$\text{O7}-\text{H21} \cdots \text{O5}$	0.85	2.07	2.8817 (19)	160
$\text{O7}-\text{H22} \cdots \text{O2}$	0.85	1.97	2.7826 (17)	161

Symmetry code: (i) $x - 1, -y - \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalClear* (Rigaku/MSK, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-III* (Burnett & Johnson, 1996), *ORTEP-3 for Windows* (Farrugia, 1997) and *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors thank the National Natural Science Foundation of China (50733002-01) and Jilin University for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2556).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
 Burnett, M. N. & Johnson, C. K. (1996). *ORTEP-III*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
 Etter, M. C., MacDonald, J. C. & Bernstein, J. (1990). *Acta Cryst.* **B46**, 256–262.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
 Mirci, L. E. (1990). Rom. Patent No. 07 43 205.
 Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
 Rigaku/MSK (2002). *CrystalClear*. Rigaku/MSK Inc., The Woodlands, Texas, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Soleimannejad, J., Aghabozorg, H., Najafi, S., Nasibipour, M. & Attar Gharamaleki, J. (2009). *Acta Cryst.* **E65**, o532–o533.
 Yu, B., Wang, X.-Q., Wang, R.-J., Shen, G.-Q. & Shen, D.-Z. (2006). *Acta Cryst.* **E62**, o2757–o2758.

supplementary materials

Acta Cryst. (2010). E66, o1129 [doi:10.1107/S1600536810013905]

4,4'-Bipyridine-2,2'-(1,2-phenylenedioxy)diacetic acid-water (1/1/1)

F.-Q. Bu, Y. Wu, L. Ye and P. Yan

Comment

Flexible carboxylic acid and 4,4'-bipyridine are often used for constructing the high dimensional structure with intriguing topology. In this paper, we report the title compound, contained flexible 1,2-phenylenedi(oxyacetic acid), 4,4'-bipyridine and water molecules, which forms a one-dimensional hydrogen bonding chain structure.

The O—H...O hydrogen bonds link the 1,2-phenylenedi(oxyacetic acid) and water molecules into a 13-membered hydrogen bonding ring, with a unitary graph-set descriptor $N1 = R^2_2(13)$ (Etter *et al.*, 1990; Bernstein *et al.*, 1995) (Figure 1, Table 1). And then, 4,4'-bipyridine molecules bridge these hydrogen bonding rings into a one-dimensional zigzag chain through the O—H...N hydrogen bonds (Table 1, Figure 2). Similar hydrogen bonding interactions involving the 4,4'-bipyridine have been frequently observed (Soleimannejad *et al.*, 2009; Yu *et al.*, 2006).

Experimental

1,2-phenylenedi(oxyacetic acid) was prepared by the reaction of chloroacetic acid with dihydroxybenzene (Mirci, 1990). All other chemicals were analytical grade reagents and used without further purification. 1,2-phenylenedi(oxyacetic acid) (0.40 g, 2 mmol) and 4,4'-bipyridine (0.31 g, 2 mmol) were dissolved in mixed solution of water (5 ml) and ethanol (5 ml). Colorless rod crystals of title compound were obtained after several days in room temperature.

Refinement

H atoms bound to C atoms were placed in calculated positions and treated as riding on their parent atoms, with C—H = 0.97 Å (methylene), C—H = 0.93 Å (aromatic) and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. H atoms O atoms were initially located in a difference Fourier map and treated as riding on their parent atoms, with O—H = 0.85 Å, $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{N/O})$.

Figures




Fig. 1. The molecular structure of the title compound with the atom labeling scheme, Ellipsoids are drawn at the 50% probability level. H atoms are represented as small spheres of arbitrary radii. Hydrogen bonds are shown as dashed lines.



Fig. 2. A partial packing view, showing the three-dimensional hydrogen bonding supra-molecular network. Hydrogen bonds are shown as dashed lines. H atoms not involved in hydrogen bondings have been omitted for clarity.

4,4'-Bipyridine-2,2'-(1,2-phenylenedioxy)diacetic acid-water (1/1/1)

Crystal data

$C_{10}H_8N_2 \cdot C_{10}H_{10}O_6 \cdot H_2O$	$F(000) = 840$
$M_r = 400.38$	$D_x = 1.408 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 11161 reflections
$a = 11.566 (5) \text{ \AA}$	$\theta = 3.0\text{--}27.5^\circ$
$b = 9.712 (5) \text{ \AA}$	$\mu = 0.11 \text{ mm}^{-1}$
$c = 16.810 (7) \text{ \AA}$	$T = 291 \text{ K}$
$\beta = 90.81 (2)^\circ$	Rod, colorless
$V = 1888.2 (15) \text{ \AA}^3$	$0.20 \times 0.18 \times 0.12 \text{ mm}$
$Z = 4$	

Data collection

Rigaku R-Axis RAPID diffractometer	4306 independent reflections
Radiation source: fine-focus sealed tube graphite	2654 reflections with $I > 2\sigma(I)$
ω scan	$R_{\text{int}} = 0.030$
Absorption correction: multi-scan (ABSCOR; Higashi, 1995)	$\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 3.0^\circ$
$T_{\text{min}} = 0.979$, $T_{\text{max}} = 0.987$	$h = -15 \rightarrow 14$
18105 measured reflections	$k = -12 \rightarrow 12$
	$l = -21 \rightarrow 21$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.039$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.113$	H-atom parameters constrained
$S = 0.97$	$w = 1/[\sigma^2(F_o^2) + (0.0647P)^2]$
4306 reflections	where $P = (F_o^2 + 2F_c^2)/3$
262 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.17 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.15 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds

in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.60888 (10)	0.06435 (13)	0.44070 (7)	0.0574 (3)
C1	0.59235 (13)	-0.04189 (17)	0.39244 (9)	0.0616 (4)
H1	0.5467	-0.0293	0.3470	0.074*
C2	0.63933 (12)	-0.16976 (16)	0.40627 (8)	0.0588 (4)
H2	0.6265	-0.2406	0.3700	0.071*
C3	0.70588 (11)	-0.19349 (14)	0.47429 (8)	0.0462 (3)
C4	0.72238 (12)	-0.08127 (15)	0.52468 (9)	0.0574 (4)
H4	0.7662	-0.0910	0.5712	0.069*
C5	0.67402 (13)	0.04364 (16)	0.50574 (9)	0.0613 (4)
H5	0.6873	0.1175	0.5399	0.074*
C6	0.75714 (11)	-0.33068 (14)	0.49263 (8)	0.0460 (3)
C7	0.72463 (14)	-0.44768 (16)	0.45093 (9)	0.0662 (4)
H7	0.6697	-0.4416	0.4101	0.079*
C8	0.77316 (14)	-0.57246 (17)	0.46966 (10)	0.0672 (4)
H8	0.7498	-0.6493	0.4406	0.081*
C9	0.88302 (13)	-0.47824 (16)	0.56821 (9)	0.0609 (4)
H9	0.9375	-0.4879	0.6091	0.073*
C10	0.83819 (12)	-0.34962 (15)	0.55282 (9)	0.0579 (4)
H10	0.8626	-0.2748	0.5832	0.069*
C11	0.27722 (11)	0.59624 (14)	0.26783 (7)	0.0458 (3)
C12	0.32396 (13)	0.71838 (15)	0.29545 (8)	0.0554 (4)
H12	0.3895	0.7167	0.3282	0.066*
C13	0.27396 (13)	0.84376 (16)	0.27477 (9)	0.0597 (4)
H13	0.3061	0.9256	0.2935	0.072*
C14	0.17787 (14)	0.84657 (15)	0.22704 (9)	0.0594 (4)
H14	0.1442	0.9305	0.2135	0.071*
C15	0.12991 (12)	0.72505 (14)	0.19859 (9)	0.0561 (4)
H15	0.0644	0.7282	0.1659	0.067*
C16	0.17855 (11)	0.59898 (13)	0.21839 (8)	0.0460 (3)
C17	0.04541 (11)	0.47710 (14)	0.13642 (8)	0.0495 (3)
H17A	0.0694	0.5276	0.0897	0.059*
H17B	-0.0208	0.5239	0.1587	0.059*
C18	0.01249 (11)	0.33352 (14)	0.11347 (8)	0.0479 (3)
C19	0.41863 (12)	0.46231 (15)	0.33609 (8)	0.0542 (4)
H19A	0.4001	0.5044	0.3867	0.065*
H19B	0.4818	0.5135	0.3130	0.065*
C20	0.45447 (12)	0.31491 (16)	0.34906 (9)	0.0548 (4)
N2	0.85154 (10)	-0.58951 (13)	0.52696 (7)	0.0564 (3)

supplementary materials

O1	0.13736 (8)	0.47386 (9)	0.19334 (6)	0.0545 (3)
O2	0.05129 (9)	0.23016 (10)	0.14397 (6)	0.0597 (3)
O3	-0.06491 (9)	0.33379 (10)	0.05595 (6)	0.0651 (3)
H23	-0.0867	0.2512	0.0481	0.098*
O4	0.32100 (8)	0.46709 (9)	0.28462 (6)	0.0531 (3)
O5	0.42135 (9)	0.21927 (11)	0.30849 (7)	0.0710 (3)
O6	0.52702 (9)	0.30628 (11)	0.40864 (6)	0.0709 (3)
H24	0.5506	0.2251	0.4185	0.106*
O7	0.17765 (9)	0.17334 (11)	0.28219 (6)	0.0712 (3)
H21	0.2472	0.2012	0.2804	0.107*
H22	0.1466	0.2093	0.2410	0.107*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0485 (7)	0.0614 (8)	0.0619 (7)	0.0020 (6)	-0.0057 (6)	0.0183 (6)
C1	0.0563 (9)	0.0752 (11)	0.0530 (9)	0.0049 (8)	-0.0126 (7)	0.0131 (8)
C2	0.0600 (9)	0.0681 (10)	0.0477 (8)	0.0021 (7)	-0.0126 (7)	0.0027 (7)
C3	0.0390 (7)	0.0545 (8)	0.0450 (7)	-0.0045 (6)	-0.0039 (6)	0.0072 (6)
C4	0.0596 (9)	0.0545 (9)	0.0576 (9)	-0.0024 (7)	-0.0206 (7)	0.0064 (7)
C5	0.0631 (9)	0.0544 (9)	0.0659 (10)	-0.0012 (7)	-0.0148 (8)	0.0071 (7)
C6	0.0416 (7)	0.0524 (8)	0.0438 (7)	-0.0029 (6)	-0.0028 (6)	0.0038 (6)
C7	0.0703 (10)	0.0606 (10)	0.0669 (10)	0.0008 (8)	-0.0296 (8)	-0.0013 (7)
C8	0.0746 (11)	0.0555 (9)	0.0707 (10)	-0.0005 (8)	-0.0216 (9)	-0.0062 (8)
C9	0.0601 (9)	0.0614 (10)	0.0605 (9)	0.0063 (7)	-0.0208 (7)	-0.0002 (7)
C10	0.0591 (8)	0.0562 (9)	0.0578 (9)	0.0015 (7)	-0.0184 (7)	-0.0038 (7)
C11	0.0460 (7)	0.0424 (7)	0.0491 (8)	0.0020 (6)	-0.0019 (6)	0.0038 (6)
C12	0.0569 (8)	0.0524 (8)	0.0564 (9)	-0.0040 (7)	-0.0123 (7)	-0.0008 (7)
C13	0.0711 (10)	0.0460 (8)	0.0619 (9)	-0.0055 (7)	-0.0024 (8)	-0.0063 (7)
C14	0.0701 (9)	0.0429 (8)	0.0653 (9)	0.0085 (7)	-0.0010 (8)	-0.0015 (7)
C15	0.0536 (8)	0.0473 (8)	0.0673 (9)	0.0062 (6)	-0.0098 (7)	-0.0008 (7)
C16	0.0434 (7)	0.0412 (7)	0.0532 (8)	0.0010 (6)	-0.0036 (6)	-0.0006 (6)
C17	0.0436 (7)	0.0486 (8)	0.0559 (8)	0.0043 (6)	-0.0107 (6)	0.0010 (6)
C18	0.0419 (7)	0.0512 (8)	0.0504 (8)	0.0016 (6)	-0.0037 (6)	0.0006 (6)
C19	0.0486 (8)	0.0563 (9)	0.0572 (8)	-0.0028 (6)	-0.0138 (7)	0.0076 (7)
C20	0.0420 (7)	0.0605 (9)	0.0615 (9)	-0.0020 (7)	-0.0072 (7)	0.0113 (7)
N2	0.0542 (7)	0.0556 (7)	0.0592 (7)	0.0048 (6)	-0.0051 (6)	0.0036 (6)
O1	0.0508 (5)	0.0404 (5)	0.0718 (6)	0.0025 (4)	-0.0210 (5)	-0.0003 (4)
O2	0.0639 (6)	0.0475 (6)	0.0672 (7)	0.0036 (5)	-0.0183 (5)	0.0005 (5)
O3	0.0661 (6)	0.0540 (6)	0.0743 (7)	-0.0046 (5)	-0.0293 (6)	0.0030 (5)
O4	0.0492 (5)	0.0462 (6)	0.0635 (6)	0.0002 (4)	-0.0167 (4)	0.0052 (4)
O5	0.0635 (7)	0.0599 (7)	0.0888 (8)	0.0060 (5)	-0.0230 (6)	-0.0003 (6)
O6	0.0665 (7)	0.0639 (7)	0.0812 (8)	0.0023 (5)	-0.0319 (6)	0.0154 (6)
O7	0.0797 (7)	0.0581 (7)	0.0750 (7)	-0.0155 (5)	-0.0260 (6)	0.0135 (5)

Geometric parameters (\AA , $^\circ$)

N1—C1	1.325 (2)	C12—H12	0.9300
N1—C5	1.3342 (18)	C13—C14	1.362 (2)

C1—C2	1.374 (2)	C13—H13	0.9300
C1—H1	0.9300	C14—C15	1.386 (2)
C2—C3	1.3885 (18)	C14—H14	0.9300
C2—H2	0.9300	C15—C16	1.3861 (19)
C3—C4	1.392 (2)	C15—H15	0.9300
C3—C6	1.489 (2)	C16—O1	1.3695 (16)
C4—C5	1.371 (2)	C17—O1	1.4208 (15)
C4—H4	0.9300	C17—C18	1.495 (2)
C5—H5	0.9300	C17—H17A	0.9700
C6—C10	1.3818 (18)	C17—H17B	0.9700
C6—C7	1.384 (2)	C18—O2	1.2105 (16)
C7—C8	1.370 (2)	C18—O3	1.3083 (16)
C7—H7	0.9300	C19—O4	1.4136 (15)
C8—N2	1.3237 (18)	C19—C20	1.505 (2)
C8—H8	0.9300	C19—H19A	0.9700
C9—N2	1.3320 (19)	C19—H19B	0.9700
C9—C10	1.376 (2)	C20—O5	1.2113 (18)
C9—H9	0.9300	C20—O6	1.3002 (16)
C10—H10	0.9300	O3—H23	0.8499
C11—O4	1.3803 (17)	O6—H24	0.8500
C11—C12	1.3812 (19)	O7—H21	0.8499
C11—C16	1.4025 (18)	O7—H22	0.8501
C12—C13	1.390 (2)		
C1—N1—C5	117.34 (13)	C14—C13—C12	119.87 (13)
N1—C1—C2	123.17 (12)	C14—C13—H13	120.1
N1—C1—H1	118.4	C12—C13—H13	120.1
C2—C1—H1	118.4	C13—C14—C15	120.36 (13)
C1—C2—C3	120.17 (14)	C13—C14—H14	119.8
C1—C2—H2	119.9	C15—C14—H14	119.8
C3—C2—H2	119.9	C16—C15—C14	120.67 (13)
C2—C3—C4	116.16 (13)	C16—C15—H15	119.7
C2—C3—C6	122.24 (13)	C14—C15—H15	119.7
C4—C3—C6	121.60 (11)	O1—C16—C15	124.87 (11)
C5—C4—C3	119.96 (12)	O1—C16—C11	116.23 (11)
C5—C4—H4	120.0	C15—C16—C11	118.90 (12)
C3—C4—H4	120.0	O1—C17—C18	109.80 (10)
N1—C5—C4	123.18 (14)	O1—C17—H17A	109.7
N1—C5—H5	118.4	C18—C17—H17A	109.7
C4—C5—H5	118.4	O1—C17—H17B	109.7
C10—C6—C7	115.98 (13)	C18—C17—H17B	109.7
C10—C6—C3	122.33 (12)	H17A—C17—H17B	108.2
C7—C6—C3	121.68 (12)	O2—C18—O3	124.09 (13)
C8—C7—C6	120.14 (13)	O2—C18—C17	124.94 (12)
C8—C7—H7	119.9	O3—C18—C17	110.98 (11)
C6—C7—H7	119.9	O4—C19—C20	109.62 (11)
N2—C8—C7	123.49 (14)	O4—C19—H19A	109.7
N2—C8—H8	118.3	C20—C19—H19A	109.7
C7—C8—H8	118.3	O4—C19—H19B	109.7
N2—C9—C10	122.64 (12)	C20—C19—H19B	109.7

supplementary materials

N2—C9—H9	118.7	H19A—C19—H19B	108.2
C10—C9—H9	118.7	O5—C20—O6	125.43 (14)
C9—C10—C6	120.57 (13)	O5—C20—C19	124.30 (12)
C9—C10—H10	119.7	O6—C20—C19	110.26 (13)
C6—C10—H10	119.7	C8—N2—C9	117.18 (13)
O4—C11—C12	124.82 (11)	C16—O1—C17	116.18 (10)
O4—C11—C16	115.56 (11)	C18—O3—H23	108.1
C12—C11—C16	119.61 (11)	C11—O4—C19	116.26 (10)
C11—C12—C13	120.59 (12)	C20—O6—H24	114.3
C11—C12—H12	119.7	H21—O7—H22	103.2
C13—C12—H12	119.7		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O3—H23 \cdots N2 ⁱ	0.85	1.76	2.6051 (19)	173
O6—H24 \cdots N1	0.85	1.74	2.5871 (19)	176
O7—H21 \cdots O5	0.85	2.07	2.8817 (19)	160
O7—H22 \cdots O2	0.85	1.97	2.7826 (17)	161

Symmetry codes: (i) $x-1, -y-1/2, z-1/2$.

Fig. 1




Fig. 2

