

3,3'-Ethylenebis(3,4-dihydro-6-chloro-2H-1,3-benzoxazine)

Augusto Rivera,^{a*} Jicli José Rojas,^a Jaime Ríos-Motta,^a Michal Dušek^b and Karla Fejfarová^b

^aDepartamento de Química, Universidad Nacional de Colombia, Bogotá, AA 14490, Colombia, and ^bInstitute of Physics, Na Slovance 2, 182 21 Praha 8, Czech Republic
Correspondence e-mail: ariverau@unal.edu.co


Received 29 March 2010; accepted 18 April 2010

Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.030; wR factor = 0.103; data-to-parameter ratio = 13.2.

The asymmetric unit of the title compound, $\text{C}_{18}\text{H}_{18}\text{Cl}_2\text{N}_2\text{O}_2$, contains one half of an independent molecule, the other half being generated *via* a centre of inversion at the molecular centroid. In the crystal structure, molecular chains are formed through non-classical $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds between an axial H atom of the oxazine ring and the O atom of a neighbouring molecule.

Related literature

For the synthesis, see: Rivera *et al.* (1989). For related structures, see: Rivera *et al.* (1986); Huerta *et al.* (2006); Chen & Wu (2007); Ranjith *et al.* (2009). For uses of benzoxazines in polymer science, see Yaggi *et al.* (2009). For the biological activity of bis-benzoxazine compounds, see: Billmann & Dorman (1963); Heinisch *et al.* (2002).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{18}\text{Cl}_2\text{N}_2\text{O}_2$	$V = 1629.03$ (12) Å ³
$M_r = 365.3$	$Z = 4$
Monoclinic, $C2/c$	Cu $K\alpha$ radiation
$a = 18.9920$ (5) Å	$\mu = 3.70$ mm ⁻¹
$b = 5.8884$ (2) Å	$T = 120$ K
$c = 17.8813$ (5) Å	$0.30 \times 0.19 \times 0.12$ mm
$\beta = 125.449$ (4)°	

Data collection

Oxford Diffraction Xcalibur diffractometer with an Atlas (Gemini ultra Cu) detector	expressions derived by Clark & Reid (1995)]
Absorption correction: analytical [CrysAlis PRO (Oxford Diffraction, 2009), using a multifaceted crystal model based on	$T_{\min} = 0.593$, $T_{\max} = 0.787$
	12716 measured reflections
	1442 independent reflections
	1344 reflections with $I > 3\sigma(I)$
	$R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$	109 parameters
$wR(F^2) = 0.103$	H-atom parameters constrained
$S = 2.26$	$\Delta\rho_{\max} = 0.25$ e Å ⁻³
1442 reflections	$\Delta\rho_{\min} = -0.25$ e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}2-\text{H}2\text{B}\cdots\text{O}1^i$	0.96	2.56	3.369 (2)	142

Symmetry code: (i) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2009); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SIR2002* (Burla *et al.*, 2003); program(s) used to refine structure: *JANA2006* (Petříček *et al.*, 2006); molecular graphics: *DIAMOND* (Brandenburg & Putz, 2005); software used to prepare material for publication: *JANA2006*.

We acknowledge the Dirección de Investigaciones Sede Bogotá (DIB) of the Universidad Nacional de Colombia, the Institutional research plan No. AVOZ10100521 of the Institute of Physics and the Praemium Academiae project of the Academy of Sciences (ASCR) for financial support of this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2292).

References

- Billmann, J. H. & Dorman, L. C. (1963). *J. Med. Chem.* **6**, 701–708.
- Brandenburg, K. & Putz, H. (2005). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Burla, M. C., Camalli, M., Carrozzini, B., Cascarano, G. L., Giacovazzo, C., Polidori, G. & Spagna, R. (2003). *J. Appl. Cryst.* **36**, 1103.
- Chen, X.-L. & Wu, M.-H. (2007). *Acta Cryst.* **E63**, o3684.
- Clark, R. C. & Reid, J. S. (1995). *Acta Cryst.* **A51**, 887–897.
- Heinisch, L., Wittmann, S., Stoiber, T., Berg, A., Ankel-Fuchs, D. & Mollmann, U. (2002). *J. Med. Chem.* **45**, 3032–3039.
- Huerta, R., Toscano, R. A. & Castillo, I. (2006). *Acta Cryst.* **E62**, o2938–o2940.
- Oxford Diffraction (2009). *CrysAlis CCD*, *CrysAlis RED* and *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
- Petříček, V., Dušek, M. & Palatinus, L. (2006). *JANA2006*. Institute of Physics, Praha, Czech Republic.
- Ranjith, S., Thenmozhi, S., Manikannan, R., Muthusubramanian, S. & Subbiahpanidi, A. (2009). *Acta Cryst.* **E65**, o581.
- Rivera, A., Aguilar, Z., Clavijo, D. & Joseph-Nathan, P. (1989). *Anal. Quim.* **85**, 9–10.
- Rivera, A., Ospina, E., Sanchez, A. & Joseph-Nathan, P. (1986). *Heterocycles*, **24**, 2507–2510.
- Yaggi, Y., Kiskan, B. & Ghosh, N. N. (2009). *J. Polym. Sci. Part A Polym. Chem.* **47**, 5565–5576.

supplementary materials

Acta Cryst. (2010). E66, o1134 [doi:10.1107/S1600536810014248]

3,3'-Ethylenebis(3,4-dihydro-6-chloro-2H-1,3-benzoxazine)

A. Rivera, J. J. Rojas, J. Ríos-Motta, M. Dusek and K. Fejfarová

Comment

1,3-Benzoxazines are heterocyclic compound obtained from condensation between phenols, formaldehyde and a primary amine. Applications of these compounds are in polymeric and pharmacological fields. Recently the structure of these compounds has attracted much attention see Huerta *et al.* (2006); Chen & Wu (2007); Ranjith *et al.* (2009). During our investigations, a series of bis-benzoxazines were prepared by reaction of phenols, formaldehyde and ethylenediamine (Rivera *et al.*, 1986). However, the crystallization of these compounds was difficult and led to crystals of bad quality. In the present work, the single crystals of the title compound were finally successfully prepared and its crystal structure has been determined herein.

The molecule contains two 1,3-benzoxazine units linked by an ethylene bridge. The asymmetric unit of the title compound $C_{18}H_{18}Cl_2N_2O_2$, contains one-half of the formula unit; a centre of inversion is located at the mid-point of the central C1—C1ⁱ bond (see Fig. 1). Both oxazine rings are in cyclohexene-like conformations with normal bond distances and angles, and their values were found in good agreement with the corresponding values in the related structures reported by Huerta *et al.* (2006), Chen & Wu (2007) and Ranjith *et al.* (2009). In the crystal structure, molecules are linked *via* C2—H2B \cdots O1 weak hydrogen bonds forming a molecular slab (see Fig 2a,b). The bond involves axial-hydrogen of oxazine ring and the oxygen atom of a neighbor molecule.

There is also possibility for very weak intermolecular interaction between the hydrogen H2A and the aromatic ring C3,C4,C6, C7, C8, C9, with the distance between H2A and the centre of the ring of 2.99 Å.

Experimental

Under vigorous stirring a mixture of ethylenediamine (0.34 ml, 5 mmol) and *p*-chlorophenol (1.3 g 10 mmol) was dissolved in dioxane (10 ml) and (1.5 ml, 20 mmol) was slowly added. Stirring was continued for 4 h at rt until a precipitate appeared. The solid was filtered off and washed with water (1.83 g, 92%). Recrystallization from ethanol gave a white solid.

Refinement

All hydrogen atoms were discernible in difference Fourier maps and could be refined to reasonable geometry. According to common practice H atoms attached to C atoms were nevertheless kept in ideal positions during the refinement. The isotropic atomic displacement parameters of hydrogen atoms were evaluated as $1.2 \cdot U_{eq}$ of the parent atom.

Figures




Fig. 1. The molecular structure of title compound, showing the atomic numbering scheme with atomic displacement ellipsoids drawn at the 50%.




Fig. 2. Perspective views of the crystal packing showing hydrogen-bonded interactions (dashed lines).

3,3'-Ethylenebis(3,4-dihydro-6-chloro-2H-1,3-benzoxazine)

Crystal data

$C_{18}H_{18}Cl_2N_2O_2$

$M_r = 365.3$

Monoclinic, $C2/c$

Hall symbol: $-C 2yc$

$a = 18.9920 (5) \text{ \AA}$

$b = 5.8884 (2) \text{ \AA}$

$c = 17.8813 (5) \text{ \AA}$

$\beta = 125.449 (4)^\circ$

$V = 1629.03 (12) \text{ \AA}^3$

$Z = 4$

$F(000) = 760$

$D_x = 1.489 \text{ Mg m}^{-3}$

Cu $K\alpha$ radiation, $\lambda = 1.54184 \text{ \AA}$

Cell parameters from 10117 reflections

$\theta = 3.0\text{--}66.8^\circ$

$\mu = 3.70 \text{ mm}^{-1}$

$T = 120 \text{ K}$

Prism, colorless

$0.30 \times 0.19 \times 0.12 \text{ mm}$

Data collection

Oxford Diffraction Xcalibur

diffractometer with an Atlas (Gemini ultra Cu) detector

1442 independent reflections

Radiation source: X-ray tube mirror

1344 reflections with $I > 3\sigma(I)$

$R_{\text{int}} = 0.024$

Detector resolution: $10.3784 \text{ pixels mm}^{-1}$

$\theta_{\text{max}} = 75.1^\circ$, $\theta_{\text{min}} = 5.4^\circ$

Rotation method data acquisition using ω scans

$h = -22 \rightarrow 22$

Absorption correction: analytical

[CrysAlis Pro (Oxford Diffraction, 2009), using a multifaceted crystal model based on expressions derived by Clark & Reid (1995)]

$k = -7 \rightarrow 6$

$T_{\text{min}} = 0.593$, $T_{\text{max}} = 0.787$

$l = -19 \rightarrow 20$

12716 measured reflections

Refinement

Refinement on F^2	36 constraints
$R[F^2 > 2\sigma(F^2)] = 0.030$	H-atom parameters constrained
$wR(F^2) = 0.103$	Weighting scheme based on measured s.u.'s $w = 1/[\sigma^2(I) + 0.0016I^2]$
$S = 2.26$	$(\Delta/\sigma)_{\max} = 0.014$
1442 reflections	$\Delta\rho_{\max} = 0.25 \text{ e } \text{\AA}^{-3}$
109 parameters	$\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$
0 restraints	

Special details

Experimental. CrysAlisPro (Oxford Diffraction Ltd., Version 1.171.33.51 (release 27-10-2009 CrysAlis171 .NET) Analytical numeric absorption correction using a multifaceted crystal model based on expressions derived by Clark & Reid (1995)

Refinement. The refinement was carried out against all reflections. The conventional R -factor is always based on F . The goodness of fit as well as the weighted R -factor are based on F and F^2 for refinement carried out on F and F^2 , respectively. The threshold expression is used only for calculating R -factors etc. and it is not relevant to the choice of reflections for refinement.

The program used for refinement, Jana2006, uses the weighting scheme based on the experimental expectations, see `_refine_ls_weighting_details`, that does not force S to be one. Therefore the values of S are usually larger than the ones from the SHELX program.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.60192 (2)	-0.00670 (6)	0.93208 (3)	0.0280 (3)
O1	0.67792 (6)	0.65142 (16)	0.72490 (7)	0.0215 (5)
N1	0.60904 (7)	0.4270 (2)	0.58616 (8)	0.0180 (5)
C1	0.53100 (8)	0.5511 (3)	0.54598 (10)	0.0185 (7)
C2	0.67854 (8)	0.5625 (3)	0.64492 (10)	0.0198 (6)
C3	0.65998 (9)	0.4905 (2)	0.77097 (11)	0.0192 (7)
C4	0.62512 (8)	0.2799 (2)	0.73128 (10)	0.0190 (6)
C5	0.61011 (8)	0.2223 (2)	0.63683 (10)	0.0191 (6)
C6	0.60690 (8)	0.1294 (2)	0.78148 (10)	0.0208 (6)
C7	0.62347 (9)	0.1873 (3)	0.86889 (10)	0.0233 (7)
C8	0.65769 (9)	0.3956 (3)	0.90753 (11)	0.0261 (7)
C9	0.67601 (10)	0.5459 (3)	0.85788 (11)	0.0246 (7)
H1a	0.54087	0.707097	0.539047	0.0222*
H1b	0.509368	0.533303	0.582633	0.0222*
H2a	0.679796	0.688363	0.61157	0.0238*
H2b	0.730991	0.479648	0.668433	0.0238*
H5a	0.65391	0.118579	0.647102	0.023*
H5b	0.556851	0.140341	0.598522	0.023*
H6	0.582529	-0.016899	0.755597	0.0249*
H8	0.668857	0.438584	0.965284	0.0313*
H9	0.700365	0.691858	0.884131	0.0295*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0306 (3)	0.0293 (3)	0.0288 (3)	0.00199 (13)	0.0199 (2)	0.00560 (14)
O1	0.0192 (5)	0.0205 (6)	0.0179 (5)	-0.0041 (4)	0.0068 (4)	-0.0012 (4)
N1	0.0124 (5)	0.0177 (6)	0.0169 (6)	0.0007 (4)	0.0044 (5)	0.0005 (5)
C1	0.0141 (6)	0.0178 (7)	0.0161 (8)	0.0023 (5)	0.0045 (6)	0.0013 (6)
C2	0.0150 (6)	0.0227 (7)	0.0174 (7)	-0.0018 (5)	0.0069 (6)	-0.0001 (6)
C3	0.0137 (6)	0.0193 (8)	0.0183 (8)	0.0015 (4)	0.0057 (6)	0.0018 (5)
C4	0.0131 (6)	0.0203 (7)	0.0172 (7)	0.0028 (5)	0.0052 (5)	0.0006 (5)
C5	0.0162 (6)	0.0173 (7)	0.0176 (7)	0.0010 (5)	0.0062 (5)	0.0011 (5)
C6	0.0146 (6)	0.0193 (7)	0.0231 (8)	0.0016 (5)	0.0079 (6)	0.0007 (6)
C7	0.0194 (6)	0.0251 (8)	0.0248 (8)	0.0042 (5)	0.0125 (6)	0.0053 (6)
C8	0.0267 (7)	0.0282 (8)	0.0210 (8)	0.0026 (6)	0.0125 (6)	-0.0014 (6)
C9	0.0233 (7)	0.0217 (7)	0.0228 (8)	-0.0005 (6)	0.0100 (6)	-0.0023 (6)

Geometric parameters (\AA , $^\circ$)

C11—C7	1.813 (2)	C3—C9	1.437 (3)
O1—C2	1.529 (2)	C4—C5	1.578 (3)
O1—C3	1.421 (2)	C4—C6	1.439 (3)
N1—C1	1.4182 (18)	C5—H5a	0.96
N1—C2	1.3690 (16)	C5—H5b	0.96
N1—C5	1.501 (2)	C6—C7	1.445 (3)
C1—C1 ⁱ	1.4853 (18)	C6—H6	0.96
C1—H1a	0.96	C7—C8	1.372 (2)
C1—H1b	0.96	C8—C9	1.432 (3)
C2—H2a	0.96	C8—H8	0.96
C2—H2b	0.96	C9—H9	0.96
C3—C4	1.3907 (19)		
C2—O1—C3	116.65 (11)	C3—C4—C6	116.51 (16)
C1—N1—C2	110.35 (12)	C5—C4—C6	125.13 (12)
C1—N1—C5	111.32 (14)	N1—C5—C4	113.84 (12)
C2—N1—C5	109.60 (10)	N1—C5—H5a	109.4703
N1—C1—C1 ⁱ	106.07 (13)	N1—C5—H5b	109.4713
N1—C1—H1a	109.4706	C4—C5—H5a	109.4721
N1—C1—H1b	109.4717	C4—C5—H5b	109.4707
C1 ⁱ —C1—H1a	109.47	H5a—C5—H5b	104.7188
C1 ⁱ —C1—H1b	109.4723	C4—C6—C7	123.28 (13)
H1a—C1—H1b	112.6688	C4—C6—H6	118.3591
O1—C2—N1	112.97 (15)	C7—C6—H6	118.3615
O1—C2—H2a	109.4708	C11—C7—C6	122.62 (11)
O1—C2—H2b	109.4711	C11—C7—C8	117.49 (15)
N1—C2—H2a	109.471	C6—C7—C8	119.88 (18)
N1—C2—H2b	109.4715	C7—C8—C9	117.04 (18)
H2a—C2—H2b	105.7297	C7—C8—H8	121.4818
O1—C3—C4	120.14 (17)	C9—C8—H8	121.4814

O1—C3—C9	120.27 (12)	C3—C9—C8	123.70 (14)
C4—C3—C9	119.58 (17)	C3—C9—H9	118.1485
C3—C4—C5	118.35 (16)	C8—C9—H9	118.1475
C2—N1—C1—C1 ⁱ	150.21 (14)	C9—C3—C4—C6	0.2 (3)
C5—N1—C1—C1 ⁱ	-87.87 (15)	O1—C3—C9—C8	178.38 (17)
C3—O1—C2—N1	46.41 (18)	C4—C3—C9—C8	-0.3 (3)
C2—O1—C3—C4	-14.6 (2)	C3—C4—C5—N1	-18.4 (2)
C2—O1—C3—C9	166.66 (16)	C6—C4—C5—N1	162.77 (15)
C1—N1—C2—O1	61.58 (16)	C3—C4—C6—C7	-0.3 (3)
C5—N1—C2—O1	-61.37 (17)	C5—C4—C6—C7	178.57 (16)
C1—N1—C5—C4	-74.43 (16)	C4—C6—C7—C11	-178.83 (13)
C2—N1—C5—C4	47.92 (19)	C4—C6—C7—C8	0.5 (3)
N1—C1—C1 ⁱ —N1 ⁱ	180.00 (13)	C11—C7—C8—C9	178.84 (14)
O1—C3—C4—C5	2.6 (2)	C6—C7—C8—C9	-0.5 (3)
O1—C3—C4—C6	-178.46 (15)	C7—C8—C9—C3	0.5 (3)
C9—C3—C4—C5	-178.74 (16)		

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C2—H2B \cdots O1 ⁱⁱ	0.96	2.56	3.369 (2)	142

Symmetry codes: (ii) $-x+3/2, y-1/2, -z+3/2$.

Fig. 1




Fig. 2

