

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Methyl 12-bromodehydroabietate

Hong Gao, Zhan-Qian Song* and Shi-Bin Shang

Institute of Chemical Industry of Forest Products, Chinese Academy of Forestry, Nanjing 210042, People's Republic of China

Correspondence e-mail: songzq@hotmail.com


Received 22 March 2010; accepted 31 March 2010

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å; R factor = 0.047; wR factor = 0.128; data-to-parameter ratio = 16.1.

The title compound [systematic name: (1*R*)-methyl 6-bromo-7-isopropyl-1,4a-dimethyl-1,2,3,4,4a,9,10,10a-octahydrophenanthrene-1-carboxylate], $\text{C}_{21}\text{H}_{29}\text{BrO}_2$, was synthesized from *N*-bromosuccinimide and methyl dehydroabietate, which was prepared through an esterification reaction using dehydroabietic acid and methanol as raw materials. The three six-membered rings adopt planar (mean deviation = 0.002 Å) half-chair and chair conformations. The two cyclohexane rings form a *trans* ring junction with the two methyl groups in axial positions. The crystal structure is stabilized by weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ contacts along the *b* axis.

Related literature

For the isolation of dehydroabietic acid, see: Halbrook & Lawrence (1966). For the preparation and use of dehydroabietic acid derivatives, see: Fonseca *et al.* (2001); Pan *et al.* (2006). For the synthesis of the title compound, see: Esteves *et al.* (1999). For related structures, see: Zhang *et al.* (2006); Rao *et al.* (2009).



Experimental

Crystal data

 $\text{C}_{21}\text{H}_{29}\text{BrO}_2$ $M_r = 393.35$

Monoclinic, C_2
 $a = 13.888$ (3) Å
 $b = 6.1260$ (12) Å
 $c = 23.382$ (5) Å
 $\beta = 103.19$ (3)°
 $V = 1936.8$ (7) Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 2.13$ mm⁻¹
 $T = 293$ K
 $0.30 \times 0.20 \times 0.20$ mm

Data collection

Enraf–Nonius CAD-4
 diffractometer
 Absorption correction: ψ scan
 (North *et al.*, 1968)
 $T_{\min} = 0.567$, $T_{\max} = 0.675$
 3557 measured reflections

3505 independent reflections
 2754 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.035$
 3 standard reflections every 200
 reflections
 intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.128$
 $S = 1.01$
 3505 reflections
 218 parameters
 1 restraint

H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.64$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.40$ e Å⁻³
 Absolute structure: Flack (1983),
 1563 Friedel pairs
 Flack parameter: 0.010 (15)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C}21-\text{H}21A\cdots\text{O}2^i$	0.96	2.72	3.652 (10)	165

Symmetry code: (i) $x, y + 1, z$.

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

This project was supported by the Forestry Commonwealth Industry Special Foundation of China (No. 200704008).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZQ2035).

References

- Enraf–Nonius (1994). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
- Esteves, M. A., Narender, N., Gigante, B., Marcelo-Curto, M. J. & Alvarez, A. (1999). *Synth. Commun.* **29**, 275–280.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Fonseca, T., Gigante, B. & Gilchrist, T. L. (2001). *Tetrahedron*, **57**, 1793–1799.
- Halbrook, N. J. & Lawrence, R. V. (1966). *J. Org. Chem.* **31**, 4246–4247.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Pan, Y.-M., Zhang, Y., Wang, H.-S. & Li, F. (2006). *Chin. J. Org. Chem.* **26**, 314–317.
- Rao, X.-P., Song, Z.-Q. & Shang, S.-B. (2009). *Acta Cryst.* **E65**, o2402.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Zhang, Y., Pan, Y.-M., Wang, H.-S., Wu, Q. & Zhang, Y. (2006). *Acta Cryst.* **E62**, o5425–o5426.

supplementary materials

Acta Cryst. (2010). E66, o1207 [doi:10.1107/S1600536810012201]

Methyl 12-bromodehydroabietate

H. Gao, Z.-Q. Song and S.-B. Shang

Comment

Dehydroabietic acid is a readily obtainable compound, which is isolated from disproportionated rosin by methods of amination (Halbrook & Lawrence, 1966). A number of new derivatives of dehydroabietic acid have been prepared (Fonseca *et al.*, 2001). The title compound is one of modified products of dehydroabietic acid, which could be used in synthesis of many fluorescence derivatization reagents (Pan *et al.*, 2006). Although, it has been first prepared by Esteves *et al.* (1999), the crystal structure of the title compound methyl 12-bromo-dehydroabietate was not yet reported. In this work, we present its crystal structure, the molecular structure is shown in Fig. 1. There are three six-membered rings, which adopt planar, half-chair and chair conformations. The two cyclohexane rings form a *trans* ring junction with the two methyl groups in axial positions. The crystal structure is stabilized by weak intermolecular O—H...O contacts and π - π stacking interactions (centroid-centroid distance = 6.126 Å) along the *b* axis (Fig. 2).

Experimental

p-Toluene sulfochloride (4.6 g), potassium carbonate (3.4 g) and methanol anhydrous (20 ml) were mixed and stirred at room temperature for 2 h, then dehydroabietic acid (6.0 g) was added into the mixture to react for 3 h. The mixture was filtered and extracted by ethyl ether and recrystallized by methanol, methyl dehydroabietate (5.3 g) was prepared. Subsequently, methyl dehydroabietate (1.0 g), *N*-bromosuccinimide (1.0 g) and acetonitrile (100 ml) were mixed and stirred for 24 h at room temperature, the precipitate was filtered and recrystallized from methanol. Suitable crystals of the title compound for X-ray diffraction were obtained by slow evaporation of a methanol solution.

Refinement

All H atoms bonded to the C atoms were placed geometrically at bond distances of 0.93–0.98 Å and included in the refinement in a riding motion approximation with $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5U_{\text{eq}}$ of the carrier atom.

Figures



Fig. 1. A view of the molecular structure of the title compound, showing displacement ellipsoids at the 30% probability level.




Fig. 2. A view of the packing of the title compound along (010).

(1R)-methyl 6-bromo-7-isopropyl-1,4a-dimethyl-1,2,3,4,4a,9,10,10a-octahydrophenanthrene-1-carboxylate

Crystal data

$C_{21}H_{29}BrO_2$

$M_r = 393.35$

Monoclinic, $C2$

Hall symbol: $C\ 2y$

$a = 13.888\ (3)\ \text{\AA}$

$b = 6.1260\ (12)\ \text{\AA}$

$c = 23.382\ (5)\ \text{\AA}$

$\beta = 103.19\ (3)^\circ$

$V = 1936.8\ (7)\ \text{\AA}^3$

$Z = 4$

$F(000) = 824$

$D_x = 1.349\ \text{Mg m}^{-3}$

Melting point: 416 K

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 25 reflections

$\theta = 10\text{--}13^\circ$

$\mu = 2.13\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Rod, colorless

$0.30 \times 0.20 \times 0.20\ \text{mm}$

Data collection

Enraf-Nonius CAD-4
diffractometer

Radiation source: fine-focus sealed tube
graphite

$\omega/2\theta$ scans

Absorption correction: ψ scan
(North *et al.*, 1968)

$T_{\min} = 0.567$, $T_{\max} = 0.675$

3657 measured reflections

3505 independent reflections

2754 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.035$

$\theta_{\max} = 25.3^\circ$, $\theta_{\min} = 1.8^\circ$

$h = 0 \rightarrow 16$

$k = -7 \rightarrow 7$

$l = -28 \rightarrow 27$

3 standard reflections every 200 reflections

intensity decay: 1%

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.047$

$wR(F^2) = 0.128$

$S = 1.01$

3505 reflections

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.083P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.64\ \text{e \AA}^{-3}$

$\Delta\rho_{\min} = -0.40\ \text{e \AA}^{-3}$

218 parameters
 1 restraint
 Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map

Extinction correction: SHELXL97 (Sheldrick, 2008),
 $F_c^* = kF_c [1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.0175 (13)
 Absolute structure: Flack (1983), 1563 Friedel pairs
 Flack parameter: 0.010 (15)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br	0.52734 (4)	0.20204 (10)	0.41747 (2)	0.0569 (3)
C1	0.4547 (4)	-0.1955 (9)	0.2063 (2)	0.0396 (12)
H1A	0.4131	-0.2111	0.2342	0.048*
H1B	0.4601	-0.0409	0.1985	0.048*
O1	0.6426 (4)	-0.1241 (7)	0.0769 (2)	0.0621 (13)
O2	0.6696 (4)	-0.4626 (8)	0.0554 (2)	0.0841 (17)
C2	0.4054 (3)	-0.3087 (14)	0.1497 (2)	0.0478 (12)
H2A	0.3425	-0.2385	0.1334	0.057*
H2B	0.3922	-0.4594	0.1581	0.057*
C3	0.4686 (3)	-0.3032 (13)	0.10451 (19)	0.0448 (11)
H3A	0.4744	-0.1532	0.0924	0.054*
H3B	0.4354	-0.3851	0.0701	0.054*
C4	0.5724 (4)	-0.3975 (9)	0.1272 (2)	0.0387 (12)
C5	0.6204 (3)	-0.2861 (11)	0.18667 (18)	0.0324 (9)
H5A	0.6258	-0.1318	0.1768	0.039*
C6	0.7266 (4)	-0.3588 (9)	0.2118 (2)	0.0423 (14)
H6A	0.7269	-0.4996	0.2309	0.051*
H6B	0.7614	-0.3732	0.1805	0.051*
C7	0.7773 (4)	-0.1913 (11)	0.2557 (3)	0.0502 (15)
H7A	0.7984	-0.0709	0.2346	0.060*
H7B	0.8361	-0.2568	0.2801	0.060*
C8	0.7139 (3)	-0.1013 (9)	0.2953 (2)	0.0361 (11)
C9	0.6128 (3)	-0.1423 (8)	0.2847 (2)	0.0320 (11)
C10	0.5593 (3)	-0.2863 (11)	0.23428 (18)	0.0319 (9)
C11	0.5608 (4)	-0.0505 (9)	0.3232 (2)	0.0357 (11)
H11A	0.4930	-0.0738	0.3170	0.043*
C12	0.6087 (4)	0.0746 (8)	0.3703 (2)	0.0339 (11)

supplementary materials

C13	0.7099 (3)	0.1180 (8)	0.3821 (2)	0.0347 (11)
C14	0.7593 (4)	0.0265 (9)	0.3430 (2)	0.0406 (12)
H14A	0.8269	0.0517	0.3490	0.049*
C15	0.7620 (4)	0.2626 (9)	0.4324 (2)	0.0419 (14)
H15A	0.7239	0.2576	0.4629	0.050*
C16	0.8676 (4)	0.1866 (15)	0.4603 (3)	0.0641 (15)
H16A	0.8662	0.0379	0.4730	0.096*
H16B	0.8952	0.2774	0.4934	0.096*
H16C	0.9076	0.1969	0.4318	0.096*
C17	0.7635 (5)	0.5004 (10)	0.4120 (3)	0.0601 (17)
H17A	0.6973	0.5469	0.3946	0.090*
H17B	0.8033	0.5113	0.3834	0.090*
H17C	0.7909	0.5918	0.4450	0.090*
C18	0.6337 (4)	-0.3401 (9)	0.0833 (2)	0.0398 (14)
C19	0.5695 (5)	-0.6506 (9)	0.1291 (3)	0.0527 (16)
H19A	0.5388	-0.7056	0.0908	0.079*
H19B	0.5321	-0.6966	0.1568	0.079*
H19C	0.6357	-0.7063	0.1408	0.079*
C20	0.5494 (4)	-0.5132 (8)	0.2616 (2)	0.0421 (13)
H20A	0.6140	-0.5693	0.2789	0.063*
H20B	0.5160	-0.6113	0.2316	0.063*
H20C	0.5121	-0.4995	0.2913	0.063*
C21	0.6987 (5)	-0.0491 (12)	0.0371 (3)	0.0620 (18)
H21A	0.6992	0.1076	0.0368	0.093*
H21B	0.6697	-0.1020	-0.0016	0.093*
H21C	0.7653	-0.1021	0.0492	0.093*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br	0.0488 (3)	0.0726 (4)	0.0544 (3)	-0.0030 (4)	0.0226 (2)	-0.0201 (4)
C1	0.033 (3)	0.045 (3)	0.040 (3)	0.000 (2)	0.008 (2)	-0.005 (2)
O1	0.100 (4)	0.038 (2)	0.066 (3)	-0.015 (2)	0.057 (3)	-0.002 (2)
O2	0.135 (5)	0.052 (3)	0.093 (4)	0.006 (3)	0.082 (4)	-0.007 (3)
C2	0.042 (3)	0.052 (3)	0.048 (3)	-0.001 (4)	0.006 (2)	-0.007 (4)
C3	0.050 (3)	0.048 (3)	0.035 (2)	-0.004 (4)	0.006 (2)	-0.002 (4)
C4	0.046 (3)	0.028 (2)	0.044 (3)	-0.005 (2)	0.015 (2)	-0.005 (2)
C5	0.038 (2)	0.023 (2)	0.038 (2)	-0.003 (3)	0.0137 (18)	0.000 (3)
C6	0.038 (3)	0.044 (4)	0.049 (3)	0.006 (2)	0.018 (2)	-0.001 (2)
C7	0.029 (3)	0.075 (4)	0.052 (3)	0.003 (3)	0.018 (3)	-0.005 (3)
C8	0.033 (3)	0.041 (3)	0.034 (3)	0.000 (2)	0.008 (2)	0.003 (2)
C9	0.028 (3)	0.029 (3)	0.038 (3)	0.004 (2)	0.005 (2)	0.006 (2)
C10	0.030 (2)	0.026 (2)	0.041 (2)	-0.002 (3)	0.0099 (17)	-0.002 (3)
C11	0.030 (3)	0.039 (3)	0.040 (3)	-0.003 (2)	0.013 (2)	0.000 (2)
C12	0.039 (3)	0.031 (3)	0.034 (3)	0.005 (2)	0.013 (2)	0.004 (2)
C13	0.036 (3)	0.034 (3)	0.031 (3)	-0.004 (2)	0.001 (2)	0.006 (2)
C14	0.027 (2)	0.049 (3)	0.045 (3)	-0.006 (2)	0.008 (2)	0.001 (3)
C15	0.043 (3)	0.046 (4)	0.035 (3)	-0.004 (2)	0.005 (2)	-0.001 (2)

C16	0.058 (3)	0.060 (4)	0.063 (3)	-0.006 (4)	-0.008 (3)	-0.006 (4)
C17	0.067 (4)	0.044 (4)	0.062 (4)	-0.008 (3)	-0.002 (3)	-0.003 (3)
C18	0.051 (3)	0.034 (4)	0.039 (3)	-0.004 (3)	0.019 (2)	-0.008 (3)
C19	0.075 (4)	0.028 (3)	0.061 (4)	-0.011 (3)	0.029 (3)	-0.010 (3)
C20	0.053 (3)	0.031 (3)	0.043 (3)	-0.005 (2)	0.013 (3)	0.004 (2)
C21	0.090 (5)	0.052 (4)	0.052 (4)	-0.026 (4)	0.034 (4)	-0.010 (3)

Geometric parameters (Å, °)

Br—C12	1.916 (5)	C9—C11	1.393 (7)
C1—C2	1.512 (8)	C9—C10	1.524 (7)
C1—C10	1.553 (7)	C10—C20	1.549 (8)
C1—H1A	0.9700	C11—C12	1.382 (7)
C1—H1B	0.9700	C11—H11A	0.9300
O1—C18	1.340 (8)	C12—C13	1.396 (7)
O1—C21	1.420 (7)	C13—C14	1.381 (7)
O2—C18	1.178 (7)	C13—C15	1.517 (7)
C2—C3	1.520 (6)	C14—H14A	0.9300
C2—H2A	0.9700	C15—C17	1.535 (8)
C2—H2B	0.9700	C15—C16	1.535 (8)
C3—C4	1.532 (7)	C15—H15A	0.9800
C3—H3A	0.9700	C16—H16A	0.9600
C3—H3B	0.9700	C16—H16B	0.9600
C4—C18	1.517 (7)	C16—H16C	0.9600
C4—C19	1.552 (8)	C17—H17A	0.9600
C4—C5	1.555 (7)	C17—H17B	0.9600
C5—C6	1.524 (6)	C17—H17C	0.9600
C5—C10	1.547 (5)	C19—H19A	0.9600
C5—H5A	0.9800	C19—H19B	0.9600
C6—C7	1.507 (8)	C19—H19C	0.9600
C6—H6A	0.9700	C20—H20A	0.9600
C6—H6B	0.9700	C20—H20B	0.9600
C7—C8	1.519 (7)	C20—H20C	0.9600
C7—H7A	0.9700	C21—H21A	0.9600
C7—H7B	0.9700	C21—H21B	0.9600
C8—C14	1.391 (7)	C21—H21C	0.9600
C8—C9	1.391 (7)		
C2—C1—C10	113.4 (4)	C20—C10—C1	109.5 (4)
C2—C1—H1A	108.9	C5—C10—C1	108.1 (4)
C10—C1—H1A	108.9	C12—C11—C9	120.8 (4)
C2—C1—H1B	108.9	C12—C11—H11A	119.6
C10—C1—H1B	108.9	C9—C11—H11A	119.6
H1A—C1—H1B	107.7	C11—C12—C13	122.9 (4)
C18—O1—C21	118.1 (5)	C11—C12—Br	116.5 (4)
C1—C2—C3	112.3 (5)	C13—C12—Br	120.5 (4)
C1—C2—H2A	109.1	C14—C13—C12	114.9 (5)
C3—C2—H2A	109.1	C14—C13—C15	122.0 (4)
C1—C2—H2B	109.1	C12—C13—C15	123.1 (5)
C3—C2—H2B	109.1	C13—C14—C8	123.8 (4)

supplementary materials

H2A—C2—H2B	107.9	C13—C14—H14A	118.1
C2—C3—C4	113.4 (4)	C8—C14—H14A	118.1
C2—C3—H3A	108.9	C13—C15—C17	110.5 (4)
C4—C3—H3A	108.9	C13—C15—C16	113.1 (5)
C2—C3—H3B	108.9	C17—C15—C16	109.9 (5)
C4—C3—H3B	108.9	C13—C15—H15A	107.7
H3A—C3—H3B	107.7	C17—C15—H15A	107.7
C18—C4—C3	107.9 (4)	C16—C15—H15A	107.7
C18—C4—C19	105.9 (4)	C15—C16—H16A	109.5
C3—C4—C19	111.0 (5)	C15—C16—H16B	109.5
C18—C4—C5	108.1 (4)	H16A—C16—H16B	109.5
C3—C4—C5	108.8 (4)	C15—C16—H16C	109.5
C19—C4—C5	115.0 (5)	H16A—C16—H16C	109.5
C6—C5—C10	111.3 (4)	H16B—C16—H16C	109.5
C6—C5—C4	113.4 (4)	C15—C17—H17A	109.5
C10—C5—C4	116.7 (4)	C15—C17—H17B	109.5
C6—C5—H5A	104.7	H17A—C17—H17B	109.5
C10—C5—H5A	104.7	C15—C17—H17C	109.5
C4—C5—H5A	104.7	H17A—C17—H17C	109.5
C7—C6—C5	109.0 (4)	H17B—C17—H17C	109.5
C7—C6—H6A	109.9	O2—C18—O1	120.4 (5)
C5—C6—H6A	109.9	O2—C18—C4	126.9 (5)
C7—C6—H6B	109.9	O1—C18—C4	112.6 (4)
C5—C6—H6B	109.9	C4—C19—H19A	109.5
H6A—C6—H6B	108.3	C4—C19—H19B	109.5
C6—C7—C8	114.6 (4)	H19A—C19—H19B	109.5
C6—C7—H7A	108.6	C4—C19—H19C	109.5
C8—C7—H7A	108.6	H19A—C19—H19C	109.5
C6—C7—H7B	108.6	H19B—C19—H19C	109.5
C8—C7—H7B	108.6	C10—C20—H20A	109.5
H7A—C7—H7B	107.6	C10—C20—H20B	109.5
C14—C8—C9	119.9 (5)	H20A—C20—H20B	109.5
C14—C8—C7	118.2 (4)	C10—C20—H20C	109.5
C9—C8—C7	121.8 (5)	H20A—C20—H20C	109.5
C8—C9—C11	117.6 (5)	H20B—C20—H20C	109.5
C8—C9—C10	122.4 (4)	O1—C21—H21A	109.5
C11—C9—C10	120.0 (4)	O1—C21—H21B	109.5
C9—C10—C20	105.9 (4)	H21A—C21—H21B	109.5
C9—C10—C5	107.8 (4)	O1—C21—H21C	109.5
C20—C10—C5	114.4 (5)	H21A—C21—H21C	109.5
C9—C10—C1	111.3 (5)	H21B—C21—H21C	109.5
C10—C1—C2—C3	-55.4 (8)	C6—C5—C10—C1	176.2 (5)
C1—C2—C3—C4	55.2 (9)	C4—C5—C10—C1	-51.6 (7)
C2—C3—C4—C18	-168.5 (6)	C2—C1—C10—C9	170.0 (5)
C2—C3—C4—C19	76.0 (7)	C2—C1—C10—C20	-73.3 (6)
C2—C3—C4—C5	-51.4 (7)	C2—C1—C10—C5	51.8 (7)
C18—C4—C5—C6	-60.0 (6)	C8—C9—C11—C12	-0.6 (7)
C3—C4—C5—C6	-176.8 (5)	C10—C9—C11—C12	177.9 (5)
C19—C4—C5—C6	58.0 (6)	C9—C11—C12—C13	0.4 (8)

C18—C4—C5—C10	168.7 (5)	C9—C11—C12—Br	177.2 (4)
C3—C4—C5—C10	51.8 (6)	C11—C12—C13—C14	0.1 (7)
C19—C4—C5—C10	-73.3 (6)	Br—C12—C13—C14	-176.5 (4)
C10—C5—C6—C7	-65.9 (6)	C11—C12—C13—C15	177.5 (5)
C4—C5—C6—C7	160.2 (5)	Br—C12—C13—C15	0.9 (7)
C5—C6—C7—C8	40.9 (7)	C12—C13—C14—C8	-0.5 (8)
C6—C7—C8—C14	169.8 (5)	C15—C13—C14—C8	-177.9 (5)
C6—C7—C8—C9	-10.7 (8)	C9—C8—C14—C13	0.3 (8)
C14—C8—C9—C11	0.2 (7)	C7—C8—C14—C13	179.8 (5)
C7—C8—C9—C11	-179.3 (5)	C14—C13—C15—C17	86.4 (6)
C14—C8—C9—C10	-178.2 (5)	C12—C13—C15—C17	-90.9 (6)
C7—C8—C9—C10	2.3 (8)	C14—C13—C15—C16	-37.3 (7)
C8—C9—C10—C20	98.6 (5)	C12—C13—C15—C16	145.4 (5)
C11—C9—C10—C20	-79.7 (6)	C21—O1—C18—O2	-1.8 (10)
C8—C9—C10—C5	-24.1 (7)	C21—O1—C18—C4	179.7 (5)
C11—C9—C10—C5	157.5 (5)	C3—C4—C18—O2	-117.3 (7)
C8—C9—C10—C1	-142.5 (5)	C19—C4—C18—O2	1.6 (9)
C11—C9—C10—C1	39.1 (6)	C5—C4—C18—O2	125.3 (7)
C6—C5—C10—C9	55.8 (6)	C3—C4—C18—O1	61.1 (6)
C4—C5—C10—C9	-172.0 (5)	C19—C4—C18—O1	179.9 (5)
C6—C5—C10—C20	-61.7 (6)	C5—C4—C18—O1	-56.4 (6)
C4—C5—C10—C20	70.6 (6)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C21—H21A \cdots O2 ⁱ	0.96	2.72	3.652 (10)	165

Symmetry codes: (i) $x, y+1, z$.

Fig. 1




Fig. 2

