

Triaqua(1,4,7-triazacyclononane- κ^3N^1,N^4,N^7)nickel(II) bromide nitrate

Changchun Wen, Jianqi Lu and Zhong Zhang*

College of Chemistry and Chemical Engineering, Guangxi Normal University, Yucui Road 15, Guilin 541004, People's Republic of China

Correspondence e-mail: zhangzhong@mailbox.gxnu.edu.cn


Received 8 April 2010; accepted 3 May 2010

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(C-C) = 0.006$ Å; R factor = 0.035; wR factor = 0.072; data-to-parameter ratio = 142.9.

In the title half-sandwich compound, $[Ni(C_6H_{15}N_3)(H_2O)_3]Br(NO_3)$, the central Ni^{II} ion, lying on a threefold rotation axis, is six-coordinated by three amine N atoms from the face-capping triaza macrocycle and three water O atoms in a slightly distorted octahedral geometry. In the crystal, $O-H \cdots O$ hydrogen bonding and weak $O-H \cdots Br$ interactions associate the Ni^{II} cations and the counter-ions into a three-dimensional supramolecular network.

Related literature

For the preparation of 1,4,7-triazacyclononane trihydrobromide, see: Koyama & Yoshino (1972). For the applications of metal complexes containing 1,4,7-triazacyclononane as small-molecule models of metalloenzymes and metalloproteins and as molecule-based magnets, see: Berseth *et al.* (2000); Chaudhury *et al.* (1985); Cheng *et al.* (2004); Deal *et al.* (1996); Hegg & Burstyn (1995); Hegg *et al.* (1997); Lin *et al.* (2001); Poganiuch *et al.* (1991); Williams *et al.* (1999). For related Ni^{II} complexes with 1,4,7-triazacyclononane, see: Bencini *et al.* (1990); Stranger *et al.* (1992); Wang *et al.* (2003, 2005); Zompa & Margulis (1978).



Experimental

Crystal data

$[Ni(C_6H_{15}N_3)(H_2O)_3]Br(NO_3)$	$Z = 4$
$M_r = 383.89$	Mo $K\alpha$ radiation
Cubic, $P2_13$	$\mu = 4.14$ mm ⁻¹
$a = 11.300$ (1) Å	$T = 298$ K
$V = 1442.9$ (3) Å ³	$0.29 \times 0.27 \times 0.18$ mm

Data collection

Bruker APEXII CCD area-detector diffractometer	15223 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 1998)	1110 independent reflections
$T_{min} = 0.320$, $T_{max} = 0.480$	985 reflections with $I > 2\sigma(I)$
	$R_{int} = 0.080$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$	$\Delta\rho_{max} = 0.36$ e Å ⁻³
$wR(F^2) = 0.072$	$\Delta\rho_{min} = -0.47$ e Å ⁻³
$S = 1.03$	Absolute structure: Flack (1983),
1110 reflections	475 Friedel pairs
61 parameters	Flack parameter: 0.01 (3)
H atoms treated by a mixture of independent and constrained refinement	

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$O1-H4A \cdots O2^i$	0.84 (4)	1.95 (5)	2.776 (5)	162 (4)
$O1-H4B \cdots Br1^{ii}$	0.85 (5)	2.48 (5)	3.312 (3)	167 (4)

 Symmetry codes: (i) $-x + 2, y - \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: APEX2 (Bruker, 2002); cell refinement: SAINT (Bruker, 2002); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The authors are grateful for financial support from the Guangxi Science Foundation (grant No. 0832023) and the Scientific Research Foundation of Guangxi Normal University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PB2027).

References

- Bencini, A., Bianchi, A., Paoli, P., Garcia-Espana, E., Julve, M. & Marcelino, V. (1990). *J. Chem. Soc. Dalton Trans.* pp. 2213–2217.
- Berseth, P. A., Sokol, J. J., Shores, M. P., Heinrich, J. L. & Long, J. R. (2000). *J. Am. Chem. Soc.* **122**, 9655–9662.
- Bruker (1998). SADABS. Bruker AXS inc., Madison, Wisconsin, USA.
- Bruker (2002). APEX2 and SAINT. Bruker AXS inc., Madison, Wisconsin, USA.
- Chaudhury, P., Wieghardt, K., Nuber, B. & Weiss, J. (1985). *Angew. Chem. Int. Ed. Engl.* **24**, 778–779.
- Cheng, P., Yan, S. P., Xie, C. Z., Zhao, B., Chen, X. Y., Liu, X. W., Li, C. H., Liao, D. Z., Jiang, Z. H. & Wang, G. L. (2004). *Eur. J. Inorg. Chem.* pp. 2369–2378.
- Deal, K. A., Hengge, A. C. & Burstyn, J. N. (1996). *J. Am. Chem. Soc.* **118**, 1713–1718.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.

- Hegg, E. L. & Burstyn, J. N. (1995). *J. Am. Chem. Soc.* **117**, 7015–7016.
- Hegg, E. L., Deal, K. A., Kiessling, L. L. & Burstyn, J. N. (1997). *Inorg. Chem.* **36**, 1715–1718.
- Koyama, H. & Yoshino, T. (1972). *Bull. Chem. Soc. Jpn.*, **45**, 481–484.
- Lin, G., Reid, G. & Bugg, T. D. H. (2001). *J. Am. Chem. Soc.* **123**, 5030–5039.
- Poganiuch, P., Liu, S. C., Papaefthymiou, G. C. & Lippard, S. J. (1991). *J. Am. Chem. Soc.* **113**, 4645–4651.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Stranger, R., Wallis, S. C., Gahan, L. R., Kennard, C. H. L. & Byriel, K. A. (1992). *J. Chem. Soc. Dalton Trans.* pp. 2971–2976.
- Wang, Z. D., Gao, F. X., Han, W., Xie, C. Z. & Yan, S. P. (2005). *Ultramicroscopy*, **38**, 80–82.
- Wang, Q. L., Yu, L. H., Liao, D. Z., Yan, S. P., Jiang, Z. H. & Cheng, P. (2003). *Helv. Chim. Acta*, **86**, 2441–2451.
- Williams, N. H., Lebus, A. M. & Chin, J. (1999). *J. Am. Chem. Soc.* **121**, 3341–3348.
- Zompa, L. J. & Margulis, T. N. (1978). *Inorg. Chim. Acta*, **28**, L157–L159.

supplementary materials

Acta Cryst. (2010). E66, m624-m625 [doi:10.1107/S160053681001620X]

Triaqua(1,4,7-triazacyclononane- κ^3N^1,N^4,N^7)nickel(II) bromide nitrate

C. Wen, J. Lu and Z. Zhang

Comment

The coordination chemistry of 1,4,7-triazacyclononane (TACN) has been extensively studied for its applications in the simulation of metalloenzymes and metalloproteins (Chaudhury *et al.*, 1985; Deal *et al.*, 1996; Hegg & Burstyn, 1995; Hegg *et al.*, 1997; Lin *et al.*, 2001; Williams *et al.*, 1999) as well as in constructing molecule-based magnetic materials (Berseth *et al.*, 2000; Cheng *et al.*, 2004; Poganiuch *et al.*, 1991). In general, TACN ligand can form stable sandwich complexes with many transition metals (Stranger *et al.*, 1992; Zompa & Margulis, 1978) or functions as a terminal chelator for the assembly of binuclear/polynuclear species and coordination polymers supported by bridging ligands (Bencini *et al.*, 1990; Wang *et al.*, 2005; Wang *et al.*, 2003). In this paper, a half-sandwich type Ni^{II} complex with TACN has been synthesized and characterized.

In the selected crystal, the title compound (I) crystallizes in a chiral space group $P2_13$ and Flack parameter of 0.01 (3) indicates that a spontaneous resolution has been achieved during crystallization. As depicted in Fig. 1, the Ni^{II} center in the complex cation lies on a three-fold rotation axis and three amine N atoms from facially coordinated TACN and three water molecules complete the slightly distorted octahedral arrangement. Upon coordination, three five-membered Ni—N—C—C—N chelating rings subtended at metal center adopt ($\lambda\lambda\lambda$) conformation, which is the source of the chirality of the crystal. Ni—N [2.091 (3) Å] and Ni—O [2.089 (3) Å] bond lengths are both in the normal ranges, meanwhile N—Ni—N bond angle is smaller than that of O—Ni—O due to the small size of TACN ring. Counter-ions NO₃⁻ and Br⁻ interconnect neighbouring cations by O—H \cdots O hydrogen bond and O—H \cdots Br⁻ weak interaction (Table 1) into three-dimensional supra-molecular network (Fig. 2).

Experimental

1,4,7-Triazacyclononane trihydrobromide (TACN·3HBr) was prepared by following a modified published method (Koyama & Yoshino, 1972).

To a solution of 0.074 g (0.02 mmol) of TACN·3HBr in water (10 ml), 0.1 M NaOH was added to adjust the pH to 6. Then aqueous solution (5 ml) of 0.058 g (0.02 mmol) of Ni(NO₃)₂·6H₂O was added and the resulting mixture was stirred under reflux for 6 h. After cooling, the mixture was filtered, and the filtrate was allowed to standing at ambient temperature. Plate-like green single crystals suitable for X-ray crystallographic analysis were collected by slow evaporation of the filtrate within two months.

Refinement

All methylene H atoms were placed at calculated positions and refined as riding on their parent atoms [C—H = 0.97 Å and $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$]. The H atoms of amine groups and water molecules were located in a difference Fourier map as riding atoms, with $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{N})$ and $1.5 U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. An ORTEP plot for the title compound (I) with the atom labelling scheme and 30% displacement ellipsoids. Symmetry codes: (i) $y+1/2, -z+3/2, -x+2$; (ii) $-z+2, x-1/2, -y+3/2$.




Fig. 2. A view of the packing diagram of the title compound (I), showing the hydrogen-bonding supramolecular network. Hydrogen bonds are drawn in dashed lines. H atoms not involved in hydrogen bonds are omitted for clarity.

Triaqua(1,4,7-triazacyclononane- κ^3N^1, N^4, N^7)nickel(II) bromide nitrate

Crystal data

$[\text{Ni}(\text{C}_6\text{H}_{15}\text{N}_3)(\text{H}_2\text{O})_3]\text{Br}(\text{NO}_3)$

$M_r = 383.89$

Cubic, $P2_13$

Hall symbol: P 2ac 2ab 3

$a = 11.300 (1) \text{ \AA}$

$V = 1442.9 (3) \text{ \AA}^3$

$Z = 4$

$F(000) = 784$

$D_x = 1.767 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 13409 reflections

$\theta = 3.1\text{--}27.4^\circ$

$\mu = 4.14 \text{ mm}^{-1}$

$T = 298 \text{ K}$

Plate, green

$0.29 \times 0.27 \times 0.18 \text{ mm}$

Data collection

Bruker APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

φ and ω scans

Absorption correction: multi-scan

1110 independent reflections

985 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.080$

$\theta_{\text{max}} = 27.4^\circ$, $\theta_{\text{min}} = 3.1^\circ$

$h = -14 \rightarrow 14$

(SADABS; Bruker, 1998)

$T_{\min} = 0.320$, $T_{\max} = 0.480$

15223 measured reflections

$k = -14 \rightarrow 14$

$l = -14 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.035$

$wR(F^2) = 0.072$

$S = 1.03$

8717 reflections

61 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0232P)^2 + 1.6516P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.36 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.46 \text{ e } \text{\AA}^{-3}$

Absolute structure: Flack (1983), 475 Friedel pairs

Flack parameter: 0.01 (3)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	1.06169 (4)	0.56169 (4)	0.93831 (4)	0.02729 (19)
Br1	0.25347 (4)	0.24653 (4)	0.75347 (4)	0.0437 (2)
C1	0.8566 (4)	0.4233 (4)	0.8872 (4)	0.0437 (11)
H1A	0.8123	0.3498	0.8858	0.052*
H1B	0.8438	0.4636	0.8125	0.052*
C2	1.0118 (4)	0.3128 (4)	0.9995 (4)	0.0449 (11)
H2A	1.0326	0.2365	0.9660	0.054*
H2B	0.9417	0.3022	1.0478	0.054*
N1	0.9850 (3)	0.3973 (3)	0.9019 (3)	0.0353 (8)
H3	1.0226	0.3631	0.8407	0.053*
N2	0.9466 (3)	0.9466 (3)	0.9466 (3)	0.0316 (11)
O1	1.0196 (3)	0.6447 (3)	0.7786 (3)	0.0391 (8)
O2	0.8854 (3)	1.0345 (3)	0.9196 (3)	0.0577 (9)

supplementary materials

H4B	0.947 (5)	0.659 (4)	0.765 (4)	0.050 (14)*
H4A	1.040 (4)	0.598 (4)	0.723 (4)	0.050 (14)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.02729 (19)	0.02729 (19)	0.02729 (19)	-0.0011 (2)	0.0011 (2)	0.0011 (2)
Br1	0.0437 (2)	0.0437 (2)	0.0437 (2)	-0.0012 (2)	0.0012 (2)	-0.0012 (2)
C1	0.041 (2)	0.042 (3)	0.049 (3)	-0.016 (2)	-0.009 (2)	0.005 (2)
C2	0.054 (3)	0.027 (2)	0.054 (3)	-0.0022 (19)	0.010 (2)	0.0063 (19)
N1	0.0365 (19)	0.0349 (19)	0.0346 (19)	-0.0022 (14)	0.0050 (14)	-0.0021 (14)
N2	0.0316 (11)	0.0316 (11)	0.0316 (11)	0.0002 (15)	0.0002 (15)	0.0002 (15)
O1	0.0424 (18)	0.0419 (18)	0.0330 (18)	0.0052 (14)	-0.0007 (13)	0.0024 (12)
O2	0.055 (2)	0.054 (2)	0.065 (2)	0.0156 (16)	0.0149 (17)	0.0133 (18)

Geometric parameters (\AA , $^\circ$)

Ni1—O1 ⁱ	2.089 (3)	C2—N1	1.490 (5)
Ni1—O1	2.089 (3)	C2—C1 ⁱⁱ	1.520 (6)
Ni1—O1 ⁱⁱ	2.089 (3)	C2—H2A	0.9700
Ni1—N1	2.091 (3)	C2—H2B	0.9700
Ni1—N1 ⁱⁱ	2.091 (3)	N1—H3	0.8987
Ni1—N1 ⁱ	2.091 (3)	N2—O2 ⁱⁱⁱ	1.248 (3)
C1—N1	1.490 (6)	N2—O2 ^{iv}	1.248 (3)
C1—C2 ⁱ	1.520 (6)	N2—O2	1.248 (3)
C1—H1A	0.9700	O1—H4B	0.85 (5)
C1—H1B	0.9700	O1—H4A	0.84 (4)
O1 ⁱ —Ni1—O1	84.90 (14)	H1A—C1—H1B	108.1
O1 ⁱ —Ni1—O1 ⁱⁱ	84.90 (14)	N1—C2—C1 ⁱⁱ	111.7 (3)
O1—Ni1—O1 ⁱⁱ	84.90 (14)	N1—C2—H2A	109.3
O1 ⁱ —Ni1—N1	177.00 (13)	C1 ⁱⁱ —C2—H2A	109.3
O1—Ni1—N1	97.72 (13)	N1—C2—H2B	109.3
O1 ⁱⁱ —Ni1—N1	93.87 (12)	C1 ⁱⁱ —C2—H2B	109.3
O1 ⁱ —Ni1—N1 ⁱⁱ	93.87 (12)	H2A—C2—H2B	108.0
O1—Ni1—N1 ⁱⁱ	177.00 (13)	C1—N1—C2	114.0 (3)
O1 ⁱⁱ —Ni1—N1 ⁱⁱ	97.72 (12)	C1—N1—Ni1	104.5 (3)
N1—Ni1—N1 ⁱⁱ	83.58 (14)	C2—N1—Ni1	109.8 (3)
O1 ⁱ —Ni1—N1 ⁱ	97.72 (12)	C1—N1—H3	117.3
O1—Ni1—N1 ⁱ	93.87 (12)	C2—N1—H3	101.4
O1 ⁱⁱ —Ni1—N1 ⁱ	177.00 (13)	Ni1—N1—H3	109.7
N1—Ni1—N1 ⁱ	83.58 (14)	O2 ⁱⁱⁱ —N2—O2 ^{iv}	119.999 (2)
N1 ⁱⁱ —Ni1—N1 ⁱ	83.58 (14)	O2 ⁱⁱⁱ —N2—O2	120.000 (3)
N1—C1—C2 ⁱ	110.3 (4)	O2 ^{iv} —N2—O2	120.000 (2)
N1—C1—H1A	109.6	Ni1—O1—H4B	117 (4)

C2 ⁱ —C1—H1A	109.6	Ni1—O1—H4A	107 (4)
N1—C1—H1B	109.6	H4B—O1—H4A	104 (5)
C2 ⁱ —C1—H1B	109.6		
C2 ⁱ —C1—N1—C2	72.1 (5)	N1 ⁱⁱ —Ni1—N1—C1	114.6 (2)
C2 ⁱ —C1—N1—Ni1	-47.8 (4)	N1 ⁱ —Ni1—N1—C1	30.4 (3)
C1 ⁱⁱ —C2—N1—C1	-133.2 (4)	O1—Ni1—N1—C2	174.6 (3)
C1 ⁱⁱ —C2—N1—Ni1	-16.3 (4)	O1 ⁱⁱ —Ni1—N1—C2	89.3 (3)
O1—Ni1—N1—C1	-62.7 (3)	N1 ⁱⁱ —Ni1—N1—C2	-8.1 (3)
O1 ⁱⁱ —Ni1—N1—C1	-148.0 (3)	N1 ⁱ —Ni1—N1—C2	-92.3 (2)

Symmetry codes: (i) $y+1/2, -z+3/2, -x+2$; (ii) $-z+2, x-1/2, -y+3/2$; (iii) z, x, y ; (iv) y, z, x .

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H4A \cdots O2 ^v	0.84 (4)	1.95 (5)	2.776 (5)	162 (4)
O1—H4B \cdots Br1 ^{vi}	0.85 (5)	2.48 (5)	3.312 (3)	167 (4)

Symmetry codes: (v) $-x+2, y-1/2, -z+3/2$; (vi) $-x+1, y+1/2, -z+3/2$.

Fig. 1




Fig. 2

