

# Poly[diethylenetriammonium [aquadi- $\mu_2$ -sulfato-sulfatocerium(III)]]

Xiu-Mei Zhang<sup>a</sup> and Ya-Feng Li<sup>b\*</sup>

<sup>a</sup>Department of X-ray, First Hospital, Jilin University, Changchun 130021, People's Republic of China, and <sup>b</sup>School of Chemical Engineering, Changchun University of Technology, Changchun 130012, People's Republic of China

Correspondence e-mail: fly012345@sohu.com


Received 30 April 2010; accepted 6 May 2010

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.015;  $wR$  factor = 0.041; data-to-parameter ratio = 15.5.

A new organically templated open-framework cerium sulfate,  $\{(\text{C}_4\text{H}_{16}\text{N}_3)[\text{Ce}(\text{SO}_4)_3(\text{H}_2\text{O})]\}_n$ , was hydrothermally synthesized. The  $\text{Ce}^{\text{III}}$  cation is nine-coordinated by nine O atoms, including one water molecule. Two of the  $\text{SO}_4$  groups afford one monodentate and bidentate linkages as the bridge to connect adjacent  $\text{Ce}^{\text{III}}$  cations, while the third  $\text{SO}_4$  group attaches the  $\text{Ce}^{\text{III}}$  cation in a bidentate mode. The crystal structure consists of layers composed of eight-membered-ring networks formed by four  $\text{CeO}_9$  polyhedra and four  $\text{SO}_4$  tetrahedra. The triply protonated diethylenetriamine cations are located between adjacent layers and connect the layers *via* hydrogen bonds.

## Related literature

For related literature, see: Choudhury *et al.* (2001); Fu *et al.* (2006); Paul *et al.* (2002); Rao *et al.* (2006); Wickleder (2002).



## Experimental

### Crystal data

$(\text{C}_4\text{H}_{16}\text{N}_3)[\text{Ce}(\text{SO}_4)_3(\text{H}_2\text{O})]$

$M_r = 552.51$

Monoclinic,  $P2_1$

$a = 6.6774$  (13) Å

$b = 10.397$  (2) Å

$c = 11.093$  (2) Å

$\beta = 93.77$  (3)°

$V = 768.5$  (3) Å<sup>3</sup>

$Z = 2$

Mo  $K\alpha$  radiation

$\mu = 3.44$  mm<sup>-1</sup>

$T = 293$  K

$0.25 \times 0.22 \times 0.19$  mm

### Data collection

Rigaku R-Axis RAPID

diffractometer

Absorption correction: empirical

(using intensity measurements)

(*ABSCOR*; Higashi, 1995)

$T_{\text{min}} = 0.480$ ,  $T_{\text{max}} = 0.561$

7575 measured reflections

3485 independent reflections

3443 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.017$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.015$

$wR(F^2) = 0.041$

$S = 1.15$

3485 reflections

225 parameters

4 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 0.56$  e Å<sup>-3</sup>

$\Delta\rho_{\text{min}} = -0.71$  e Å<sup>-3</sup>

Absolute structure: Flack (1983)

Flack parameter:  $-0.009$  (8)

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1W}-\text{H1F}\cdots\text{O4}$	0.83 (2)	1.98 (2)	2.766 (3)	159 (4)
$\text{O1W}-\text{H1G}\cdots\text{O11}^i$	0.81 (2)	2.06 (2)	2.850 (3)	164 (4)
$\text{N1}-\text{H1A}\cdots\text{O8}^{ii}$	0.89	2.02	2.769 (3)	141
$\text{N1}-\text{H1C}\cdots\text{O9}^{ii}$	0.89	2.02	2.883 (3)	162
$\text{N1}-\text{H1B}\cdots\text{O6}^{iii}$	0.89	2.05	2.852 (3)	150
$\text{N2}-\text{H2B}\cdots\text{O11}$	0.90	1.92	2.764 (4)	156
$\text{N2}-\text{H2A}\cdots\text{O2}^{iv}$	0.90	2.16	2.993 (3)	154
$\text{N2}-\text{H2A}\cdots\text{O4}^{iv}$	0.90	2.30	2.997 (3)	134
$\text{N3}-\text{H3A}\cdots\text{O5}^v$	0.89	2.17	2.808 (3)	128
$\text{N3}-\text{H3A}\cdots\text{O3}^{vi}$	0.89	2.26	3.059 (4)	150
$\text{N3}-\text{H3C}\cdots\text{O12}^v$	0.89	1.91	2.799 (4)	173
$\text{N3}-\text{H3B}\cdots\text{O10}^{vii}$	0.89	2.04	2.763 (4)	137

Symmetry codes: (i)  $-x + 1, y + \frac{1}{2}, -z + 1$ ; (ii)  $-x + 1, y - \frac{1}{2}, -z + 2$ ; (iii)  $x - 1, y, z$ ; (iv)  $x, y - 1, z$ ; (v)  $-x + 1, y - \frac{1}{2}, -z + 1$ ; (vi)  $-x, y - \frac{1}{2}, -z + 1$ ; (vii)  $x - 1, y - 1, z$ .

Data collection: *PROCESS-AUTO* (Rigaku, 1998); cell refinement: *PROCESS-AUTO*; data reduction: *CrystalStructure* (Rigaku/MSK, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 2000); software used to prepare material for publication: *SHELXL97*.

This project was sponsored by the Scientific Research Foundation for Returned Overseas Chinese Scholars, Chinese Education Ministry (grant No. 20071108) and the Scientific Research Foundation for the Returned Overseas Team, Chinese Education Ministry.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5433).

## References

- Brandenburg, K. (2000). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Choudhury, A., Krishnamoorthy, J. & Rao, C. N. R. (2001). *Chem. Commun.* pp. 2610–2611.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Fu, Y., Xu, Z. & Ren, J. (2006). *J. Mol. Struct.* **788**, 190–193.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.

## metal-organic compounds

---

Paul, G., Choudhury, A. & Rao, C. N. R. (2002). *J. Chem. Soc. Dalton Trans.* pp. 3859–3867.

Rao, C. N. R., Behera, J. N. & Dan, M. (2006). *Chem. Soc. Rev.* **35**, 375–387.

Rigaku (1998). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.

Rigaku/MSC (2002). *CrystalStructure*. Rigaku/MSC, The Woodlands, Texas, USA.

Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

Wickleder, M. S. (2002). *Chem. Rev.* **102**, 2011–2087.

**supplementary materials**

*Acta Cryst.* (2010). E66, m649-m650 [ doi:10.1107/S1600536810016600 ]

**Poly[diethylenetriammonium [ $\mu_2$ -sulfato-sulfatocerium(III)]]**

**X.-M. Zhang and Y.-F. Li**

**Comment**

The hydrous and anhydrous lanthanide sulfates have been intensively studied due to use of the separation of rare earth elements (Wickleder, 2002). Since the pioneering works of Rao *et al.* (Choudhury, *et al.*, 2001; Paul, *et al.*, 2002; Rao, *et al.*, 2006) on the preparation of organically templated open-framework metal sulfates, a remarkable plenty of organically templated open-framework rare-earth sulfates have been described also. The example of organically templated cerium sulfate is few reported except for  $(C_4H_{12}N_2)_4[Ce_8(SO_4)_{16}(H_2O)_8]$  and  $(C_2H_{10}N_2)_2[Ce_2(SO_4)_5(H_2O)_2]$  (Fu, *et al.*, 2006). In this work, a new layer cerium sulfate,  $\{(C_4H_{16}N_3)[Ce(SO_4)_3(H_2O)]\}_n$ , is obtained.

The asymmetric unit of (I) comprises of one  $Ce^{III}$  cation, three  $SO_4$  groups, one coordination water and one protonated diethylene triamine cation, as shown in Fig.1. The  $Ce^{III}$  cation is 9-coordinated by nine oxygen including one water molecule with the bond distances from 2.468 (2) Å to 2.588 (27) Å and the angles of O—Ce—O between 54.18 (10)° and 149.13 (10)°. Three  $SO_4$  can be divided into two modes: S(1) and S(3) consist of three S—O—Ce linkages and links adjacent Ce atoms through one bidentate and one monodentate; S(2) makes two S—O—Ce linkages as a ligand of one Ce atom through bidentate. The bond angles of S—O—Ce of bidentate coordination range from 99.23 (10)° to 101.8 (1)°, and the S—O—Ce of monodentate coordination is at 141.81 (9)° and 144.17 (13)°.

As shown in Fig.2, the layer of (I) is accomplished by connect the Ce cations by  $\mu_2$ -S(1)O<sub>4</sub> and  $\mu_2$ -S(3)O<sub>4</sub> as the bridge along (100) and (010) direction, respectively. The S(2)O<sub>4</sub> do not take part in the formation of layer and coordinates to Ce cation by the bidentate mode. The protonated H<sub>3</sub>DETA interacts with the layer by the H-bond of N—H $\cdots$ O.

**Experimental**

(I) was synthesized under hydrothermal condition. In a typically route,  $Ce(NO_3)_3 \cdot 6H_2O$  (0.30 g, 0.7 mmol) was dissolved in 5 ml deionized water under stirring, and then  $H_2SO_4$  (95%, 0.25 ml, 4.55 mmol) and DETA (0.22 ml, 2.8 mmol) were dropwisely added to a clear solution with pH=3.0. After continuously stirred for 3 h, the solution with the molar ratio of  $Ce(NO_3)_3 \cdot 6H_2O : 6.5H_2SO_4 : 2.8DETA : 397H_2O$  was transferred into 23 ml autoclave and heated at 438 K for 5 days. After naturally cooling to room temperature, colorless block soluble product was collected by filtration as a single phase. The atomic ratio of Ce : S determined by EDX was 1 : 3, in consistence with the results of structural determination of (I).

**Refinement**

Water H atoms were located in a difference Fourier map and were refined with O—H = 0.82 (2) Å, H $\cdots$ H = 1.37 (2) Å and  $U_{iso}(H) = 1.2U_{eq}(O)$ . The remaining H-atoms were placed in calculated positions (C—H = 0.89 Å, N—H = 0.89-0.90 Å) and were included in the refinement in the riding-model approximation, with  $U(H) = 1.2U_{eq}(C, N)$ .

Figures




Fig. 1. The unit cell of (I), showing the atomic labelling scheme and displacement ellipsoids at the 50% probability level. [Symmetry codes: (i)  $1+x, y, z$ ; (ii)  $1-x, 0.5+y, -z$ .]




Fig. 2. The stick plot of (I), displaying the layer along (101) direction composed by linking the Ce cation with  $\mu_2$ -S(1)O<sub>4</sub> and  $\mu_2$ -S(3)O<sub>4</sub>. S(1) is shown in yellow, S(2) in green and S(3) in blue.

**Poly[diethylenetriammonium [aquadi- $\mu_2$ -sulfato-sulfatocerium(III)]]**

*Crystal data*

(C<sub>4</sub>H<sub>16</sub>N<sub>3</sub>)[Ce(SO<sub>4</sub>)<sub>3</sub>(H<sub>2</sub>O)]

$M_r = 552.51$

Monoclinic,  $P2_1$

Hall symbol: P 2yb

$a = 6.6774$  (13) Å

$b = 10.397$  (2) Å

$c = 11.093$  (2) Å

$\beta = 93.77$  (3)°

$V = 768.5$  (3) Å<sup>3</sup>

$Z = 2$

$F(000) = 546$

$D_x = 2.388$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 1000 reflections

$\theta = 3.1$ – $24.8^\circ$

$\mu = 3.44$  mm<sup>-1</sup>

$T = 293$  K

Block, colorless

$0.25 \times 0.22 \times 0.19$  mm

*Data collection*

Rigaku R-Axis RAPID  
diffractometer

Radiation source: fine-focus sealed tube  
graphite

Detector resolution: 10.00 pixels mm<sup>-1</sup>

$\omega$  scans

Absorption correction: empirical (using intensity  
measurements)

(ABSCOR; Higashi, 1995)

$T_{\min} = 0.480$ ,  $T_{\max} = 0.561$

7575 measured reflections

3485 independent reflections

3443 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.017$

$\theta_{\max} = 27.5^\circ$ ,  $\theta_{\min} = 3.1^\circ$

$h = -7 \rightarrow 8$

$k = -13 \rightarrow 13$

$l = -14 \rightarrow 14$

*Refinement*

Refinement on  $F^2$

Secondary atom site location: difference Fourier map

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.015$$

$$wR(F^2) = 0.041$$

$$S = 1.15$$

3485 reflections

225 parameters

4 restraints

Primary atom site location: structure-invariant direct methods

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.005P)^2 + 0.1027P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.56 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.71 \text{ e } \text{\AA}^{-3}$$

Absolute structure: Flack (1983)

Flack parameter:  $-0.009(8)$

### Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Ce1	0.468643 (16)	0.636197 (16)	0.819119 (10)	0.00928 (4)
S1	1.00549 (10)	0.68984 (7)	0.75724 (6)	0.01204 (13)
S2	0.57509 (11)	0.41208 (7)	0.63748 (6)	0.01388 (14)
S3	0.57345 (10)	0.97128 (6)	0.93682 (6)	0.01258 (13)
O1	0.5483 (3)	0.8315 (2)	0.9420 (2)	0.0190 (4)
O2	0.3896 (3)	1.0370 (2)	0.97444 (19)	0.0198 (4)
O3	0.5087 (3)	0.5445 (2)	0.60536 (19)	0.0199 (4)
O4	0.6164 (4)	1.0119 (2)	0.81523 (19)	0.0249 (5)
O5	1.1298 (3)	0.5804 (2)	0.71992 (18)	0.0181 (4)
O6	0.5651 (3)	0.40504 (19)	0.77192 (18)	0.0172 (4)
O7	0.8387 (2)	0.6423 (3)	0.82503 (16)	0.0198 (4)
O8	1.1447 (3)	0.7667 (2)	0.83878 (18)	0.0160 (4)
O9	0.7341 (3)	1.01219 (19)	1.02740 (19)	0.0178 (4)
O10	0.9302 (3)	0.7643 (2)	0.6533 (2)	0.0264 (5)
O11	0.4326 (3)	0.3179 (2)	0.5804 (2)	0.0242 (5)
O12	0.7762 (3)	0.3875 (2)	0.6036 (2)	0.0276 (5)
O1W	0.4833 (4)	0.8105 (2)	0.66809 (19)	0.0199 (4)
H1F	0.529 (5)	0.879 (2)	0.696 (3)	0.024*
H1G	0.520 (5)	0.799 (3)	0.600 (2)	0.024*
N1	-0.0899 (4)	0.3572 (3)	0.9309 (2)	0.0214 (5)

## supplementary materials

---

H1A	-0.1665	0.3382	0.9912	0.026*
H1B	-0.1653	0.3921	0.8703	0.026*
H1C	0.0049	0.4127	0.9567	0.026*
N2	0.2464 (4)	0.1635 (2)	0.7422 (2)	0.0204 (6)
H2A	0.3274	0.1334	0.8039	0.025*
H2B	0.3257	0.1928	0.6856	0.025*
N3	-0.0952 (4)	-0.0122 (3)	0.5183 (2)	0.0245 (6)
H3A	-0.1851	0.0142	0.4609	0.029*
H3B	-0.1518	-0.0693	0.5653	0.029*
H3C	0.0079	-0.0486	0.4845	0.029*
C1	0.0052 (5)	0.2382 (3)	0.8884 (3)	0.0223 (6)
H1D	-0.0969	0.1769	0.8604	0.027*
H1E	0.0873	0.1992	0.9540	0.027*
C2	0.1330 (5)	0.2739 (3)	0.7870 (3)	0.0182 (6)
H2C	0.0477	0.3090	0.7208	0.022*
H2D	0.2268	0.3405	0.8147	0.022*
C3	0.1272 (5)	0.0532 (3)	0.6894 (3)	0.0234 (7)
H3D	0.0579	0.0108	0.7524	0.028*
H3E	0.2169	-0.0087	0.6558	0.028*
C4	-0.0234 (5)	0.0989 (3)	0.5919 (3)	0.0279 (8)
H4A	-0.1355	0.1398	0.6280	0.034*
H4B	0.0383	0.1615	0.5412	0.034*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Ce1	0.00800 (7)	0.00962 (6)	0.01010 (6)	-0.00022 (6)	-0.00020 (5)	-0.00028 (7)
S1	0.0082 (3)	0.0151 (3)	0.0127 (3)	-0.0003 (2)	-0.0003 (3)	0.0010 (2)
S2	0.0137 (3)	0.0150 (3)	0.0128 (3)	0.0005 (3)	-0.0002 (3)	-0.0029 (2)
S3	0.0141 (3)	0.0122 (3)	0.0113 (3)	-0.0004 (2)	-0.0007 (3)	-0.0016 (2)
O1	0.0245 (11)	0.0130 (10)	0.0190 (10)	-0.0016 (8)	-0.0016 (9)	-0.0026 (8)
O2	0.0165 (10)	0.0207 (10)	0.0216 (10)	0.0066 (8)	-0.0027 (9)	-0.0042 (8)
O3	0.0222 (11)	0.0194 (10)	0.0178 (10)	0.0059 (8)	-0.0012 (9)	0.0004 (8)
O4	0.0369 (13)	0.0251 (12)	0.0131 (10)	-0.0070 (10)	0.0043 (10)	0.0000 (8)
O5	0.0118 (9)	0.0214 (10)	0.0208 (10)	0.0010 (8)	-0.0019 (9)	-0.0082 (8)
O6	0.0212 (10)	0.0181 (10)	0.0118 (9)	-0.0005 (8)	-0.0022 (9)	0.0000 (8)
O7	0.0102 (7)	0.0279 (10)	0.0216 (8)	-0.0029 (11)	0.0029 (7)	0.0037 (12)
O8	0.0133 (9)	0.0177 (10)	0.0171 (9)	-0.0015 (7)	0.0012 (8)	-0.0040 (8)
O9	0.0142 (9)	0.0185 (10)	0.0202 (10)	-0.0012 (8)	-0.0033 (9)	-0.0053 (8)
O10	0.0243 (11)	0.0317 (13)	0.0220 (11)	0.0019 (10)	-0.0074 (10)	0.0097 (9)
O11	0.0277 (12)	0.0270 (12)	0.0176 (10)	-0.0095 (10)	-0.0012 (10)	-0.0062 (9)
O12	0.0199 (10)	0.0321 (13)	0.0317 (12)	0.0067 (9)	0.0077 (10)	-0.0061 (10)
O1W	0.0268 (11)	0.0179 (11)	0.0153 (10)	-0.0013 (10)	0.0034 (9)	0.0011 (8)
N1	0.0179 (13)	0.0301 (15)	0.0164 (11)	-0.0004 (10)	0.0017 (11)	0.0016 (10)
N2	0.0166 (11)	0.0171 (16)	0.0268 (12)	0.0019 (9)	-0.0046 (11)	-0.0027 (9)
N3	0.0334 (15)	0.0221 (14)	0.0174 (12)	-0.0037 (11)	-0.0025 (12)	0.0021 (10)
C1	0.0245 (16)	0.0212 (15)	0.0213 (14)	-0.0046 (12)	0.0027 (13)	0.0043 (11)
C2	0.0202 (14)	0.0151 (14)	0.0193 (13)	-0.0008 (11)	0.0025 (12)	-0.0001 (11)

C3	0.0255 (16)	0.0140 (14)	0.0300 (16)	0.0022 (11)	-0.0046 (14)	-0.0006 (12)
C4	0.0401 (19)	0.0169 (15)	0.0250 (15)	0.0030 (12)	-0.0111 (16)	-0.0025 (11)

*Geometric parameters (Å, °)*

Ce1—O7	2.4685 (17)	N1—C1	1.481 (4)
Ce1—O1W	2.474 (2)	N1—H1A	0.8900
Ce1—O1	2.484 (2)	N1—H1B	0.8900
Ce1—O5 <sup>i</sup>	2.518 (2)	N1—H1C	0.8900
Ce1—O6	2.551 (2)	N2—C2	1.479 (4)
Ce1—O8 <sup>i</sup>	2.575 (2)	N2—C3	1.493 (4)
Ce1—O3	2.586 (2)	N2—H2A	0.9000
Ce1—O9 <sup>ii</sup>	2.588 (2)	N2—H2B	0.9000
Ce1—O2 <sup>ii</sup>	2.631 (2)	N3—C4	1.476 (4)
S1—O10	1.451 (2)	N3—H3A	0.8900
S1—O7	1.470 (2)	N3—H3B	0.8900
S1—O5	1.483 (2)	N3—H3C	0.8900
S1—O8	1.487 (2)	C1—C2	1.502 (4)
S2—O12	1.441 (2)	C1—H1D	0.9700
S2—O11	1.478 (2)	C1—H1E	0.9700
S2—O3	1.482 (2)	C2—H2C	0.9700
S2—O6	1.499 (2)	C2—H2D	0.9700
S3—O4	1.460 (2)	C3—C4	1.504 (4)
S3—O1	1.465 (2)	C3—H3D	0.9700
S3—O9	1.483 (2)	C3—H3E	0.9700
S3—O2	1.488 (2)	C4—H4A	0.9700
O1W—H1F	0.828 (18)	C4—H4B	0.9700
O1W—H1G	0.813 (18)		
O7—Ce1—O1W	85.13 (8)	O11—S2—O3	109.76 (14)
O7—Ce1—O1	77.68 (8)	O11—S2—O3	109.76 (14)
O1W—Ce1—O1	75.90 (8)	O12—S2—O6	110.75 (13)
O7—Ce1—O5 <sup>i</sup>	152.47 (7)	O11—S2—O6	108.94 (13)
O1W—Ce1—O5 <sup>i</sup>	86.97 (8)	O11—S2—O6	108.94 (13)
O1—Ce1—O5 <sup>i</sup>	125.59 (7)	O3—S2—O6	104.62 (12)
O7—Ce1—O6	76.34 (8)	O12—S2—Ce1	122.98 (10)
O1W—Ce1—O6	121.95 (7)	O11—S2—Ce1	126.18 (10)
O1—Ce1—O6	146.61 (7)	O11—S2—Ce1	126.18 (10)
O5 <sup>i</sup> —Ce1—O6	85.70 (7)	O3—S2—Ce1	53.00 (8)
O7—Ce1—O8 <sup>i</sup>	146.03 (8)	O6—S2—Ce1	51.77 (8)
O1W—Ce1—O8 <sup>i</sup>	75.06 (7)	O4—S3—O1	110.70 (14)
O1—Ce1—O8 <sup>i</sup>	70.94 (7)	O4—S3—O1	110.70 (14)
O5 <sup>i</sup> —Ce1—O8 <sup>i</sup>	54.73 (6)	O4—S3—O9	111.45 (13)
O6—Ce1—O8 <sup>i</sup>	137.61 (6)	O4—S3—O9	111.45 (13)
O7—Ce1—O3	82.53 (7)	O1—S3—O9	109.74 (12)
O1W—Ce1—O3	68.78 (7)	O4—S3—O2	109.99 (14)
O1—Ce1—O3	140.66 (7)	O4—S3—O2	109.99 (14)



## supplementary materials

---

O5 <sup>i</sup> —Ce1—O3	70.03 (7)	O1—S3—O2	110.21 (14)
O6—Ce1—O3	54.67 (6)	O9—S3—O2	104.58 (12)
O8 <sup>i</sup> —Ce1—O3	114.22 (7)	O4—S3—Ce1 <sup>iv</sup>	130.80 (10)
O7—Ce1—O9 <sup>ii</sup>	124.15 (7)	O4—S3—Ce1 <sup>iv</sup>	130.80 (10)
O1W—Ce1—O9 <sup>ii</sup>	148.88 (7)	O1—S3—Ce1 <sup>iv</sup>	118.49 (10)
O1—Ce1—O9 <sup>ii</sup>	98.55 (7)	O9—S3—Ce1 <sup>iv</sup>	51.63 (9)
O5 <sup>i</sup> —Ce1—O9 <sup>ii</sup>	71.27 (7)	O2—S3—Ce1 <sup>iv</sup>	53.35 (8)
O6—Ce1—O9 <sup>ii</sup>	79.38 (7)	S3—O1—Ce1	144.17 (13)
O8 <sup>i</sup> —Ce1—O9 <sup>ii</sup>	74.23 (7)	S3—O2—Ce1 <sup>iv</sup>	99.66 (10)
O3—Ce1—O9 <sup>ii</sup>	120.64 (7)	S2—O3—Ce1	99.75 (10)
O7—Ce1—O2 <sup>ii</sup>	71.62 (7)	S1—O5—Ce1 <sup>iii</sup>	101.82 (10)
O1W—Ce1—O2 <sup>ii</sup>	148.08 (7)	S2—O6—Ce1	100.75 (10)
O1—Ce1—O2 <sup>ii</sup>	77.94 (7)	S1—O7—Ce1	141.80 (12)
O5 <sup>i</sup> —Ce1—O2 <sup>ii</sup>	123.49 (7)	S1—O8—Ce1 <sup>iii</sup>	99.21 (10)
O6—Ce1—O2 <sup>ii</sup>	74.22 (7)	S3—O9—Ce1 <sup>iv</sup>	101.66 (10)
O8 <sup>i</sup> —Ce1—O2 <sup>ii</sup>	112.91 (7)	Ce1—O1W—H1F	114 (3)
O3—Ce1—O2 <sup>ii</sup>	126.92 (7)	Ce1—O1W—H1G	123 (3)
O9 <sup>ii</sup> —Ce1—O2 <sup>ii</sup>	53.54 (6)	H1F—O1W—H1G	110 (3)
O7—Ce1—S1 <sup>i</sup>	163.95 (6)	C1—N1—H1A	109.5
O1W—Ce1—S1 <sup>i</sup>	78.82 (6)	C1—N1—H1B	109.5
O1—Ce1—S1 <sup>i</sup>	98.37 (5)	H1A—N1—H1B	109.5
O5 <sup>i</sup> —Ce1—S1 <sup>i</sup>	27.23 (5)	C1—N1—H1C	109.5
O6—Ce1—S1 <sup>i</sup>	112.13 (5)	H1A—N1—H1C	109.5
O8 <sup>i</sup> —Ce1—S1 <sup>i</sup>	27.55 (4)	H1B—N1—H1C	109.5
O3—Ce1—S1 <sup>i</sup>	91.41 (6)	C2—N2—C3	117.1 (2)
O9 <sup>ii</sup> —Ce1—S1 <sup>i</sup>	71.66 (5)	C2—N2—H2A	108.0
O2 <sup>ii</sup> —Ce1—S1 <sup>i</sup>	123.17 (5)	C3—N2—H2A	108.0
O7—Ce1—S2	76.78 (6)	C2—N2—H2B	108.0
O1W—Ce1—S2	94.98 (6)	C3—N2—H2B	108.0
O1—Ce1—S2	153.51 (6)	H2A—N2—H2B	107.3
O5 <sup>i</sup> —Ce1—S2	77.70 (5)	C4—N3—H3A	109.5
O6—Ce1—S2	27.48 (4)	C4—N3—H3B	109.5
O8 <sup>i</sup> —Ce1—S2	131.42 (5)	H3A—N3—H3B	109.5
O3—Ce1—S2	27.25 (5)	C4—N3—H3C	109.5
O9 <sup>ii</sup> —Ce1—S2	101.51 (5)	H3A—N3—H3C	109.5
O2 <sup>ii</sup> —Ce1—S2	100.43 (5)	H3B—N3—H3C	109.5
S1 <sup>i</sup> —Ce1—S2	104.25 (3)	N1—C1—C2	108.0 (2)
O7—Ce1—S3 <sup>ii</sup>	97.66 (6)	N1—C1—H1D	110.1
O1W—Ce1—S3 <sup>ii</sup>	164.83 (5)	C2—C1—H1D	110.1
O1—Ce1—S3 <sup>ii</sup>	90.07 (5)	N1—C1—H1E	110.1
O5 <sup>i</sup> —Ce1—S3 <sup>ii</sup>	96.83 (6)	C2—C1—H1E	110.1
O6—Ce1—S3 <sup>ii</sup>	73.10 (5)	H1D—C1—H1E	108.4

O8 <sup>i</sup> —Ce1—S3 <sup>ii</sup>	95.10 (5)	N2—C2—C1	112.9 (2)
O3—Ce1—S3 <sup>ii</sup>	126.32 (5)	N2—C2—H2C	109.0
O9 <sup>ii</sup> —Ce1—S3 <sup>ii</sup>	26.70 (4)	C1—C2—H2C	109.0
O2 <sup>ii</sup> —Ce1—S3 <sup>ii</sup>	26.99 (4)	N2—C2—H2D	109.0
S1 <sup>i</sup> —Ce1—S3 <sup>ii</sup>	97.89 (3)	C1—C2—H2D	109.0
S2—Ce1—S3 <sup>ii</sup>	100.18 (2)	H2C—C2—H2D	107.8
O10—S1—O7	110.58 (13)	N2—C3—C4	110.7 (2)
O10—S1—O5	111.01 (14)	N2—C3—H3D	109.5
O7—S1—O5	109.98 (15)	C4—C3—H3D	109.5
O10—S1—O8	111.51 (13)	N2—C3—H3E	109.5
O7—S1—O8	109.51 (12)	C4—C3—H3E	109.5
O5—S1—O8	104.08 (11)	H3D—C3—H3E	108.1
O10—S1—Ce1 <sup>iii</sup>	123.20 (11)	N3—C4—C3	109.2 (3)
O7—S1—Ce1 <sup>iii</sup>	126.20 (9)	N3—C4—H4A	109.8
O5—S1—Ce1 <sup>iii</sup>	50.96 (8)	C3—C4—H4A	109.8
O8—S1—Ce1 <sup>iii</sup>	53.24 (8)	N3—C4—H4B	109.8
O12—S2—O11	110.80 (15)	C3—C4—H4B	109.8
O12—S2—O11	110.80 (15)	H4A—C4—H4B	108.3
O12—S2—O3	111.77 (14)		
O7—Ce1—S2—O12	-5.76 (14)	O4—S3—O2—Ce1 <sup>iv</sup>	-126.59 (11)
O1W—Ce1—S2—O12	78.04 (14)	O4—S3—O2—Ce1 <sup>iv</sup>	-126.59 (11)
O1—Ce1—S2—O12	9.97 (17)	O1—S3—O2—Ce1 <sup>iv</sup>	111.08 (11)
O5 <sup>i</sup> —Ce1—S2—O12	163.83 (14)	O9—S3—O2—Ce1 <sup>iv</sup>	-6.81 (12)
O6—Ce1—S2—O12	-91.51 (16)	O12—S2—O3—Ce1	-115.66 (13)
O8 <sup>i</sup> —Ce1—S2—O12	152.48 (14)	O11—S2—O3—Ce1	120.98 (12)
O3—Ce1—S2—O12	93.70 (17)	O11—S2—O3—Ce1	120.98 (12)
O9 <sup>ii</sup> —Ce1—S2—O12	-128.49 (13)	O6—S2—O3—Ce1	4.23 (12)
O2 <sup>ii</sup> —Ce1—S2—O12	-73.88 (13)	O7—Ce1—O3—S2	75.58 (12)
S1 <sup>i</sup> —Ce1—S2—O12	157.75 (12)	O1W—Ce1—O3—S2	163.23 (14)
S3 <sup>ii</sup> —Ce1—S2—O12	-101.32 (13)	O1—Ce1—O3—S2	135.61 (11)
O7—Ce1—S2—O11	172.06 (13)	O5 <sup>i</sup> —Ce1—O3—S2	-102.14 (12)
O1W—Ce1—S2—O11	-104.14 (14)	O6—Ce1—O3—S2	-2.95 (8)
O1—Ce1—S2—O11	-172.22 (17)	O8 <sup>i</sup> —Ce1—O3—S2	-135.32 (10)
O5 <sup>i</sup> —Ce1—S2—O11	-18.35 (13)	O9 <sup>ii</sup> —Ce1—O3—S2	-49.90 (13)
O6—Ce1—S2—O11	86.31 (16)	O2 <sup>ii</sup> —Ce1—O3—S2	15.33 (15)
O8 <sup>i</sup> —Ce1—S2—O11	-29.70 (14)	S1 <sup>i</sup> —Ce1—O3—S2	-119.34 (10)
O3—Ce1—S2—O11	-88.48 (17)	S3 <sup>ii</sup> —Ce1—O3—S2	-18.47 (13)
O9 <sup>ii</sup> —Ce1—S2—O11	49.33 (13)	O1—S3—O4—O4	0.0 (7)
O2 <sup>ii</sup> —Ce1—S2—O11	103.93 (13)	O9—S3—O4—O4	0.0 (7)
S1 <sup>i</sup> —Ce1—S2—O11	-24.43 (12)	O2—S3—O4—O4	0.0 (7)
S3 <sup>ii</sup> —Ce1—S2—O11	76.49 (12)	Ce1 <sup>iv</sup> —S3—O4—O4	0.0 (8)
O7—Ce1—S2—O11	172.06 (13)	O10—S1—O5—Ce1 <sup>iii</sup>	116.35 (13)

## supplementary materials

---

O1W—Ce1—S2—O11	-104.14 (14)	O7—S1—O5—Ce1 <sup>iii</sup>	-120.96 (10)
O1—Ce1—S2—O11	-172.22 (17)	O8—S1—O5—Ce1 <sup>iii</sup>	-3.73 (13)
O5 <sup>i</sup> —Ce1—S2—O11	-18.35 (13)	O12—S2—O6—Ce1	116.27 (13)
O6—Ce1—S2—O11	86.31 (16)	O11—S2—O6—Ce1	-121.62 (12)
O8 <sup>i</sup> —Ce1—S2—O11	-29.70 (14)	O11—S2—O6—Ce1	-121.62 (12)
O3—Ce1—S2—O11	-88.48 (17)	O3—S2—O6—Ce1	-4.30 (12)
O9 <sup>ii</sup> —Ce1—S2—O11	49.33 (13)	O7—Ce1—O6—S2	-87.55 (11)
O2 <sup>ii</sup> —Ce1—S2—O11	103.93 (13)	O1W—Ce1—O6—S2	-12.30 (13)
S1 <sup>i</sup> —Ce1—S2—O11	-24.43 (12)	O1—Ce1—O6—S2	-127.41 (12)
S3 <sup>ii</sup> —Ce1—S2—O11	76.49 (12)	O5 <sup>i</sup> —Ce1—O6—S2	71.42 (11)
O7—Ce1—S2—O3	-99.46 (13)	O8 <sup>i</sup> —Ce1—O6—S2	91.15 (13)
O1W—Ce1—S2—O3	-15.66 (13)	O3—Ce1—O6—S2	2.92 (8)
O1—Ce1—S2—O3	-83.74 (16)	O9 <sup>ii</sup> —Ce1—O6—S2	143.15 (11)
O5 <sup>i</sup> —Ce1—S2—O3	70.13 (13)	O2 <sup>ii</sup> —Ce1—O6—S2	-161.97 (11)
O6—Ce1—S2—O3	174.79 (15)	S1 <sup>i</sup> —Ce1—O6—S2	78.11 (10)
O8 <sup>i</sup> —Ce1—S2—O3	58.78 (13)	S3 <sup>ii</sup> —Ce1—O6—S2	169.90 (10)
O9 <sup>ii</sup> —Ce1—S2—O3	137.81 (12)	O10—S1—O7—Ce1	3.9 (3)
O2 <sup>ii</sup> —Ce1—S2—O3	-167.59 (12)	O5—S1—O7—Ce1	-119.0 (2)
S1 <sup>i</sup> —Ce1—S2—O3	64.05 (11)	O8—S1—O7—Ce1	127.2 (2)
S3 <sup>ii</sup> —Ce1—S2—O3	164.97 (11)	Ce1 <sup>iii</sup> —S1—O7—Ce1	-174.67 (17)
O7—Ce1—S2—O6	85.75 (12)	O1W—Ce1—O7—S1	-16.4 (3)
O1W—Ce1—S2—O6	169.55 (12)	O1—Ce1—O7—S1	-93.0 (3)
O1—Ce1—S2—O6	101.48 (15)	O5 <sup>i</sup> —Ce1—O7—S1	57.5 (4)
O5 <sup>i</sup> —Ce1—S2—O6	-104.66 (12)	O6—Ce1—O7—S1	108.2 (3)
O8 <sup>i</sup> —Ce1—S2—O6	-116.01 (12)	O8 <sup>i</sup> —Ce1—O7—S1	-70.3 (3)
O3—Ce1—S2—O6	-174.79 (15)	O3—Ce1—O7—S1	52.8 (3)
O9 <sup>ii</sup> —Ce1—S2—O6	-36.98 (11)	O9 <sup>ii</sup> —Ce1—O7—S1	175.0 (2)
O2 <sup>ii</sup> —Ce1—S2—O6	17.63 (11)	O2 <sup>ii</sup> —Ce1—O7—S1	-174.2 (3)
S1 <sup>i</sup> —Ce1—S2—O6	-110.74 (10)	S1 <sup>i</sup> —Ce1—O7—S1	-15.7 (5)
S3 <sup>ii</sup> —Ce1—S2—O6	-9.81 (10)	S2—Ce1—O7—S1	79.9 (3)
O4—S3—O1—Ce1	-21.6 (3)	S3 <sup>ii</sup> —Ce1—O7—S1	178.6 (3)
O4—S3—O1—Ce1	-21.6 (3)	O10—S1—O8—Ce1 <sup>iii</sup>	-116.13 (12)
O9—S3—O1—Ce1	-145.0 (2)	O7—S1—O8—Ce1 <sup>iii</sup>	121.17 (12)
O2—S3—O1—Ce1	100.3 (2)	O5—S1—O8—Ce1 <sup>iii</sup>	3.62 (12)
Ce1 <sup>iv</sup> —S3—O1—Ce1	158.70 (17)	O4—S3—O9—Ce1 <sup>iv</sup>	125.76 (12)
O7—Ce1—O1—S3	93.1 (2)	O4—S3—O9—Ce1 <sup>iv</sup>	125.76 (12)
O1W—Ce1—O1—S3	5.1 (2)	O1—S3—O9—Ce1 <sup>iv</sup>	-111.24 (12)
O5 <sup>i</sup> —Ce1—O1—S3	-70.6 (3)	O2—S3—O9—Ce1 <sup>iv</sup>	6.96 (13)
O6—Ce1—O1—S3	132.7 (2)	O12—S2—O11—O11	0.0 (2)
O8 <sup>i</sup> —Ce1—O1—S3	-73.7 (2)	O3—S2—O11—O11	0.0 (2)
O3—Ce1—O1—S3	31.6 (3)	O6—S2—O11—O11	0.00 (19)
O9 <sup>ii</sup> —Ce1—O1—S3	-143.6 (2)	Ce1—S2—O11—O11	0.00 (19)

O2 <sup>ii</sup> —Ce1—O1—S3	166.7 (2)	C3—N2—C2—C1	61.8 (3)
S1 <sup>i</sup> —Ce1—O1—S3	-71.1 (2)	N1—C1—C2—N2	176.3 (2)
S2—Ce1—O1—S3	77.5 (3)	C2—N2—C3—C4	53.3 (4)
S3 <sup>ii</sup> —Ce1—O1—S3	-169.0 (2)	N2—C3—C4—N3	163.5 (3)

Symmetry codes: (i)  $x-1, y, z$ ; (ii)  $-x+1, y-1/2, -z+2$ ; (iii)  $x+1, y, z$ ; (iv)  $-x+1, y+1/2, -z+2$ .

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O1W—H1F···O4	0.83 (2)	1.98 (2)	2.766 (3)	159 (4)
O1W—H1G···O11 <sup>v</sup>	0.81 (2)	2.06 (2)	2.850 (3)	164 (4)
N1—H1A···O8 <sup>ii</sup>	0.89	2.02	2.769 (3)	141.
N1—H1C···O9 <sup>ii</sup>	0.89	2.02	2.883 (3)	162.
N1—H1B···O6 <sup>i</sup>	0.89	2.05	2.852 (3)	150.
N2—H2B···O11	0.90	1.92	2.764 (4)	156.
N2—H2A···O2 <sup>vi</sup>	0.90	2.16	2.993 (3)	154.
N2—H2A···O4 <sup>vi</sup>	0.90	2.30	2.997 (3)	134.
N3—H3A···O5 <sup>vii</sup>	0.89	2.17	2.808 (3)	128.
N3—H3A···O3 <sup>viii</sup>	0.89	2.26	3.059 (4)	150.
N3—H3C···O12 <sup>vii</sup>	0.89	1.91	2.799 (4)	173.
N3—H3B···O10 <sup>ix</sup>	0.89	2.04	2.763 (4)	137.

Symmetry codes: (v)  $-x+1, y+1/2, -z+1$ ; (ii)  $-x+1, y-1/2, -z+2$ ; (i)  $x-1, y, z$ ; (vi)  $x, y-1, z$ ; (vii)  $-x+1, y-1/2, -z+1$ ; (viii)  $-x, y-1/2, -z+1$ ; (ix)  $x-1, y-1, z$ .

Fig. 1




Fig. 2

