

Bis(acetato- κ O)[1,2-bis(2-pyridyl-methoxy)benzene- κ^4 N,O,O',N']-copper(II) monohydrate

Shuang Zhang,^a Yu-Jie Wang,^{a,b} Dong-Sheng Ma,^b Ying Liu^b and Jin-Sheng Gao^{a,b*}

^aEngineering Research Center of Pesticide of Heilongjiang Province, Heilongjiang University, Harbin 150080, People's Republic of China, and ^bCollege of Chemistry and Materials Science, Heilongjiang University, Harbin 150080, People's Republic of China

Correspondence e-mail: hgf1000@163.com


Received 18 May 2010; accepted 19 May 2010

Key indicators: single-crystal X-ray study; $T = 291$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.034; wR factor = 0.089; data-to-parameter ratio = 17.0.

In the title compound, $[\text{Cu}(\text{CH}_3\text{COO})_2(\text{C}_{18}\text{H}_{16}\text{N}_2\text{O}_2)] \cdot \text{H}_2\text{O}$, the Cu^{II} ion is six-coordinated in a typically Jahn–Teller distorted octahedral environment defined by two O and two N atoms from the ligand and two O atoms from acetate anions. A linear chain structure propagating in [010] is built up by intermolecular O—H···O hydrogen bonds involving the uncoordinated water molecules.

Related literature

For the synthesis and for general background to flexible pyridyl-based ligands, see: Liu *et al.* (2010a,b).



Experimental

Crystal data

$[\text{Cu}(\text{C}_2\text{H}_3\text{O}_2)_2(\text{C}_{18}\text{H}_{16}\text{N}_2\text{O}_2)] \cdot \text{H}_2\text{O}$
 $M_r = 491.98$
Monoclinic, $P2_{1}/c$
 $a = 11.661$ (3) Å
 $b = 14.689$ (6) Å
 $c = 15.553$ (4) Å
 $\beta = 123.540$ (11)°
 $V = 2220.5$ (12) Å³

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 1.03$ mm⁻¹

$T = 291$ K
 $0.20 \times 0.19 \times 0.18$ mm

Data collection

Rigaku R-AXIS RAPID diffractometer
Absorption correction: multi-scan (*ABSCOR*; Higashi, 1995)
 $T_{\min} = 0.818$, $T_{\max} = 0.838$

21143 measured reflections
4960 independent reflections
3854 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.043$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.089$
 $S = 1.05$
4960 reflections

291 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.28$ e Å⁻³
 $\Delta\rho_{\min} = -0.28$ e Å⁻³

Table 1
Selected bond lengths (Å).

Cu1—O5	1.9529 (16)	Cu1—N2	2.0823 (18)
Cu1—O3	1.9571 (16)	Cu1—O2	2.4719 (15)
Cu1—N1	2.0580 (18)	Cu1—O1	2.5353 (16)

Table 2
Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
O7—H71···O4 ⁱ	0.85	1.92	2.772 (3)	174
O7—H72···O5	0.85	2.14	2.986 (2)	178

Symmetry code: (i) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalClear* (Rigaku/MSC, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors thank the Special Funds for the Research of Scientific and Technological Innovative Talents of Harbin Municipal Science and Technology Bureau (2006RFQXG093) and Heilongjiang University for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2778).

References

- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Liu, Y., Yan, P.-F., Yu, Y.-H., Hou, G.-F. & Gao, J.-S. (2010a). *Cryst. Growth Des.* **10**, 1559–1568.
- Liu, Y., Yan, P.-F., Yu, Y.-H., Hou, G.-F. & Gao, J.-S. (2010b). *Inorg. Chem. Commun.* **13**, 630–632.
- Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSC (2002). *CrystalClear*. Rigaku/MSC Inc., The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2010). E66, m701 [doi:10.1107/S160053681001874X]

Bis(acetato- κO)[1,2-bis(2-pyridylmethoxy)benzene- $\kappa^4 N,O,O',N'$]copper(II) monohydrate

S. Zhang, Y.-J. Wang, D.-S. Ma, Y. Liu and J.-S. Gao

Comment

N-Heterocyclic ligands coordinated with transition metal ions can form a variety of topology structures, including macrocycles, polyhedra and linear and helical polymers. Our group has report three kinds of flexible pyridyl-based ligands in previous reports (Liu *et al.* 20010a; Liu *et al.* 20010 b). As a part of our continuing work for bipyridyl aromatic ligands, we report the crystal structure of the title compound here.

1,2-Bis(pyridin-2-ylmethoxy)benzene molecule act as a chelating ligand to coordinate with Cu^{II} ion forming a discrete strucutre. Two acetate counter ions also coordinate to the center Cu^{II} ion, resulting the Cu^{II} ion is six-coordinated in quadrangular bipyramidal geometry (Figure 1, Table 1).

A one-dimensional chain structure is built up by intermolecular hydrogen bonds involving the uncoordinated water molecules (Figure 2, Table 2).

Experimental

The 1,2-Bis(pyridin-2-ylmethoxy)benzene was synthesized by the reaction of o-dihydroxybenzene and 2-chloromethylpyridine hydrochloride under nitrogen atmosphere and alkaline condition (Liu *et al.*, 2010a). Title ligand (0.58 g, 2 mmol) and Cu(CH₃COO)₂·H₂O (0.40 g, 2 mmol) were dissolved in 15 mL e ethanol, and then the mixture keep stirring for 30 minute. The resulting solution was filtered, and the filtrate was allowed to stand in a desiccator at room temperature for several days. Bule needle crystals were obtained with yield 34 %.

Refinement

H atoms bound to C atoms were placed in calculated positions and treated as riding on their parent atoms, with C—H = 0.93 Å (aromatic C), C—H = 0.97 Å (methene C), C—H = 0.98 Å (methyl C), and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. Water H atoms were initially located in a difference Fourier map but they were treated as riding on their parent atoms with O—H = 0.85 Å, and with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound, showing displacement ellipsoids at the 30% probability level for non-H atoms.




Fig. 2. A partial packing view, showing the one-dimensional hydrogen bonding structure. Dashed lines indicate the hydrogen bonds, no involving H atoms have been omitted.

Bis(acetato- κ O)[1,2-bis(2-pyridylmethoxy)benzene- κ^4N,O,O',N']copper(II) monohydrate

Crystal data

$[Cu(C_2H_3O_2)_2(C_{18}H_{16}N_2O_2)] \cdot H_2O$	$F(000) = 1020$
$M_r = 491.98$	$D_x = 1.472 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 15798 reflections
$a = 11.661 (3) \text{ \AA}$	$\theta = 3.0\text{--}27.5^\circ$
$b = 14.689 (6) \text{ \AA}$	$\mu = 1.03 \text{ mm}^{-1}$
$c = 15.553 (4) \text{ \AA}$	$T = 291 \text{ K}$
$\beta = 123.540 (11)^\circ$	Block, blue
$V = 2220.5 (12) \text{ \AA}^3$	$0.20 \times 0.19 \times 0.18 \text{ mm}$
$Z = 4$	

Data collection

Rigaku R-AXIS RAPID diffractometer	4960 independent reflections
Radiation source: fine-focus sealed tube graphite	3854 reflections with $I > 2\sigma(I)$
ω scan	$R_{\text{int}} = 0.043$
Absorption correction: multi-scan (<i>ABSCOR</i> ; Higashi, 1995)	$\theta_{\max} = 27.5^\circ, \theta_{\min} = 3.0^\circ$
$T_{\min} = 0.818, T_{\max} = 0.838$	$h = -14 \rightarrow 14$
21143 measured reflections	$k = -19 \rightarrow 19$
	$l = -19 \rightarrow 18$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.034$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.089$	H-atom parameters constrained
$S = 1.05$	$w = 1/[\sigma^2(F_o^2) + (0.0469P)^2 + 0.2754P]$ where $P = (F_o^2 + 2F_c^2)/3$
4960 reflections	$(\Delta/\sigma)_{\max} = 0.001$
291 parameters	$\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.19750 (2)	0.638427 (17)	0.351821 (17)	0.02824 (9)
O1	0.43387 (14)	0.58511 (11)	0.40262 (11)	0.0360 (3)
O2	0.23187 (14)	0.59585 (11)	0.21381 (11)	0.0379 (4)
O3	0.26556 (16)	0.75769 (10)	0.34343 (12)	0.0394 (4)
O4	0.13698 (19)	0.81284 (13)	0.39686 (14)	0.0531 (5)
O5	0.15262 (15)	0.51145 (10)	0.35769 (11)	0.0352 (3)
O6	0.00215 (19)	0.56093 (13)	0.39381 (14)	0.0533 (5)
N1	0.33015 (19)	0.64394 (11)	0.50978 (13)	0.0319 (4)
N2	0.01307 (18)	0.65453 (12)	0.20902 (13)	0.0315 (4)
C1	0.2732 (3)	0.66269 (16)	0.56418 (17)	0.0400 (5)
H1	0.1779	0.6651	0.5294	0.048*
C2	0.3524 (3)	0.67823 (18)	0.66939 (19)	0.0513 (6)
H2	0.3115	0.6921	0.7049	0.062*
C3	0.4931 (3)	0.6726 (2)	0.72009 (18)	0.0585 (7)
H3	0.5486	0.6836	0.7907	0.070*
C4	0.5530 (3)	0.65064 (18)	0.66613 (18)	0.0503 (6)
H4	0.6479	0.6443	0.7003	0.060*
C5	0.4679 (2)	0.63844 (15)	0.56047 (16)	0.0347 (5)
C6	0.5334 (2)	0.62142 (17)	0.50060 (17)	0.0400 (5)
H6A	0.5692	0.6780	0.4924	0.048*
H6B	0.6093	0.5790	0.5381	0.048*
C7	0.4697 (2)	0.58313 (15)	0.33102 (16)	0.0336 (5)
C8	0.6037 (2)	0.57611 (17)	0.35554 (19)	0.0432 (6)
H8	0.6772	0.5743	0.4242	0.052*

supplementary materials

C9	0.6276 (3)	0.57171 (19)	0.2769 (2)	0.0532 (7)
H9	0.7170	0.5673	0.2929	0.064*
C10	0.5183 (3)	0.57400 (19)	0.1761 (2)	0.0525 (7)
H10	0.5343	0.5704	0.1238	0.063*
C11	0.3831 (3)	0.58167 (17)	0.15022 (19)	0.0440 (6)
H11	0.3098	0.5833	0.0814	0.053*
C12	0.3594 (2)	0.58687 (14)	0.22831 (16)	0.0325 (5)
C13	0.1171 (2)	0.61228 (17)	0.11292 (16)	0.0387 (5)
H13A	0.0978	0.5588	0.0704	0.046*
H13B	0.1364	0.6627	0.0825	0.046*
C14	-0.0064 (2)	0.63471 (15)	0.11764 (16)	0.0328 (4)
C15	-0.1366 (2)	0.63615 (18)	0.02673 (17)	0.0467 (6)
H15	-0.1481	0.6222	-0.0359	0.056*
C16	-0.2490 (3)	0.65856 (19)	0.0306 (2)	0.0526 (7)
H16	-0.3370	0.6584	-0.0293	0.063*
C17	-0.2301 (2)	0.68105 (18)	0.12371 (19)	0.0454 (6)
H17	-0.3041	0.6975	0.1275	0.054*
C18	-0.0983 (2)	0.67845 (16)	0.21095 (17)	0.0369 (5)
H18	-0.0849	0.6937	0.2739	0.044*
C19	0.2749 (4)	0.91611 (19)	0.3717 (2)	0.0691 (9)
H19A	0.3666	0.9223	0.4319	0.104*
H19B	0.2168	0.9618	0.3728	0.104*
H19C	0.2763	0.9235	0.3110	0.104*
C20	0.2198 (3)	0.82288 (17)	0.37095 (16)	0.0406 (5)
C21	0.0351 (3)	0.40102 (19)	0.3948 (2)	0.0531 (6)
H21A	-0.0609	0.3877	0.3477	0.080*
H21B	0.0603	0.3919	0.4643	0.080*
H21C	0.0887	0.3614	0.3811	0.080*
C22	0.0622 (2)	0.49872 (16)	0.38126 (16)	0.0368 (5)
O7	0.1077 (2)	0.40557 (15)	0.17760 (17)	0.0729 (6)
H71	0.0314	0.3772	0.1505	0.109*
H72	0.1223	0.4355	0.2295	0.109*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.02986 (14)	0.03193 (14)	0.02291 (13)	-0.00047 (11)	0.01457 (11)	0.00016 (10)
O1	0.0302 (7)	0.0476 (10)	0.0287 (8)	-0.0046 (7)	0.0153 (7)	-0.0046 (7)
O2	0.0285 (7)	0.0602 (11)	0.0254 (7)	0.0034 (7)	0.0152 (7)	-0.0004 (7)
O3	0.0487 (9)	0.0370 (9)	0.0358 (8)	-0.0042 (7)	0.0254 (8)	-0.0001 (7)
O4	0.0606 (11)	0.0524 (11)	0.0561 (11)	0.0162 (9)	0.0384 (10)	0.0136 (9)
O5	0.0353 (8)	0.0358 (8)	0.0345 (8)	-0.0041 (6)	0.0193 (7)	-0.0006 (6)
O6	0.0604 (11)	0.0581 (12)	0.0542 (11)	0.0101 (9)	0.0397 (10)	0.0089 (9)
N1	0.0400 (10)	0.0301 (9)	0.0247 (8)	-0.0018 (8)	0.0173 (8)	0.0000 (7)
N2	0.0314 (9)	0.0360 (10)	0.0272 (9)	0.0014 (7)	0.0163 (8)	0.0017 (7)
C1	0.0528 (14)	0.0394 (13)	0.0335 (12)	-0.0003 (10)	0.0274 (12)	0.0016 (9)
C2	0.0784 (19)	0.0500 (15)	0.0348 (13)	-0.0035 (14)	0.0372 (14)	-0.0017 (11)
C3	0.077 (2)	0.0650 (18)	0.0212 (11)	-0.0089 (15)	0.0193 (13)	-0.0039 (11)

C4	0.0500 (15)	0.0558 (17)	0.0300 (12)	-0.0082 (12)	0.0125 (12)	0.0019 (11)
C5	0.0396 (12)	0.0301 (11)	0.0275 (10)	-0.0034 (9)	0.0141 (10)	0.0020 (9)
C6	0.0328 (11)	0.0457 (14)	0.0327 (11)	-0.0070 (10)	0.0126 (10)	-0.0034 (10)
C7	0.0347 (11)	0.0347 (12)	0.0363 (12)	0.0012 (9)	0.0226 (10)	0.0000 (9)
C8	0.0326 (12)	0.0491 (15)	0.0467 (14)	0.0044 (10)	0.0213 (12)	0.0004 (11)
C9	0.0452 (14)	0.0641 (18)	0.0663 (18)	0.0055 (12)	0.0408 (15)	-0.0027 (14)
C10	0.0540 (15)	0.0651 (18)	0.0588 (17)	0.0042 (13)	0.0439 (15)	-0.0010 (13)
C11	0.0454 (13)	0.0550 (16)	0.0400 (13)	0.0024 (11)	0.0288 (12)	-0.0016 (11)
C12	0.0327 (11)	0.0340 (12)	0.0354 (11)	0.0003 (9)	0.0216 (10)	-0.0022 (9)
C13	0.0365 (12)	0.0523 (14)	0.0260 (11)	0.0034 (10)	0.0164 (10)	0.0011 (9)
C14	0.0341 (11)	0.0356 (12)	0.0274 (10)	0.0001 (9)	0.0163 (10)	0.0012 (9)
C15	0.0424 (13)	0.0627 (17)	0.0261 (11)	0.0087 (12)	0.0133 (11)	0.0013 (11)
C16	0.0332 (12)	0.074 (2)	0.0350 (13)	0.0103 (12)	0.0089 (11)	0.0045 (12)
C17	0.0366 (12)	0.0519 (15)	0.0456 (14)	0.0120 (11)	0.0215 (12)	0.0096 (11)
C18	0.0376 (12)	0.0398 (13)	0.0344 (11)	0.0070 (10)	0.0206 (11)	0.0040 (10)
C19	0.116 (3)	0.0406 (16)	0.0568 (17)	-0.0208 (16)	0.0513 (19)	-0.0090 (13)
C20	0.0542 (15)	0.0365 (13)	0.0269 (11)	0.0013 (11)	0.0199 (11)	0.0057 (9)
C21	0.0548 (15)	0.0511 (16)	0.0530 (16)	-0.0142 (13)	0.0295 (14)	0.0008 (12)
C22	0.0347 (11)	0.0441 (13)	0.0277 (11)	-0.0024 (10)	0.0146 (10)	0.0012 (9)
O7	0.0755 (14)	0.0811 (16)	0.0852 (15)	-0.0315 (12)	0.0590 (14)	-0.0353 (12)

Geometric parameters (Å, °)

Cu1—O5	1.9529 (16)	C7—C8	1.393 (3)
Cu1—O3	1.9571 (16)	C8—C9	1.398 (3)
Cu1—N1	2.0580 (18)	C8—H8	0.9300
Cu1—N2	2.0823 (18)	C9—C10	1.370 (4)
Cu1—O2	2.4719 (15)	C9—H9	0.9300
Cu1—O1	2.5353 (16)	C10—C11	1.401 (3)
O1—C7	1.389 (2)	C10—H10	0.9300
O1—C6	1.414 (3)	C11—C12	1.389 (3)
O2—C12	1.381 (2)	C11—H11	0.9300
O2—C13	1.411 (3)	C13—C14	1.519 (3)
O3—C20	1.280 (3)	C13—H13A	0.9700
O4—C20	1.244 (3)	C13—H13B	0.9700
O5—C22	1.307 (3)	C14—C15	1.390 (3)
O6—C22	1.231 (3)	C15—C16	1.385 (3)
N1—C5	1.345 (3)	C15—H15	0.9300
N1—C1	1.362 (3)	C16—C17	1.378 (3)
N2—C14	1.341 (3)	C16—H16	0.9300
N2—C18	1.362 (3)	C17—C18	1.378 (3)
C1—C2	1.383 (3)	C17—H17	0.9300
C1—H1	0.9300	C18—H18	0.9300
C2—C3	1.376 (4)	C19—C20	1.510 (3)
C2—H2	0.9300	C19—H19A	0.9600
C3—C4	1.394 (4)	C19—H19B	0.9600
C3—H3	0.9300	C19—H19C	0.9600
C4—C5	1.384 (3)	C21—C22	1.509 (3)
C4—H4	0.9300	C21—H21A	0.9600

supplementary materials

C5—C6	1.516 (3)	C21—H21B	0.9600
C6—H6A	0.9700	C21—H21C	0.9600
C6—H6B	0.9700	O7—H71	0.8527
C7—C12	1.392 (3)	O7—H72	0.8496
O5—Cu1—O3	170.76 (6)	C7—C8—H8	120.0
O5—Cu1—N1	91.48 (6)	C9—C8—H8	120.0
O3—Cu1—N1	88.82 (7)	C10—C9—C8	119.4 (2)
O5—Cu1—N2	90.74 (7)	C10—C9—H9	120.3
O3—Cu1—N2	92.58 (7)	C8—C9—H9	120.3
N1—Cu1—N2	157.06 (7)	C9—C10—C11	121.2 (2)
O5—Cu1—O2	88.58 (6)	C9—C10—H10	119.4
O3—Cu1—O2	84.40 (6)	C11—C10—H10	119.4
N1—Cu1—O2	132.15 (6)	C12—C11—C10	119.3 (2)
N2—Cu1—O2	70.73 (6)	C12—C11—H11	120.3
O5—Cu1—O1	87.92 (6)	C10—C11—H11	120.3
O3—Cu1—O1	83.47 (6)	O2—C12—C11	125.4 (2)
N1—Cu1—O1	70.70 (6)	O2—C12—C7	114.82 (17)
N2—Cu1—O1	132.21 (6)	C11—C12—C7	119.81 (19)
O2—Cu1—O1	61.48 (5)	O2—C13—C14	109.00 (16)
C7—O1—C6	116.14 (16)	O2—C13—H13A	109.9
C7—O1—Cu1	120.87 (12)	C14—C13—H13A	109.9
C6—O1—Cu1	108.85 (12)	O2—C13—H13B	109.9
C12—O2—C13	118.16 (16)	C14—C13—H13B	109.9
C12—O2—Cu1	123.96 (12)	H13A—C13—H13B	108.3
C13—O2—Cu1	114.41 (12)	N2—C14—C15	121.7 (2)
C20—O3—Cu1	112.84 (14)	N2—C14—C13	119.23 (18)
C22—O5—Cu1	115.36 (14)	C15—C14—C13	119.09 (19)
C5—N1—C1	118.86 (19)	C16—C15—C14	119.2 (2)
C5—N1—Cu1	124.32 (14)	C16—C15—H15	120.4
C1—N1—Cu1	116.53 (15)	C14—C15—H15	120.4
C14—N2—C18	118.30 (18)	C17—C16—C15	119.8 (2)
C14—N2—Cu1	125.17 (14)	C17—C16—H16	120.1
C18—N2—Cu1	116.19 (14)	C15—C16—H16	120.1
N1—C1—C2	122.2 (2)	C18—C17—C16	118.2 (2)
N1—C1—H1	118.9	C18—C17—H17	120.9
C2—C1—H1	118.9	C16—C17—H17	120.9
C3—C2—C1	118.3 (2)	N2—C18—C17	122.9 (2)
C3—C2—H2	120.9	N2—C18—H18	118.5
C1—C2—H2	120.9	C17—C18—H18	118.5
C2—C3—C4	120.3 (2)	C20—C19—H19A	109.5
C2—C3—H3	119.9	C20—C19—H19B	109.5
C4—C3—H3	119.9	H19A—C19—H19B	109.5
C5—C4—C3	118.4 (2)	C20—C19—H19C	109.5
C5—C4—H4	120.8	H19A—C19—H19C	109.5
C3—C4—H4	120.8	H19B—C19—H19C	109.5
N1—C5—C4	121.9 (2)	O4—C20—O3	124.2 (2)
N1—C5—C6	119.58 (18)	O4—C20—C19	120.5 (2)
C4—C5—C6	118.5 (2)	O3—C20—C19	115.3 (2)
O1—C6—C5	109.39 (18)	C22—C21—H21A	109.5

O1—C6—H6A	109.8	C22—C21—H21B	109.5
C5—C6—H6A	109.8	H21A—C21—H21B	109.5
O1—C6—H6B	109.8	C22—C21—H21C	109.5
C5—C6—H6B	109.8	H21A—C21—H21C	109.5
H6A—C6—H6B	108.2	H21B—C21—H21C	109.5
O1—C7—C12	115.01 (17)	O6—C22—O5	123.8 (2)
O1—C7—C8	124.8 (2)	O6—C22—C21	120.1 (2)
C12—C7—C8	120.17 (19)	O5—C22—C21	116.0 (2)
C7—C8—C9	120.0 (2)	H71—O7—H72	109.2

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O7—H71···O4 ⁱ	0.85	1.92	2.772 (3)	174
O7—H72···O5	0.85	2.14	2.986 (2)	178

Symmetry codes: (i) $-x, y-1/2, -z+1/2$.

supplementary materials

Fig. 1




Fig. 2

