

**(E)-4-Bromo-2-[(2-hydroxyphenyl)-
imino]methyl]phenolate**Naser Eltaher Eltayeb,^{a,‡} Siang Guan Teoh,^a Hoong-Kun
Fun^{b,*§} and Suchada Chantrapromma^{c¶}^aSchool of Chemical Sciences, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, ^bX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^cCrystal Materials Research Unit, Department of Chemistry, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand

Correspondence e-mail: hkfun@usm.my

Received 5 April 2010; accepted 26 April 2010

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.018; wR factor = 0.041; data-to-parameter ratio = 16.3.

The title compound, $\text{C}_{13}\text{H}_{10}\text{BrNO}_2$, crystallizes in a zwitterionic form. The zwitterion exists in a *trans* configuration about the $\text{C}=\text{N}$ bond and is almost planar, the dihedral angle between the two benzene rings being 2.29 (9)°. An intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond formed between the iminium NH^+ and the phenolate O^- atoms generates an $S(6)$ ring motif. In the crystal, the zwitterions are linked through $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds into chains along $[101]$ and these chains are further connected through $\text{C}-\text{H}\cdots\text{Br}$ interactions into a two-dimensional network perpendicular to (101) . $\text{C}\cdots\text{C}$ [3.572 (3)- 3.592 (3) Å] and $\text{C}\cdots\text{Br}$ [3.5633 (19)- 3.7339 (18) Å] short contacts are observed. The crystal studied was a twin with twin law $\bar{1}00, 0\bar{1}0, 001$ with a domain ratio of 0.09919 (2): 0.90081 (2).


Related literature

For bond-length data, see: Allen *et al.* (1987). For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For background to Schiff bases and their applications, see: Dao *et al.* (2000); Kagkelari *et al.* (2009); Karthikeyan *et al.* (2006); Sriram *et al.* (2006); Wei & Atwood (1998). For related structures, see: Eltayeb *et al.* (2009; 2010); Tan & Liu (2009). For the stability of the temperature controller used in the data collection, see Cosier & Glazer, (1986).

[‡] On study leave from Department of Chemistry, International University of Africa, Khartoum, Sudan; e-mail: nasertaha90@hotmail.com.

[§] Thomson Reuters ResearcherID: A-3561-2009.

[¶] Thomson Reuters ResearcherID: A-5085-2009.

**Experimental***Crystal data*

$\text{C}_{13}\text{H}_{10}\text{BrNO}_2$
 $M_r = 291.12$
 Monoclinic, $P2_1$
 $a = 4.6387$ (3) Å
 $b = 18.9379$ (13) Å
 $c = 6.2270$ (4) Å
 $\beta = 90.144$ (3)°

$V = 547.02$ (6) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 3.74$ mm⁻¹
 $T = 100$ K
 $0.43 \times 0.14 \times 0.14$ mm

Data collection

Bruker APEXII DUO CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2009)
 $T_{\min} = 0.295$, $T_{\max} = 0.628$

8575 measured reflections
 3120 independent reflections
 3034 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.018$
 $wR(F^2) = 0.041$
 $S = 1.02$
 3120 reflections
 191 parameters
 1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.59$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.29$ e Å⁻³
 Absolute structure: Flack (1983), 1480 Friedel pairs
 Flack parameter: 0.027 (7)

Table 1

Hydrogen-bond geometry (Å, °).

| $D-\text{H}\cdots A$ | $D-\text{H}$ | $\text{H}\cdots A$ | $D\cdots A$ | $D-\text{H}\cdots A$ |
|---|--------------|--------------------|-------------|----------------------|
| $\text{O2}-\text{H1O2}\cdots\text{O1}^i$ | 0.82 | 1.76 | 2.5641 (19) | 169 |
| $\text{N1}-\text{H1N1}\cdots\text{O1}$ | 0.89 (3) | 1.84 (3) | 2.6129 (18) | 143 (3) |
| $\text{C7}-\text{H7A}\cdots\text{O2}$ | 0.95 (2) | 2.12 (2) | 2.794 (2) | 127.1 (18) |
| $\text{C11}-\text{H11A}\cdots\text{Br1}^{ii}$ | 0.96 (3) | 2.89 (3) | 3.6982 (19) | 143.1 (19) |

Symmetry codes: (i) $x + 1, y, z + 1$; (ii) $-x + 2, y + \frac{1}{2}, -z + 2$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

The authors thank the Malaysian Government, the Ministry of Science, Technology and Innovation (MOSTI) and Universiti Sains Malaysia for the RU research grants (PKIMIA/815002 and PKIMIA/811120). NEE would like to acknowledge Universiti Sains Malaysia for a post-doctoral fellowship. The International University of Africa (Sudan) is acknowledged for providing study leave to NEE. The authors thank the Malaysian Government and Universiti Sains Malaysia for the Research University Golden Goose grant No. 1001/PFIZIK/811012.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2436).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2009). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
- Dao, V.-T., Gaspard, C., Mayer, M., Werner, G. H., Nguyen, S. N. & Michelot, R. J. (2000). *Eur. J. Med. Chem.* **35**, 805–813.
- Eltayeb, N. E., Teoh, S. G., Chantrapromma, S. & Fun, H.-K. (2010). *Acta Cryst.* **E66**, o934–o935.
- Eltayeb, N. E., Teoh, S. G., Yeap, C. S., Fun, H.-K. & Adnan, R. (2009). *Acta Cryst.* **E65**, o2065–o2066.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Kagkelari, A., Papaefstahiou, G. S., Raptopoulou, C. P. & Zafiroopoulos, T. F. (2009). *Polyhedron*, **28**, 3279–3283.
- Karthikeyan, M. S., Prasad, D. J., Poojary, B., Bhat, K. S., Holla, B. S. & Kumari, N. S. (2006). *Bioorg. Med. Chem.* **14**, 7482–7489.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Sriram, D., Yogeewari, P., Myneedu, N. S. & Saraswat, V. (2006). *Bioorg. Med. Chem. Lett.* **16**, 2127–2129.
- Tan, G.-X. & Liu, X.-C. (2009). *Acta Cryst.* **E65**, o559.
- Wei, P. & Atwood, D. A. (1998). *Inorg. Chem.* **37**, 4934–4938.

supplementary materials

Acta Cryst. (2010). E66, o1262-o1263 [doi:10.1107/S1600536810015230]

(*E*)-4-Bromo-2-[(2-hydroxyphenyl)imino]methylphenolate

N. E. Eltayeb, S. G. Teoh, H.-K. Fun and S. Chantrapromma

Comment

Much attention has been given to Schiff base ligands due to their applications such as in coordination chemistry (Kagkelari *et al.*, 2009), chelated boron catalyst (Wei & Atwood, 1998), pharmacological activities, anticancer (Dao *et al.*, 2000), anti-HIV (Sriram *et al.*, 2006), antibacterial and antifungal (Karthikeyan *et al.*, 2006) activities. We have reported the crystal structures of Schiff base ligands which existed in a zwitterionic form *i.e.* 2-((*E*)-{2-[(*E*)-2,3-dihydroxybenzylideneamino]-5-methylphenyl}-imino)methyl)-6-hydroxyphenolate (Eltayeb *et al.*, 2009) and (*E*)-4-allyl-2-[(2-hydroxyphenyl)imino]methyl-6-methoxyphenolate (Eltayeb *et al.*, 2010). Herein we report the crystal structure of the title zwitterionic Schiff base ligand (I).

The molecule of (I) (Fig. 1), C₁₃H₉BrNO₂, crystallizes in a zwitterionic form with cationic iminium and anionic enolate, and exists in a *trans* configuration about the C=N bond [1.310 (2) Å]; the torsion angle C8–N1–C7–C6 is 179.25 (17)°. The molecule is almost planar with the dihedral angle between the two benzene rings of 2.31 (9)°. The hydroxy group is co-planar with the attached C8–C13 benzene ring with the r.m.s. of 0.0102 (2) Å for the seven non H atoms. Intramolecular N—H···O hydrogen bond between the NH⁺ and the phenolate O[−] generates an S(6) ring motif (Fig. 1; Table 1) which help to stabilize the planarity of the molecule (Bernstein *et al.*, 1995). The bond distances are in normal ranges (Allen *et al.*, 1987) and comparable with those found in related structures (Eltayeb *et al.*, 2009, 2010; Tan & Liu, 2009).

In the crystal packing (Fig. 2), the zwitterions are linked through O2–H1O2···O1 hydrogen bonds into chains along the [101] and these chains are further connected through C11—H11A···Br1 interactions into a 2-D network perpendicular to the (101)-plane. The crystal is stabilized by O—H···O and weak C—H···Br interactions (Table 1). C···C [3.572 (3)–3.592 (3) Å] and C···Br [3.5633 (19)–3.7339 (18) Å] short contacts are observed.

Experimental

The title compound was synthesized by adding 5-bromo-2-hydroxybenzaldehyde (0.402 g, 2 mmol) to a solution of 2-aminophenol (0.218 g, 2 mmol) in ethanol (30 ml). The mixture was refluxed with stirring for half an hour. The resultant yellow solution was filtered and the filtrate was evaporated to give a yellow solid product. Yellow needle-shaped single crystals of the title compound suitable for *x*-ray structure determination were obtained from ethanol by slow evaporation at room temperature after nine days.

Refinement

Hydroxyl H atom was placed in calculated positions with $d(\text{O—H}) = 0.82$ Å and the U_{iso} values was constrained to be $1.5U_{\text{eq}}$ of the carrier atom. The remaining H atoms were located from the difference map and isotropically refined. The highest residual electron density peak is located at 0.80 Å from Br1 and the deepest hole is located at 0.99 Å from Br1.

supplementary materials

The crystal studied was a twin with twin law $\bar{1} 0 0, 0 \bar{1} 0, 0 0 1$, leading to a distribution (refined BASF parameter) of 0.09919/0.90081 (2).

Figures




Fig. 1. The molecular structure of the title compound, with 50% probability displacement ellipsoids and the atom-numbering scheme. Hydrogen bond is shown as dashed lines.



Fig. 2. The crystal packing showing 2-D networks perpendicular to the (101)-plane.

(E)-4-Bromo-2-[(2-hydroxyphenyl)imino]methylphenolate

Crystal data

$C_{13}H_{10}BrNO_2$

$M_r = 291.12$

Monoclinic, $P2_1$

Hall symbol: P 2yb

$a = 4.6387$ (3) Å

$b = 18.9379$ (13) Å

$c = 6.2270$ (4) Å

$\beta = 90.144$ (3)°

$V = 547.02$ (6) Å³

$Z = 2$

$F(000) = 292$

$D_x = 1.773$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3120 reflections

$\theta = 1.1$ – 30.0 °

$\mu = 3.74$ mm⁻¹

$T = 100$ K

Needle, yellow

$0.43 \times 0.14 \times 0.14$ mm

Data collection

Bruker APEXII DUO CCD area-detector diffractometer

Radiation source: sealed tube graphite

φ and ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2009)

$T_{\min} = 0.295$, $T_{\max} = 0.628$

8575 measured reflections

3120 independent reflections

3034 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.026$

$\theta_{\max} = 30.0$ °, $\theta_{\min} = 1.1$ °

$h = -6 \rightarrow 6$

$k = -26 \rightarrow 26$

$l = -8 \rightarrow 8$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.018$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

| | |
|--|--|
| $wR(F^2) = 0.041$ | $w = 1/[\sigma^2(F_o^2) + (0.0035P)^2]$ |
| $S = 1.02$ | where $P = (F_o^2 + 2F_c^2)/3$ |
| 3120 reflections | $(\Delta/\sigma)_{\max} < 0.001$ |
| 191 parameters | $\Delta\rho_{\max} = 0.59 \text{ e } \text{\AA}^{-3}$ |
| 1 restraint | $\Delta\rho_{\min} = -0.29 \text{ e } \text{\AA}^{-3}$ |
| Primary atom site location: structure-invariant direct methods | Absolute structure: Flack (1983), 1480 Friedel pairs Flack parameter: 0.027 (7) |

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | x | y | z | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|------|-------------|---------------|--------------|----------------------------------|
| Br1 | 0.00526 (4) | 0.442840 (15) | 0.87673 (2) | 0.01554 (4) |
| O1 | 0.5191 (3) | 0.66300 (7) | 0.29008 (19) | 0.0152 (2) |
| O2 | 1.1314 (3) | 0.70088 (8) | 1.0194 (2) | 0.0166 (3) |
| H1O2 | 1.2697 | 0.6925 | 1.0976 | 0.025* |
| N1 | 0.8565 (3) | 0.69973 (8) | 0.6076 (2) | 0.0121 (3) |
| H1N1 | 0.802 (6) | 0.6976 (16) | 0.470 (5) | 0.026 (7)* |
| C1 | 0.4132 (4) | 0.61426 (10) | 0.4149 (3) | 0.0126 (3) |
| C2 | 0.1934 (4) | 0.56695 (10) | 0.3446 (3) | 0.0140 (3) |
| H2A | 0.133 (6) | 0.5756 (14) | 0.207 (4) | 0.020 (6)* |
| C3 | 0.0797 (4) | 0.51628 (10) | 0.4781 (3) | 0.0138 (3) |
| H3A | -0.064 (6) | 0.4852 (14) | 0.440 (4) | 0.021 (6)* |
| C4 | 0.1776 (4) | 0.51058 (9) | 0.6919 (3) | 0.0129 (3) |
| C5 | 0.3894 (4) | 0.55482 (9) | 0.7685 (3) | 0.0125 (3) |
| H5A | 0.450 (6) | 0.5535 (16) | 0.905 (4) | 0.026 (7)* |
| C6 | 0.5112 (4) | 0.60653 (9) | 0.6331 (3) | 0.0122 (3) |
| C7 | 0.7261 (4) | 0.65154 (9) | 0.7227 (3) | 0.0123 (3) |
| H7A | 0.787 (5) | 0.6470 (12) | 0.868 (4) | 0.009 (5)* |
| C8 | 1.0707 (4) | 0.74911 (9) | 0.6695 (3) | 0.0117 (3) |
| C9 | 1.2064 (4) | 0.74952 (9) | 0.8722 (3) | 0.0125 (3) |
| C10 | 1.4149 (4) | 0.80132 (10) | 0.9137 (3) | 0.0148 (3) |
| H10A | 1.494 (6) | 0.8047 (15) | 1.048 (5) | 0.022 (6)* |
| C11 | 1.4940 (4) | 0.84982 (10) | 0.7569 (3) | 0.0159 (3) |

supplementary materials

| | | | | |
|------|------------|--------------|------------|------------|
| H11A | 1.639 (6) | 0.8847 (13) | 0.786 (4) | 0.018 (6)* |
| C12 | 1.3635 (4) | 0.84799 (10) | 0.5548 (3) | 0.0160 (3) |
| H12A | 1.427 (7) | 0.8788 (17) | 0.453 (5) | 0.036 (8)* |
| C13 | 1.1523 (4) | 0.79797 (9) | 0.5130 (3) | 0.0139 (3) |
| H13A | 1.045 (5) | 0.7956 (13) | 0.368 (3) | 0.014 (6)* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|-------------|-------------|-------------|--------------|--------------|-------------|
| Br1 | 0.01876 (7) | 0.01522 (7) | 0.01262 (6) | -0.00425 (9) | -0.00307 (5) | 0.00324 (8) |
| O1 | 0.0161 (6) | 0.0183 (6) | 0.0113 (5) | -0.0031 (5) | -0.0034 (5) | 0.0033 (5) |
| O2 | 0.0167 (6) | 0.0224 (7) | 0.0106 (6) | -0.0044 (5) | -0.0054 (5) | 0.0042 (5) |
| N1 | 0.0122 (7) | 0.0140 (7) | 0.0102 (7) | -0.0004 (5) | -0.0033 (5) | 0.0003 (5) |
| C1 | 0.0122 (7) | 0.0150 (8) | 0.0106 (7) | 0.0001 (6) | -0.0012 (5) | -0.0002 (6) |
| C2 | 0.0156 (8) | 0.0172 (8) | 0.0092 (8) | -0.0007 (6) | -0.0030 (6) | -0.0007 (6) |
| C3 | 0.0136 (8) | 0.0141 (8) | 0.0137 (8) | -0.0022 (6) | -0.0024 (6) | -0.0018 (6) |
| C4 | 0.0142 (8) | 0.0120 (7) | 0.0126 (8) | -0.0007 (6) | -0.0003 (6) | 0.0015 (6) |
| C5 | 0.0150 (8) | 0.0137 (8) | 0.0087 (8) | 0.0011 (6) | -0.0030 (6) | 0.0007 (6) |
| C6 | 0.0116 (7) | 0.0140 (7) | 0.0109 (7) | 0.0014 (6) | -0.0014 (6) | -0.0021 (6) |
| C7 | 0.0123 (8) | 0.0139 (8) | 0.0108 (8) | 0.0003 (6) | -0.0020 (6) | -0.0007 (6) |
| C8 | 0.0114 (7) | 0.0114 (7) | 0.0122 (7) | 0.0001 (6) | -0.0033 (5) | -0.0010 (6) |
| C9 | 0.0127 (7) | 0.0146 (8) | 0.0103 (7) | 0.0003 (6) | -0.0012 (6) | -0.0007 (6) |
| C10 | 0.0146 (8) | 0.0174 (9) | 0.0123 (8) | -0.0012 (6) | -0.0044 (6) | -0.0020 (6) |
| C11 | 0.0141 (8) | 0.0154 (8) | 0.0182 (8) | -0.0021 (7) | -0.0016 (7) | -0.0017 (6) |
| C12 | 0.0172 (9) | 0.0147 (8) | 0.0161 (8) | -0.0007 (7) | -0.0011 (6) | 0.0022 (7) |
| C13 | 0.0128 (8) | 0.0160 (8) | 0.0129 (8) | 0.0008 (6) | -0.0026 (6) | 0.0011 (6) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|------------|-------------|-----------|-------------|
| Br1—C4 | 1.9011 (18) | C5—C6 | 1.411 (2) |
| O1—C1 | 1.304 (2) | C5—H5A | 0.89 (2) |
| O2—C9 | 1.346 (2) | C6—C7 | 1.424 (2) |
| O2—H1O2 | 0.8200 | C7—H7A | 0.95 (2) |
| N1—C7 | 1.310 (2) | C8—C13 | 1.397 (2) |
| N1—C8 | 1.417 (2) | C8—C9 | 1.409 (2) |
| N1—H1N1 | 0.89 (3) | C9—C10 | 1.401 (2) |
| C1—C2 | 1.425 (2) | C10—C11 | 1.391 (3) |
| C1—C6 | 1.439 (2) | C10—H10A | 0.92 (3) |
| C2—C3 | 1.376 (3) | C11—C12 | 1.396 (3) |
| C2—H2A | 0.91 (3) | C11—H11A | 0.96 (3) |
| C3—C4 | 1.409 (2) | C12—C13 | 1.387 (3) |
| C3—H3A | 0.92 (3) | C12—H12A | 0.91 (3) |
| C4—C5 | 1.376 (3) | C13—H13A | 1.03 (2) |
| C9—O2—H1O2 | 109.5 | N1—C7—C6 | 121.74 (16) |
| C7—N1—C8 | 129.42 (16) | N1—C7—H7A | 116.6 (14) |
| C7—N1—H1N1 | 111.3 (19) | C6—C7—H7A | 121.6 (14) |
| C8—N1—H1N1 | 119.2 (18) | C13—C8—C9 | 120.02 (16) |
| O1—C1—C2 | 122.15 (15) | C13—C8—N1 | 115.98 (15) |

| | | | |
|--------------|--------------|-----------------|--------------|
| O1—C1—C6 | 121.08 (16) | C9—C8—N1 | 123.97 (16) |
| C2—C1—C6 | 116.76 (16) | O2—C9—C10 | 122.23 (15) |
| C3—C2—C1 | 121.85 (16) | O2—C9—C8 | 119.39 (15) |
| C3—C2—H2A | 125.0 (16) | C10—C9—C8 | 118.39 (16) |
| C1—C2—H2A | 113.1 (17) | C11—C10—C9 | 121.08 (16) |
| C2—C3—C4 | 120.08 (16) | C11—C10—H10A | 119.3 (18) |
| C2—C3—H3A | 124.6 (16) | C9—C10—H10A | 119.6 (17) |
| C4—C3—H3A | 115.3 (16) | C10—C11—C12 | 120.15 (17) |
| C5—C4—C3 | 120.59 (16) | C10—C11—H11A | 120.5 (15) |
| C5—C4—Br1 | 120.14 (13) | C12—C11—H11A | 119.4 (15) |
| C3—C4—Br1 | 119.24 (13) | C13—C12—C11 | 119.38 (17) |
| C4—C5—C6 | 120.14 (16) | C13—C12—H12A | 122 (2) |
| C4—C5—H5A | 122 (2) | C11—C12—H12A | 118 (2) |
| C6—C5—H5A | 118 (2) | C12—C13—C8 | 120.94 (16) |
| C5—C6—C7 | 117.51 (15) | C12—C13—H13A | 122.2 (13) |
| C5—C6—C1 | 120.57 (16) | C8—C13—H13A | 116.8 (13) |
| C7—C6—C1 | 121.89 (16) | | |
| O1—C1—C2—C3 | -179.04 (17) | C1—C6—C7—N1 | -3.8 (3) |
| C6—C1—C2—C3 | 0.1 (3) | C7—N1—C8—C13 | -175.04 (17) |
| C1—C2—C3—C4 | 0.8 (3) | C7—N1—C8—C9 | 7.1 (3) |
| C2—C3—C4—C5 | -0.8 (3) | C13—C8—C9—O2 | -178.12 (16) |
| C2—C3—C4—Br1 | 177.31 (14) | N1—C8—C9—O2 | -0.3 (3) |
| C3—C4—C5—C6 | 0.0 (3) | C13—C8—C9—C10 | 2.3 (3) |
| Br1—C4—C5—C6 | -178.09 (13) | N1—C8—C9—C10 | -179.92 (17) |
| C4—C5—C6—C7 | 178.93 (16) | O2—C9—C10—C11 | 178.26 (18) |
| C4—C5—C6—C1 | 0.8 (3) | C8—C9—C10—C11 | -2.1 (3) |
| O1—C1—C6—C5 | 178.25 (17) | C9—C10—C11—C12 | 0.6 (3) |
| C2—C1—C6—C5 | -0.8 (2) | C10—C11—C12—C13 | 0.8 (3) |
| O1—C1—C6—C7 | 0.2 (3) | C11—C12—C13—C8 | -0.6 (3) |
| C2—C1—C6—C7 | -178.85 (16) | C9—C8—C13—C12 | -0.9 (3) |
| C8—N1—C7—C6 | 179.25 (17) | N1—C8—C13—C12 | -178.91 (16) |
| C5—C6—C7—N1 | 178.12 (16) | | |

Hydrogen-bond geometry (Å, °)

| <i>D</i> —H \cdots <i>A</i> | <i>D</i> —H | H \cdots <i>A</i> | <i>D</i> \cdots <i>A</i> | <i>D</i> —H \cdots <i>A</i> |
|-------------------------------------|-------------|---------------------|----------------------------|-------------------------------|
| O2—H1O2 \cdots O1 ⁱ | 0.82 | 1.76 | 2.5641 (19) | 169 |
| N1—H1N1 \cdots O1 | 0.89 (3) | 1.84 (3) | 2.6129 (18) | 143 (3) |
| C7—H7A \cdots O2 | 0.95 (2) | 2.12 (2) | 2.794 (2) | 127.1 (18) |
| C11—H11A \cdots Br1 ⁱⁱ | 0.96 (3) | 2.89 (3) | 3.6982 (19) | 143.1 (19) |

Symmetry codes: (i) $x+1, y, z+1$; (ii) $-x+2, y+1/2, -z+2$.

Fig. 1




Fig. 2

