

## 2-Chloro-N-(2-methylbenzoyl)benzenesulfonamide

P. A. Suchetan,<sup>a</sup> B. Thimme Gowda,<sup>a\*</sup> Sabine Foro<sup>b</sup> and Hartmut Fuess<sup>b</sup>

<sup>a</sup>Department of Chemistry, Mangalore University, Mangalagangotri 574 199, Mangalore, India, and <sup>b</sup>Institute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany  
Correspondence e-mail: gowdabt@yahoo.com


Received 29 April 2010; accepted 30 April 2010

Key indicators: single-crystal X-ray study;  $T = 299\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$ ;  $R$  factor = 0.041;  $wR$  factor = 0.110; data-to-parameter ratio = 15.7.

In the title compound,  $\text{C}_{14}\text{H}_{12}\text{ClNO}_3\text{S}$ , the N—H bond is antiperiplanar to the C=O bond. The dihedral angle between the two aromatic rings is  $78.7(1)^\circ$ . The crystal structure features inversion-related dimers linked by pairs of N—H···O(S) hydrogen bonds.

### Related literature

For background to our study of the effect of ring and side-chain substitutions on the crystal structures of *N*-aryl sulfonamides and for related structures, see: Gowda *et al.* (2009, 2010a,b).



### Experimental

#### Crystal data

$\text{C}_{14}\text{H}_{12}\text{ClNO}_3\text{S}$	$c = 20.080(2)\text{ \AA}$
$M_r = 309.76$	$\beta = 95.586(8)^\circ$
Monoclinic, $P2_1/n$	$V = 1447.8(2)\text{ \AA}^3$
$a = 6.6086(5)\text{ \AA}$	$Z = 4$
$b = 10.9621(9)\text{ \AA}$	Mo $K\alpha$ radiation

$\mu = 0.41\text{ mm}^{-1}$   
 $T = 299\text{ K}$

$0.34 \times 0.32 \times 0.28\text{ mm}$

#### Data collection

Oxford Diffraction Xcalibur diffractometer with a Sapphire CCD detector  
Absorption correction: multi-scan (*CrysAlis RED*; Oxford)

Diffraction, 2009)  
 $T_{\min} = 0.872$ ,  $T_{\max} = 0.893$   
5395 measured reflections  
2917 independent reflections  
2592 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.017$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$   
 $wR(F^2) = 0.110$   
 $S = 1.07$   
2917 reflections  
186 parameters  
1 restraint

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.31\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.33\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1N}\cdots\text{O2}^i$	0.84 (2)	2.11 (2)	2.937 (2)	172 (2)

Symmetry code: (i)  $-x, -y + 1, -z + 1$ .

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2009); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

PAS thanks the Council of Scientific and Industrial Research (CSIR), Government of India, New Delhi, for the award of a research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5259).

### References

- Gowda, B. T., Foro, S., Suchetan, P. A. & Fuess, H. (2009). *Acta Cryst. E65*, o2516.
- Gowda, B. T., Foro, S., Suchetan, P. A. & Fuess, H. (2010a). *Acta Cryst. E66*, o794.
- Gowda, B. T., Foro, S., Suchetan, P. A. & Fuess, H. (2010b). *Acta Cryst. E66*, o326.
- Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, England.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

## **supplementary materials**

*Acta Cryst.* (2010). E66, o1281 [doi:10.1107/S1600536810015990]

## 2-Chloro-*N*-(2-methylbenzoyl)benzenesulfonamide

**P. A. Suchetan, B. T. Gowda, S. Foro and H. Fuess**

### Comment

In the present work, as a part of studying the effect of ring and the side chain substitutions on the crystal structures of *N*-aryl sulfonamides (Gowda *et al.*, 2009, 2010*a,b*), the structure of 2-chloro-*N*-(2-methylbenzoyl)benzenesulfonamide (I) has been determined. In the C—SO<sub>2</sub>—NH—C(O) segment, the N—H bond is *anti* to the C=O bond (Fig. 1), similar to those observed in *N*-(2-chlorobenzoyl)2-chlorobenzenesulfonamide (II) (Suchetan *et al.*, 2010), *N*-(benzoyl)benzenesulfonamide (III) (Gowda *et al.*, 2009), and *N*-(benzoyl)2-chlorobenzenesulfonamide (IV) (Gowda *et al.*, 2010*a*).

Further, the conformation of the C=O bond in the C—SO<sub>2</sub>—NH—C(O) segment of (I) is *syn* to the *ortho*-methyl group in the benzoyl ring, similar to that observed between the *ortho*-Cl and the C=O bond in (II).

The molecules are twisted at the S atom with the torsional angle of -64.0 (2) $^{\circ}$ , compared to those of 66.5 (2) $^{\circ}$  in (II), -66.9 (3) $^{\circ}$  in (III), and 66.7 (2) $^{\circ}$  in (IV).

The dihedral angles between the sulfonyl benzene ring and the —SO<sub>2</sub>—NH—C—O segment is 88.4 (1) $^{\circ}$ , compared to the values of 86.9 (1) $^{\circ}$  in (II), 86.5(0.1) in (III) and 87.3 (1) $^{\circ}$  in (IV). Furthermore, the dihedral angle between the sulfonyl and the benzoyl benzene rings is 78.7 (1) $^{\circ}$ , compared to the values of 76.9 (1) $^{\circ}$  in (II), of 80.3(0.1) in (III), 69.8 (1) $^{\circ}$  (molecule 1) and 89.8 (1) $^{\circ}$  (molecule 2) in (III) and 73.3 (1) $^{\circ}$  in (IV).

The packing of molecules linked by N—H···O(S) hydrogen bonds (Table 1) is shown in Fig. 2.

### Experimental

The title compound was prepared by refluxing a mixture of 2-methylbenzoic acid, 2-chlorobenzenesulfonamide and phosphorous oxy chloride for 3 h on a water bath. The resultant mixture was cooled and poured into ice cold water. The solid obtained was filtered, washed thoroughly with water and then dissolved in sodium bicarbonate solution. The compound was later reprecipitated by acidifying the filtered solution with dilute HCl. It was filtered, dried and recrystallized.

Prism like colourless single crystals of the title compound used in X-ray diffraction studies were obtained by slow evaporation of its toluene solution at room temperature.

### Refinement

The H atom of the NH group was located in a difference map and its coordinates were refined with a distance restraint of N—H = 0.86 (2) Å. The other H atoms were positioned with idealized geometry using a riding model with C—H = 0.93–0.96 Å. All H atoms were refined with isotropic displacement parameters set to 1.2 times of the *U*<sub>eq</sub> of the parent atom.

# supplementary materials

---

## Figures




Fig. 1. Molecular structure of the title compound, showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. Molecular packing in the title compound. Hydrogen bonds are shown as dashed lines.

## 2-Chloro-N-(2-methylbenzoyl)benzenesulfonamide

### Crystal data

C <sub>14</sub> H <sub>12</sub> ClNO <sub>3</sub> S	<i>F</i> (000) = 640
<i>M<sub>r</sub></i> = 309.76	<i>D<sub>x</sub></i> = 1.421 Mg m <sup>-3</sup>
Monoclinic, <i>P</i> 2 <sub>1</sub> /n	Mo <i>K</i> α radiation, $\lambda$ = 0.71073 Å
Hall symbol: -P 2yn	Cell parameters from 3866 reflections
<i>a</i> = 6.6086 (5) Å	$\theta$ = 2.7–27.8°
<i>b</i> = 10.9621 (9) Å	$\mu$ = 0.41 mm <sup>-1</sup>
<i>c</i> = 20.080 (2) Å	<i>T</i> = 299 K
$\beta$ = 95.586 (8)°	Prism, colourless
<i>V</i> = 1447.8 (2) Å <sup>3</sup>	0.34 × 0.32 × 0.28 mm
<i>Z</i> = 4	

### Data collection

Oxford Diffraction Xcalibur diffractometer with a Sapphire CCD detector	2917 independent reflections
Radiation source: fine-focus sealed tube graphite	2592 reflections with $I > 2\sigma(I)$
Rotation method data acquisition using ω and phi scans	$R_{\text{int}} = 0.017$
Absorption correction: multi-scan (CrysAlis RED; Oxford Diffraction, 2009)	$\theta_{\text{max}} = 26.4^\circ$ , $\theta_{\text{min}} = 2.8^\circ$
$T_{\text{min}} = 0.872$ , $T_{\text{max}} = 0.893$	$h = -7 \rightarrow 8$
5395 measured reflections	$k = -13 \rightarrow 7$
	$l = -25 \rightarrow 25$

## *Refinement*

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.041$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.110$	$w = 1/[\sigma^2(F_o^2) + (0.0502P)^2 + 0.7805P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.07$	$(\Delta/\sigma)_{\max} < 0.001$
2917 reflections	$\Delta\rho_{\max} = 0.31 \text{ e \AA}^{-3}$
186 parameters	$\Delta\rho_{\min} = -0.32 \text{ e \AA}^{-3}$
1 restraint	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001xF_c^2\lambda^3/\sin(2\theta)]^{1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.146 (5)

## *Special details*

**Experimental.** CrysAlis RED (Oxford Diffraction, 2009) Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm.

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

## *Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.2188 (3)	0.22154 (17)	0.45434 (9)	0.0389 (4)
C2	0.3529 (3)	0.2921 (2)	0.42162 (11)	0.0504 (5)
C3	0.5045 (4)	0.2357 (3)	0.38928 (13)	0.0717 (8)
H3	0.5940	0.2821	0.3667	0.086*
C4	0.5208 (4)	0.1097 (3)	0.39104 (14)	0.0771 (9)
H4	0.6237	0.0719	0.3702	0.092*
C5	0.3890 (4)	0.0403 (3)	0.42274 (13)	0.0658 (7)
H5	0.4005	-0.0443	0.4228	0.079*
C6	0.2392 (4)	0.0956 (2)	0.45452 (10)	0.0502 (5)
H6	0.1500	0.0480	0.4765	0.060*
C7	0.2598 (3)	0.30250 (17)	0.61222 (9)	0.0392 (4)
C8	0.3281 (3)	0.38164 (17)	0.67032 (9)	0.0389 (4)
C9	0.5185 (3)	0.3631 (2)	0.70605 (11)	0.0492 (5)

## supplementary materials

---

C10	0.5679 (4)	0.4362 (3)	0.76164 (13)	0.0661 (7)
H10	0.6937	0.4257	0.7860	0.079*
C11	0.4385 (4)	0.5231 (3)	0.78199 (13)	0.0680 (7)
H11	0.4759	0.5693	0.8201	0.082*
C12	0.2534 (4)	0.5422 (2)	0.74615 (12)	0.0608 (6)
H12	0.1660	0.6020	0.7594	0.073*
C13	0.1982 (3)	0.4717 (2)	0.69029 (10)	0.0472 (5)
H13	0.0731	0.4845	0.6658	0.057*
C14	0.6730 (4)	0.2723 (3)	0.68618 (15)	0.0696 (7)
H14A	0.6814	0.2770	0.6388	0.083*
H14B	0.6325	0.1915	0.6977	0.083*
H14C	0.8034	0.2905	0.7094	0.083*
N1	0.1386 (3)	0.36014 (14)	0.56089 (8)	0.0412 (4)
H1N	0.135 (3)	0.4361 (15)	0.5574 (11)	0.049*
O1	-0.0894 (2)	0.18713 (14)	0.52207 (8)	0.0559 (4)
O2	-0.0831 (2)	0.37563 (12)	0.45683 (8)	0.0497 (4)
O3	0.3005 (3)	0.19555 (13)	0.60869 (7)	0.0554 (4)
Cl1	0.34073 (12)	0.45035 (6)	0.42249 (4)	0.0807 (3)
S1	0.02324 (7)	0.28443 (4)	0.49721 (2)	0.03857 (18)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0448 (10)	0.0355 (10)	0.0351 (9)	0.0007 (8)	-0.0027 (7)	-0.0024 (7)
C2	0.0472 (11)	0.0577 (13)	0.0447 (11)	-0.0085 (10)	-0.0030 (9)	0.0012 (9)
C3	0.0462 (12)	0.117 (3)	0.0523 (13)	-0.0120 (14)	0.0068 (10)	-0.0090 (15)
C4	0.0600 (15)	0.108 (2)	0.0607 (15)	0.0241 (16)	-0.0054 (12)	-0.0369 (16)
C5	0.0772 (17)	0.0629 (16)	0.0551 (14)	0.0211 (13)	-0.0052 (12)	-0.0178 (12)
C6	0.0670 (13)	0.0384 (11)	0.0440 (11)	0.0063 (10)	0.0001 (9)	-0.0051 (9)
C7	0.0477 (10)	0.0337 (9)	0.0366 (9)	0.0017 (8)	0.0067 (8)	0.0037 (7)
C8	0.0473 (10)	0.0349 (9)	0.0351 (9)	-0.0035 (8)	0.0076 (8)	0.0031 (7)
C9	0.0507 (11)	0.0501 (12)	0.0465 (11)	-0.0030 (9)	0.0038 (9)	0.0045 (9)
C10	0.0616 (14)	0.0768 (18)	0.0574 (14)	-0.0082 (13)	-0.0074 (11)	-0.0048 (13)
C11	0.0824 (18)	0.0715 (17)	0.0497 (13)	-0.0146 (14)	0.0038 (12)	-0.0161 (12)
C12	0.0743 (15)	0.0569 (14)	0.0534 (13)	0.0010 (12)	0.0185 (11)	-0.0140 (11)
C13	0.0527 (11)	0.0465 (11)	0.0432 (10)	0.0024 (9)	0.0085 (9)	-0.0018 (9)
C14	0.0565 (14)	0.0680 (17)	0.0833 (18)	0.0118 (12)	0.0019 (13)	0.0033 (14)
N1	0.0570 (10)	0.0242 (7)	0.0411 (8)	0.0016 (7)	-0.0015 (7)	-0.0002 (6)
O1	0.0591 (9)	0.0400 (8)	0.0701 (10)	-0.0135 (7)	0.0146 (8)	-0.0011 (7)
O2	0.0509 (8)	0.0349 (7)	0.0597 (9)	0.0040 (6)	-0.0124 (7)	-0.0026 (6)
O3	0.0846 (11)	0.0320 (7)	0.0483 (8)	0.0118 (7)	-0.0006 (7)	0.0017 (6)
Cl1	0.0810 (5)	0.0573 (4)	0.1049 (6)	-0.0258 (3)	0.0148 (4)	0.0216 (4)
S1	0.0428 (3)	0.0266 (3)	0.0456 (3)	-0.00243 (18)	0.00085 (19)	-0.00064 (18)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

C1—C6	1.387 (3)	C9—C10	1.387 (3)
C1—C2	1.389 (3)	C9—C14	1.507 (3)
C1—S1	1.761 (2)	C10—C11	1.369 (4)

C2—C3	1.391 (4)	C10—H10	0.9300
C2—Cl1	1.737 (3)	C11—C12	1.373 (4)
C3—C4	1.386 (4)	C11—H11	0.9300
C3—H3	0.9300	C12—C13	1.382 (3)
C4—C5	1.361 (4)	C12—H12	0.9300
C4—H4	0.9300	C13—H13	0.9300
C5—C6	1.370 (3)	C14—H14A	0.9600
C5—H5	0.9300	C14—H14B	0.9600
C6—H6	0.9300	C14—H14C	0.9600
C7—O3	1.206 (2)	N1—S1	1.6478 (17)
C7—N1	1.394 (2)	N1—H1N	0.836 (16)
C7—C8	1.488 (3)	O1—S1	1.4188 (15)
C8—C13	1.393 (3)	O2—S1	1.4283 (14)
C8—C9	1.401 (3)		
C6—C1—C2	119.3 (2)	C11—C10—C9	122.5 (2)
C6—C1—S1	117.65 (17)	C11—C10—H10	118.7
C2—C1—S1	123.06 (16)	C9—C10—H10	118.7
C1—C2—C3	119.7 (2)	C10—C11—C12	120.0 (2)
C1—C2—Cl1	121.28 (18)	C10—C11—H11	120.0
C3—C2—Cl1	119.0 (2)	C12—C11—H11	120.0
C4—C3—C2	119.2 (3)	C11—C12—C13	119.5 (2)
C4—C3—H3	120.4	C11—C12—H12	120.3
C2—C3—H3	120.4	C13—C12—H12	120.3
C5—C4—C3	121.2 (2)	C12—C13—C8	120.6 (2)
C5—C4—H4	119.4	C12—C13—H13	119.7
C3—C4—H4	119.4	C8—C13—H13	119.7
C4—C5—C6	119.7 (3)	C9—C14—H14A	109.5
C4—C5—H5	120.2	C9—C14—H14B	109.5
C6—C5—H5	120.2	H14A—C14—H14B	109.5
C5—C6—C1	120.9 (2)	C9—C14—H14C	109.5
C5—C6—H6	119.5	H14A—C14—H14C	109.5
C1—C6—H6	119.5	H14B—C14—H14C	109.5
O3—C7—N1	120.83 (18)	C7—N1—S1	122.33 (13)
O3—C7—C8	124.07 (18)	C7—N1—H1N	121.6 (16)
N1—C7—C8	115.09 (16)	S1—N1—H1N	115.4 (16)
C13—C8—C9	120.23 (19)	O1—S1—O2	118.67 (10)
C13—C8—C7	119.36 (18)	O1—S1—N1	108.94 (9)
C9—C8—C7	120.38 (18)	O2—S1—N1	104.66 (8)
C10—C9—C8	117.2 (2)	O1—S1—C1	108.22 (9)
C10—C9—C14	118.8 (2)	O2—S1—C1	109.90 (9)
C8—C9—C14	123.9 (2)	N1—S1—C1	105.69 (9)
C6—C1—C2—C3	0.4 (3)	C8—C9—C10—C11	0.2 (4)
S1—C1—C2—C3	179.48 (17)	C14—C9—C10—C11	178.2 (3)
C6—C1—C2—Cl1	-177.48 (16)	C9—C10—C11—C12	-1.3 (4)
S1—C1—C2—Cl1	1.6 (2)	C10—C11—C12—C13	1.0 (4)
C1—C2—C3—C4	-0.8 (3)	C11—C12—C13—C8	0.2 (4)
Cl1—C2—C3—C4	177.1 (2)	C9—C8—C13—C12	-1.2 (3)
C2—C3—C4—C5	1.2 (4)	C7—C8—C13—C12	176.6 (2)

## supplementary materials

---


C3—C4—C5—C6	−1.1 (4)	O3—C7—N1—S1	7.1 (3)
C4—C5—C6—C1	0.6 (4)	C8—C7—N1—S1	−171.69 (14)
C2—C1—C6—C5	−0.2 (3)	C7—N1—S1—O1	52.08 (18)
S1—C1—C6—C5	−179.40 (17)	C7—N1—S1—O2	179.96 (16)
O3—C7—C8—C13	−143.2 (2)	C7—N1—S1—C1	−64.00 (18)
N1—C7—C8—C13	35.5 (3)	C6—C1—S1—O1	−3.36 (19)
O3—C7—C8—C9	34.6 (3)	C2—C1—S1—O1	177.51 (16)
N1—C7—C8—C9	−146.66 (19)	C6—C1—S1—O2	−134.38 (16)
C13—C8—C9—C10	1.0 (3)	C2—C1—S1—O2	46.49 (18)
C7—C8—C9—C10	−176.8 (2)	C6—C1—S1—N1	113.21 (16)
C13—C8—C9—C14	−176.9 (2)	C2—C1—S1—N1	−65.91 (18)
C7—C8—C9—C14	5.4 (3)		

### Hydrogen-bond geometry ( $\text{\AA}$ , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N1—H1N···O2 <sup>i</sup>	0.84 (2)	2.11 (2)	2.937 (2)	172 (2)

Symmetry codes: (i)  $-x, -y+1, -z+1$ .

Fig. 1



## supplementary materials

---

Fig. 2

