

1-(1-Decyl-2-oxoindolin-3-ylidene)thiosemicarbazide

Khalil Mamari,^a Abdelfettah Zerzouf,^b Hafid Zouihri,^c El Mokhtar Essassi^a and Seik Weng Ng^{d*}

^aLaboratoire de Chimie Organique Hétérocyclique, Pôle de Compétences Pharmacochimie, Université Mohammed V-Agdal, BP 1014 Avenue Ibn Batout, Rabat, Morocco, ^bLaboratoire de Chimie Organique et Etudes Physicochimiques, ENS Rabat, Morocco, ^cCNRST Division UATRS, Angle Allal Fassi/FAR, BP 8027 Hay Riad, Rabat, Morocco, and ^dDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 4 May 2010; accepted 5 May 2010

Key indicators: single-crystal X-ray study; $T = 180\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.042; wR factor = 0.135; data-to-parameter ratio = 24.0.

In the title 1-alkylisatin 3-thiosemicarbazone, $\text{C}_{19}\text{H}_{28}\text{N}_4\text{OS}$, the imine $\text{C}=\text{N}$ bond has a *Z* configuration, whereas the $\text{N}-\text{C}=\text{S}$ unit has an *E* conformation. In the crystal, molecules are connected by $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, forming zigzag chains running along the *b* axis.

Related literature

For background to *N*-substituted isatins and their derivatives, see: Bouhfid *et al.* (2008).



Experimental

Crystal data

$\text{C}_{19}\text{H}_{28}\text{N}_4\text{OS}$
 $M_r = 360.51$

Monoclinic, $P2_1/n$
 $a = 11.8856 (2)\text{ \AA}$

$b = 11.0055 (2)\text{ \AA}$
 $c = 15.1638 (3)\text{ \AA}$
 $\beta = 99.468 (1)^\circ$
 $V = 1956.51 (6)\text{ \AA}^3$
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.18\text{ mm}^{-1}$
 $T = 180\text{ K}$
 $0.23 \times 0.14 \times 0.12\text{ mm}$

Data collection

Bruker X8 APEXII diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.960$, $T_{\max} = 0.979$

26304 measured reflections
5715 independent reflections
4227 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.043$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.135$
 $S = 1.06$
5715 reflections
238 parameters
3 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.30\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.36\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}1-\text{H}1\cdots\text{O}1^i$	0.86 (1)	2.19 (1)	2.986 (2)	154 (2)
Symmetry code: (i) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$				

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *pubLCIF* (Westrip, 2010).

We thank the Université Mohammed V-Agdal and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5265).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bouhfid, R., Joly, N., Ohmani, F., Essassi, E. M., Massoui, M. & Martin, P. (2008). *Lett. Org. Chem.* pp. 3–7.
- Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**. Submitted.

supplementary materials

Acta Cryst. (2010). E66, o1333 [doi:10.1107/S1600536810016521]

1-(1-Decyl-2-oxoindolin-3-ylidene)thiosemicarbazide

K. Mamari, A. Zerzouf, H. Zouihri, E. M. Essassi and S. W. Ng

Comment

N-Substituted isatins (Bouhfid *et al.*, 2008) represent a large family of heterocyclic compounds reported to show a wide range of useful medicinal properties. These condense readily with thiosemicarbazides to form crystalline thiosemicarbazones. The title 1-decylisatin derivative has a *Z* configuration around the imine C=N bond and *E* configuration the C(=S)=NH₂ bond (Scheme I, Fig. 1). The molecules form zigzag chains through N—H···O hydrogen bonds along the *b*-axis of the monoclinic unit cell (Fig. 2). The decyl chain adopts an irregular zigzag conformation; the irregular nature lead to a compact packing and there are no voids in the crystal.

Experimental

1-Decyl-isatin (1 g, 3.1 mmol) and thiosemicarbazide (0.31 g, 3.4 mmol) were dissolved in aqueous ethanol (50 ml); a few drops of glacial acetic acid were added. The mixture was heated for 4 hours. Yellow crystals separated from the cool solution in 80% yield.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.95–0.99 Å) and were included in the refinement in the riding model approximation, with *U*(H) set to 1.2–1.5*U*_{eq}(C). The amino-H atoms were located in a difference Fourier map, and they were refined isotropically with the N—H distance restrained to 0.86±0.01 Å.

Figures




Fig. 1. Anisotropic displacement ellipsoid plot (Barbour, 2001) of the molecule of C₁₉H₂₈N₄OS at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.




Fig. 2. Packing diagram of the title compound with hydrogen bonds drawn as dashed lines.

supplementary materials

1-(1-Decyl-2-oxoindolin-3-ylidene)thiosemicarbazide

Crystal data

C ₁₉ H ₂₈ N ₄ OS	<i>F</i> (000) = 776
<i>M_r</i> = 360.51	<i>D_x</i> = 1.224 Mg m ⁻³
Monoclinic, <i>P2₁/n</i>	Mo <i>Kα</i> radiation, λ = 0.71073 Å
Hall symbol: -P 2yn	Cell parameters from 5658 reflections
<i>a</i> = 11.8856 (2) Å	θ = 2.3–28.7°
<i>b</i> = 11.0055 (2) Å	μ = 0.18 mm ⁻¹
<i>c</i> = 15.1638 (3) Å	<i>T</i> = 180 K
β = 99.468 (1)°	Prism, yellow
<i>V</i> = 1956.51 (6) Å ³	0.23 × 0.14 × 0.12 mm
<i>Z</i> = 4	

Data collection

Bruker X8 APEXII diffractometer	5715 independent reflections
Radiation source: fine-focus sealed tube graphite	4227 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.043$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$\theta_{\text{max}} = 30.0^\circ$, $\theta_{\text{min}} = 2.0^\circ$
$T_{\text{min}} = 0.960$, $T_{\text{max}} = 0.979$	$h = -16 \rightarrow 16$
26304 measured reflections	$k = -13 \rightarrow 15$
	$l = -18 \rightarrow 21$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.042$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.135$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.06$	$w = 1/[\sigma^2(F_o^2) + (0.0721P)^2 + 0.276P]$ where $P = (F_o^2 + 2F_c^2)/3$
5715 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
238 parameters	$\Delta\rho_{\text{max}} = 0.30 \text{ e \AA}^{-3}$
3 restraints	$\Delta\rho_{\text{min}} = -0.36 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.80459 (3)	0.78130 (4)	0.66953 (3)	0.03232 (12)
O1	0.66844 (10)	0.67702 (9)	0.91795 (7)	0.0330 (3)

N1	0.78312 (11)	1.00870 (12)	0.72565 (9)	0.0304 (3)
N2	0.73941 (11)	0.85612 (11)	0.81634 (8)	0.0277 (3)
N3	0.71786 (10)	0.93926 (10)	0.87721 (8)	0.0241 (2)
N4	0.61721 (10)	0.76378 (10)	1.04405 (8)	0.0232 (2)
C1	0.77480 (11)	0.89065 (12)	0.73872 (9)	0.0240 (3)
C2	0.68081 (11)	0.89632 (12)	0.94641 (9)	0.0221 (3)
C3	0.65089 (11)	0.96609 (12)	1.02047 (9)	0.0214 (3)
C4	0.65109 (12)	1.08920 (12)	1.03890 (10)	0.0250 (3)
H4	0.6761	1.1465	0.9994	0.030*
C5	0.61370 (12)	1.12684 (13)	1.11687 (10)	0.0289 (3)
H5	0.6136	1.2109	1.1310	0.035*
C6	0.57670 (12)	1.04303 (14)	1.17397 (10)	0.0296 (3)
H6	0.5518	1.0710	1.2268	0.035*
C7	0.57503 (12)	0.91890 (13)	1.15612 (9)	0.0266 (3)
H7	0.5491	0.8618	1.1953	0.032*
C8	0.61282 (11)	0.88257 (11)	1.07871 (9)	0.0214 (3)
C9	0.65662 (12)	0.76577 (12)	0.96510 (9)	0.0240 (3)
C10	0.58378 (12)	0.65187 (12)	1.08470 (10)	0.0258 (3)
H10A	0.5452	0.5974	1.0372	0.031*
H10B	0.5286	0.6716	1.1249	0.031*
C11	0.68480 (13)	0.58614 (14)	1.13747 (11)	0.0329 (3)
H11A	0.7232	0.6404	1.1852	0.040*
H11B	0.7402	0.5667	1.0974	0.040*
C12	0.64970 (14)	0.46857 (13)	1.17964 (11)	0.0319 (3)
H12A	0.5931	0.4257	1.1352	0.038*
H12B	0.7175	0.4154	1.1935	0.038*
C13	0.59920 (13)	0.48675 (12)	1.26434 (11)	0.0305 (3)
H13A	0.5286	0.5355	1.2500	0.037*
H13B	0.6539	0.5334	1.3079	0.037*
C14	0.57148 (13)	0.36767 (13)	1.30730 (11)	0.0314 (3)
H14A	0.6407	0.3161	1.3168	0.038*
H14B	0.5121	0.3243	1.2655	0.038*
C15	0.52981 (13)	0.38398 (14)	1.39615 (11)	0.0321 (3)
H15A	0.5876	0.4308	1.4369	0.039*
H15B	0.4588	0.4326	1.3860	0.039*
C16	0.50680 (14)	0.26574 (14)	1.44180 (11)	0.0341 (3)
H16A	0.5792	0.2204	1.4577	0.041*
H16B	0.4540	0.2153	1.3996	0.041*
C17	0.45517 (16)	0.28689 (16)	1.52591 (13)	0.0431 (4)
H17A	0.3889	0.3420	1.5111	0.052*
H17B	0.5123	0.3286	1.5706	0.052*
C18	0.41613 (17)	0.17156 (19)	1.56828 (14)	0.0511 (5)
H18A	0.3704	0.1947	1.6146	0.061*
H18B	0.3663	0.1242	1.5219	0.061*
C19	0.5135 (2)	0.09282 (19)	1.60994 (14)	0.0567 (5)
H19A	0.4836	0.0208	1.6363	0.085*
H19B	0.5625	0.1386	1.6567	0.085*
H19C	0.5579	0.0676	1.5641	0.085*
H11	0.8010 (16)	1.0349 (19)	0.6760 (9)	0.048 (6)*

supplementary materials

H12	0.7606 (16)	1.0593 (15)	0.7628 (11)	0.044 (5)*
H2	0.7284 (16)	0.7811 (10)	0.8277 (13)	0.044 (5)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0421 (2)	0.0309 (2)	0.0277 (2)	0.00066 (15)	0.01700 (16)	-0.00417 (15)
O1	0.0523 (7)	0.0207 (5)	0.0300 (6)	0.0001 (4)	0.0186 (5)	-0.0034 (4)
N1	0.0425 (7)	0.0262 (6)	0.0262 (6)	0.0024 (5)	0.0164 (6)	0.0043 (5)
N2	0.0427 (7)	0.0197 (5)	0.0247 (6)	0.0004 (5)	0.0175 (5)	0.0007 (5)
N3	0.0297 (6)	0.0216 (5)	0.0227 (6)	0.0011 (4)	0.0095 (5)	0.0007 (4)
N4	0.0316 (6)	0.0180 (5)	0.0222 (6)	0.0008 (4)	0.0106 (5)	0.0025 (4)
C1	0.0247 (6)	0.0266 (6)	0.0216 (6)	0.0011 (5)	0.0069 (5)	0.0015 (5)
C2	0.0268 (6)	0.0193 (6)	0.0214 (6)	0.0013 (5)	0.0074 (5)	0.0007 (5)
C3	0.0242 (6)	0.0199 (6)	0.0207 (6)	0.0006 (5)	0.0052 (5)	0.0003 (5)
C4	0.0287 (7)	0.0208 (6)	0.0262 (7)	-0.0004 (5)	0.0062 (5)	0.0009 (5)
C5	0.0350 (7)	0.0226 (6)	0.0293 (8)	0.0010 (5)	0.0064 (6)	-0.0063 (5)
C6	0.0346 (8)	0.0320 (7)	0.0233 (7)	0.0021 (6)	0.0081 (6)	-0.0056 (6)
C7	0.0323 (7)	0.0281 (7)	0.0209 (7)	0.0007 (5)	0.0087 (5)	0.0007 (5)
C8	0.0246 (6)	0.0198 (6)	0.0202 (6)	0.0020 (5)	0.0048 (5)	0.0005 (5)
C9	0.0306 (7)	0.0204 (6)	0.0226 (6)	0.0006 (5)	0.0091 (5)	0.0015 (5)
C10	0.0313 (7)	0.0202 (6)	0.0274 (7)	-0.0029 (5)	0.0095 (6)	0.0035 (5)
C11	0.0336 (7)	0.0268 (7)	0.0400 (9)	-0.0001 (6)	0.0107 (6)	0.0105 (6)
C12	0.0383 (8)	0.0219 (6)	0.0370 (8)	0.0034 (6)	0.0104 (7)	0.0079 (6)
C13	0.0341 (7)	0.0207 (6)	0.0373 (8)	-0.0011 (5)	0.0077 (6)	0.0058 (6)
C14	0.0350 (8)	0.0240 (7)	0.0361 (8)	0.0001 (6)	0.0082 (6)	0.0077 (6)
C15	0.0345 (8)	0.0281 (7)	0.0345 (8)	-0.0001 (6)	0.0076 (6)	0.0064 (6)
C16	0.0367 (8)	0.0304 (7)	0.0366 (9)	0.0013 (6)	0.0102 (7)	0.0080 (6)
C17	0.0451 (9)	0.0427 (10)	0.0453 (10)	0.0034 (7)	0.0186 (8)	0.0095 (8)
C18	0.0539 (11)	0.0584 (12)	0.0445 (11)	-0.0115 (9)	0.0185 (9)	0.0095 (9)
C19	0.0836 (15)	0.0492 (11)	0.0392 (10)	-0.0012 (10)	0.0159 (10)	0.0104 (9)

Geometric parameters (\AA , $^\circ$)

S1—C1	1.6723 (14)	C11—C12	1.531 (2)
O1—C9	1.2320 (17)	C11—H11A	0.9900
N1—C1	1.3204 (18)	C11—H11B	0.9900
N1—H11	0.864 (9)	C12—C13	1.518 (2)
N1—H12	0.865 (9)	C12—H12A	0.9900
N2—N3	1.3537 (16)	C12—H12B	0.9900
N2—C1	1.3673 (17)	C13—C14	1.5235 (19)
N2—H2	0.858 (9)	C13—H13A	0.9900
N3—C2	1.2924 (17)	C13—H13B	0.9900
N4—C9	1.3556 (17)	C14—C15	1.521 (2)
N4—C8	1.4132 (16)	C14—H14A	0.9900
N4—C10	1.4613 (17)	C14—H14B	0.9900
C2—C3	1.4521 (18)	C15—C16	1.520 (2)
C2—C9	1.5014 (18)	C15—H15A	0.9900
C3—C4	1.3833 (18)	C15—H15B	0.9900

C3—C8	1.4004 (18)	C16—C17	1.521 (2)
C4—C5	1.3928 (19)	C16—H16A	0.9900
C4—H4	0.9500	C16—H16B	0.9900
C5—C6	1.385 (2)	C17—C18	1.529 (3)
C5—H5	0.9500	C17—H17A	0.9900
C6—C7	1.392 (2)	C17—H17B	0.9900
C6—H6	0.9500	C18—C19	1.499 (3)
C7—C8	1.3831 (18)	C18—H18A	0.9900
C7—H7	0.9500	C18—H18B	0.9900
C10—C11	1.513 (2)	C19—H19A	0.9800
C10—H10A	0.9900	C19—H19B	0.9800
C10—H10B	0.9900	C19—H19C	0.9800
C1—N1—H11	119.6 (15)	C13—C12—C11	114.51 (13)
C1—N1—H12	119.8 (13)	C13—C12—H12A	108.6
H11—N1—H12	119.9 (19)	C11—C12—H12A	108.6
N3—N2—C1	121.24 (12)	C13—C12—H12B	108.6
N3—N2—H2	117.5 (13)	C11—C12—H12B	108.6
C1—N2—H2	121.3 (13)	H12A—C12—H12B	107.6
C2—N3—N2	115.75 (11)	C12—C13—C14	113.07 (12)
C9—N4—C8	110.65 (11)	C12—C13—H13A	109.0
C9—N4—C10	122.93 (11)	C14—C13—H13A	109.0
C8—N4—C10	126.41 (11)	C12—C13—H13B	109.0
N1—C1—N2	116.36 (12)	C14—C13—H13B	109.0
N1—C1—S1	125.81 (11)	H13A—C13—H13B	107.8
N2—C1—S1	117.83 (10)	C13—C14—C15	113.72 (13)
N3—C2—C3	126.45 (12)	C13—C14—H14A	108.8
N3—C2—C9	127.20 (12)	C15—C14—H14A	108.8
C3—C2—C9	106.33 (11)	C13—C14—H14B	108.8
C4—C3—C8	120.36 (12)	C15—C14—H14B	108.8
C4—C3—C2	132.97 (12)	H14A—C14—H14B	107.7
C8—C3—C2	106.64 (11)	C16—C15—C14	114.32 (13)
C3—C4—C5	118.21 (13)	C16—C15—H15A	108.7
C3—C4—H4	120.9	C14—C15—H15A	108.7
C5—C4—H4	120.9	C16—C15—H15B	108.7
C6—C5—C4	120.72 (13)	C14—C15—H15B	108.7
C6—C5—H5	119.6	H15A—C15—H15B	107.6
C4—C5—H5	119.6	C15—C16—C17	112.24 (14)
C5—C6—C7	121.85 (13)	C15—C16—H16A	109.2
C5—C6—H6	119.1	C17—C16—H16A	109.2
C7—C6—H6	119.1	C15—C16—H16B	109.2
C8—C7—C6	116.91 (13)	C17—C16—H16B	109.2
C8—C7—H7	121.5	H16A—C16—H16B	107.9
C6—C7—H7	121.5	C16—C17—C18	114.67 (16)
C7—C8—C3	121.95 (12)	C16—C17—H17A	108.6
C7—C8—N4	128.32 (12)	C18—C17—H17A	108.6
C3—C8—N4	109.70 (11)	C16—C17—H17B	108.6
O1—C9—N4	126.06 (12)	C18—C17—H17B	108.6
O1—C9—C2	127.26 (12)	H17A—C17—H17B	107.6
N4—C9—C2	106.67 (11)	C19—C18—C17	112.98 (16)

supplementary materials


N4—C10—C11	112.26 (12)	C19—C18—H18A	109.0
N4—C10—H10A	109.2	C17—C18—H18A	109.0
C11—C10—H10A	109.2	C19—C18—H18B	109.0
N4—C10—H10B	109.2	C17—C18—H18B	109.0
C11—C10—H10B	109.2	H18A—C18—H18B	107.8
H10A—C10—H10B	107.9	C18—C19—H19A	109.5
C10—C11—C12	112.20 (12)	C18—C19—H19B	109.5
C10—C11—H11A	109.2	H19A—C19—H19B	109.5
C12—C11—H11A	109.2	C18—C19—H19C	109.5
C10—C11—H11B	109.2	H19A—C19—H19C	109.5
C12—C11—H11B	109.2	H19B—C19—H19C	109.5
H11A—C11—H11B	107.9		
C1—N2—N3—C2	177.49 (13)	C10—N4—C8—C7	-1.2 (2)
N3—N2—C1—N1	-2.0 (2)	C9—N4—C8—C3	-0.06 (16)
N3—N2—C1—S1	177.49 (10)	C10—N4—C8—C3	-179.47 (12)
N2—N3—C2—C3	-179.58 (13)	C8—N4—C9—O1	-178.82 (14)
N2—N3—C2—C9	-1.7 (2)	C10—N4—C9—O1	0.6 (2)
N3—C2—C3—C4	0.7 (2)	C8—N4—C9—C2	0.34 (15)
C9—C2—C3—C4	-177.54 (15)	C10—N4—C9—C2	179.77 (12)
N3—C2—C3—C8	178.72 (13)	N3—C2—C9—O1	0.4 (2)
C9—C2—C3—C8	0.44 (14)	C3—C2—C9—O1	178.66 (14)
C8—C3—C4—C5	0.6 (2)	N3—C2—C9—N4	-178.74 (13)
C2—C3—C4—C5	178.39 (14)	C3—C2—C9—N4	-0.48 (15)
C3—C4—C5—C6	-0.4 (2)	C9—N4—C10—C11	83.30 (17)
C4—C5—C6—C7	-0.1 (2)	C8—N4—C10—C11	-97.36 (16)
C5—C6—C7—C8	0.4 (2)	N4—C10—C11—C12	-179.72 (12)
C6—C7—C8—C3	-0.2 (2)	C10—C11—C12—C13	-78.90 (18)
C6—C7—C8—N4	-178.24 (14)	C11—C12—C13—C14	-176.76 (13)
C4—C3—C8—C7	-0.4 (2)	C12—C13—C14—C15	175.33 (13)
C2—C3—C8—C7	-178.65 (12)	C13—C14—C15—C16	-177.44 (13)
C4—C3—C8—N4	178.04 (12)	C14—C15—C16—C17	-174.78 (14)
C2—C3—C8—N4	-0.25 (15)	C15—C16—C17—C18	172.62 (15)
C9—N4—C8—C7	178.21 (14)	C16—C17—C18—C19	69.9 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H12···S1 ⁱ	0.87 (1)	2.81 (1)	3.628 (1)	159 (2)
N1—H11···O1 ⁱ	0.86 (1)	2.19 (1)	2.986 (2)	154 (2)

Symmetry codes: (i) $-x+3/2, y+1/2, -z+3/2$.

Fig. 1



supplementary materials

Fig. 2

