

2-Hydroxy-N'-(2-hydroxy-4-methoxybenzylidene)-3-methylbenzohydrazide monohydrate

You-Yue Han,* Yong-Hong Li and Qiu-Rong Zhao

Department of Chemistry and Life Science, Chuzhou University, Chuzhou, Anhui 239000, People's Republic of China
Correspondence e-mail: hanyouyue@126.com


Received 26 April 2010; accepted 7 May 2010

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.057; wR factor = 0.136; data-to-parameter ratio = 16.3.

In the title compound, $\text{C}_{16}\text{H}_{16}\text{N}_2\text{O}_4\cdot\text{H}_2\text{O}$, the dihedral angle between the two benzene rings is $12.4(2)^\circ$ and the molecule adopts an *E* configuration with respect to the $\text{C}=\text{N}$ bond. There are intramolecular $\text{O}-\text{H}\cdots\text{N}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds in the hydrazone molecule, which both generate $S(6)$ rings. In the crystal structure, molecules are linked by $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, forming layers parallel to the *ab* plane. The crystal studied was a non-merohedral twin with a domain ratio of 0.887 (3):0.113 (3).

Related literature

For our previous studies on hydrazones and for background information, see: Han & Zhao (2010*a,b*). For reference bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{16}\text{N}_2\text{O}_4\cdot\text{H}_2\text{O}$
 $M_r = 318.32$
Monoclinic, $P2_1/c$
 $a = 4.488(1)\text{ \AA}$
 $b = 13.494(2)\text{ \AA}$
 $c = 26.089(3)\text{ \AA}$
 $\beta = 91.630(2)^\circ$

$V = 1579.3(5)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.10\text{ mm}^{-1}$
 $T = 298\text{ K}$
 $0.20 \times 0.20 \times 0.18\text{ mm}$

Data collection

Bruker SMART CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2007)
 $T_{\min} = 0.980$, $T_{\max} = 0.982$

3429 measured reflections
3429 independent reflections
1584 reflections with $I > 2\sigma(I)$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.057$
 $wR(F^2) = 0.136$
 $S = 0.82$
3429 reflections

211 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.17\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.19\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O4—H4 \cdots O3	0.82	1.80	2.528 (2)	148
O1—H1 \cdots N1	0.82	1.93	2.650 (2)	146
N2—H2 \cdots O5	0.90	2.01	2.899 (2)	172
O5—H5B \cdots O3 ⁱ	0.84	1.92	2.745 (2)	167
O5—H5A \cdots O1 ⁱⁱ	0.84	2.06	2.849 (3)	156

Symmetry codes: (i) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

This work was supported by the Applied Chemistry Key Subject of Anhui Province (No. 200802187 C).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5423).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Bruker (2007). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Han, Y.-Y. & Zhao, Q.-R. (2010*a*). *Acta Cryst. E66*, o1025.
Han, Y.-Y. & Zhao, Q.-R. (2010*b*). *Acta Cryst. E66*, o1026.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.

supplementary materials

Acta Cryst. (2010). E66, o1340 [doi:10.1107/S1600536810016855]

2-Hydroxy-N-(2-hydroxy-4-methoxybenzylidene)-3-methylbenzohydrazide monohydrate

Y.-Y. Han, Y.-H. Li and Q.-R. Zhao

Comment

As a continuation of our work on the structural characterization of hydrazones (Han & Zhao, 2010a,b), we report here the crystal structure of the title compound. The title compound, Fig. 1, consists of a hydrazone molecule and a water molecule of crystallization. The dihedral angle between the two benzene rings is $12.4(2)^\circ$. The molecule adopts an *E* configuration with respect to the C=N bond. There are intramolecular O—H···N and O—H···O hydrogen bonds in the hydrazone molecule (Table 1). All the bond lengths are within normal ranges (Allen *et al.*, 1987).

In the crystal structure, molecules are linked through intermolecular N—H···O and O—H···O hydrogen bonds (Table 1) to form layers parallel to the *ab* plane (Fig. 2).

Experimental

A mixture of 4-methoxysalicylaldehyde (0.152 g, 1 mmol) and 2-hydroxy-3-methylbenzohydrazide (0.166 g, 1 mmol) in 50 ml methanol was stirred at room temperature for 1 h. The mixture was filtered to remove impurities, and then left at room temperature. After a few days, colourless blocks of (I) were formed.

Refinement

The crystal turned out to be a non-merohedral twin (twin law: -1 0 0/0 -1 0/0 0.331 0 1) with a fractional contribution of the minor component of 0.113 (3). Amino H and water H atoms were located from a difference Fourier map and refined isotropically, with N—H, O—H, and H···H distances restrained to 0.90 (1), 0.85 (1), and 1.37 (2) Å, respectively. Other H atoms were positioned geometrically and refined using the riding-model approximation, with C—H = 0.93 or 0.96 Å, O—H = 0.82 Å, and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{methyl C and O})$.

Figures




Fig. 1. The molecular structure of (I) with 30% probability displacement ellipsoids for non-H atoms. Intramolecular hydrogen bonds are shown as dashed lines.




Fig. 2. The crystal packing of (I). Hydrogen bonds are shown as dashed lines.

supplementary materials

2-Hydroxy-N¹-(2-hydroxy-4-methoxybenzylidene)-3-methylbenzohydrazide monohydrate

Crystal data

C ₁₆ H ₁₆ N ₂ O ₄ ·H ₂ O	<i>F</i> (000) = 672
<i>M_r</i> = 318.32	<i>D_x</i> = 1.339 Mg m ⁻³
Monoclinic, <i>P</i> 2 ₁ /c	Mo <i>Kα</i> radiation, λ = 0.71073 Å
Hall symbol: -P 2ybc	Cell parameters from 1505 reflections
<i>a</i> = 4.488 (1) Å	θ = 2.7–24.3°
<i>b</i> = 13.494 (2) Å	μ = 0.10 mm ⁻¹
<i>c</i> = 26.089 (3) Å	<i>T</i> = 298 K
β = 91.630 (2)°	Block, colorless
<i>V</i> = 1579.3 (5) Å ³	0.20 × 0.20 × 0.18 mm
<i>Z</i> = 4	

Data collection

Bruker SMART CCD diffractometer	3429 independent reflections
Radiation source: fine-focus sealed tube graphite	1584 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.000$
Absorption correction: multi-scan (SADABS; Bruker, 2007)	$\theta_{\text{max}} = 27.0^\circ$, $\theta_{\text{min}} = 1.6^\circ$
$T_{\text{min}} = 0.980$, $T_{\text{max}} = 0.982$	$h = -5 \rightarrow 5$
3429 measured reflections	$k = -17 \rightarrow 17$
	$l = 0 \rightarrow 33$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.057$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.136$	H-atom parameters constrained
$S = 0.82$	$w = 1/[\sigma^2(F_o^2) + (0.0467P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
3429 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
211 parameters	$\Delta\rho_{\text{max}} = 0.17 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.19 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds

in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	-0.0603 (4)	0.34424 (11)	0.17742 (7)	0.0707 (5)
H1	0.0416	0.3619	0.2024	0.106*
O2	-0.7266 (4)	0.47988 (12)	0.05036 (6)	0.0682 (5)
O3	0.4757 (4)	0.34626 (11)	0.30448 (6)	0.0750 (6)
O4	0.8037 (4)	0.30326 (11)	0.38168 (7)	0.0774 (6)
H4	0.6857	0.2940	0.3575	0.116*
N1	0.2277 (4)	0.47228 (14)	0.23782 (7)	0.0548 (5)
N2	0.4287 (4)	0.50268 (14)	0.27598 (7)	0.0529 (5)
H2	0.4617	0.5681	0.2779	0.063*
C1	-0.1155 (5)	0.52186 (16)	0.17115 (8)	0.0501 (6)
C2	-0.1891 (5)	0.42542 (16)	0.15477 (8)	0.0507 (6)
C3	-0.3910 (5)	0.40878 (16)	0.11519 (8)	0.0528 (6)
H3	-0.4381	0.3443	0.1053	0.063*
C4	-0.5233 (5)	0.48799 (16)	0.09027 (8)	0.0516 (6)
C5	-0.4595 (6)	0.58423 (17)	0.10571 (9)	0.0579 (6)
H5	-0.5539	0.6375	0.0895	0.069*
C6	-0.2570 (5)	0.60000 (17)	0.14486 (9)	0.0571 (7)
H6	-0.2116	0.6648	0.1544	0.069*
C7	-0.7804 (7)	0.38437 (19)	0.02939 (10)	0.0779 (9)
H7A	-0.5970	0.3571	0.0177	0.117*
H7B	-0.9211	0.3894	0.0011	0.117*
H7C	-0.8597	0.3420	0.0552	0.117*
C8	0.0983 (5)	0.54235 (18)	0.21216 (9)	0.0533 (6)
H8	0.1437	0.6078	0.2203	0.064*
C9	0.5468 (6)	0.43567 (17)	0.30853 (9)	0.0527 (6)
C10	0.7606 (5)	0.46971 (15)	0.34903 (8)	0.0482 (6)
C11	0.8747 (6)	0.40063 (16)	0.38452 (9)	0.0560 (6)
C12	1.0734 (6)	0.42876 (19)	0.42418 (9)	0.0626 (7)
C13	1.1545 (6)	0.52613 (19)	0.42727 (10)	0.0713 (8)
H13	1.2865	0.5462	0.4534	0.086*
C14	1.0462 (6)	0.59542 (18)	0.39284 (10)	0.0693 (8)
H14	1.1057	0.6612	0.3959	0.083*
C15	0.8526 (6)	0.56823 (17)	0.35429 (9)	0.0586 (7)
H15	0.7808	0.6157	0.3312	0.070*
C16	1.1935 (7)	0.3513 (2)	0.46131 (10)	0.0915 (10)
H16A	1.3286	0.3820	0.4857	0.137*
H16B	1.2969	0.3011	0.4428	0.137*
H16C	1.0313	0.3218	0.4790	0.137*

supplementary materials

O5	0.4807 (5)	0.71679 (11)	0.27610 (6)	0.0787 (6)
H5A	0.3414	0.7383	0.2940	0.118*
H5B	0.5224	0.7569	0.2528	0.118*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0818 (13)	0.0494 (10)	0.0792 (12)	-0.0028 (9)	-0.0243 (10)	0.0088 (8)
O2	0.0772 (13)	0.0656 (11)	0.0607 (10)	0.0033 (10)	-0.0179 (10)	-0.0049 (8)
O3	0.1079 (16)	0.0447 (10)	0.0717 (12)	-0.0132 (10)	-0.0130 (10)	-0.0080 (8)
O4	0.1079 (16)	0.0432 (10)	0.0804 (13)	0.0016 (10)	-0.0083 (11)	0.0055 (8)
N1	0.0563 (13)	0.0572 (12)	0.0505 (11)	-0.0060 (10)	-0.0044 (10)	-0.0047 (10)
N2	0.0592 (13)	0.0466 (10)	0.0524 (12)	-0.0060 (10)	-0.0057 (10)	-0.0047 (9)
C1	0.0540 (15)	0.0509 (14)	0.0454 (13)	-0.0051 (12)	0.0009 (12)	-0.0036 (11)
C2	0.0554 (16)	0.0456 (13)	0.0512 (13)	-0.0011 (12)	0.0017 (12)	0.0048 (11)
C3	0.0545 (15)	0.0468 (13)	0.0568 (15)	-0.0027 (12)	-0.0045 (12)	-0.0056 (11)
C4	0.0558 (15)	0.0541 (14)	0.0447 (13)	0.0000 (12)	-0.0014 (12)	-0.0044 (11)
C5	0.0660 (17)	0.0483 (13)	0.0591 (15)	0.0080 (13)	-0.0030 (13)	0.0046 (11)
C6	0.0646 (17)	0.0449 (13)	0.0616 (16)	-0.0051 (12)	0.0000 (14)	-0.0026 (11)
C7	0.086 (2)	0.0762 (19)	0.0699 (18)	-0.0058 (17)	-0.0192 (15)	-0.0202 (15)
C8	0.0521 (15)	0.0521 (14)	0.0556 (14)	-0.0075 (12)	0.0005 (12)	-0.0052 (12)
C9	0.0694 (17)	0.0417 (13)	0.0470 (13)	-0.0014 (12)	0.0046 (12)	-0.0046 (11)
C10	0.0573 (15)	0.0429 (12)	0.0445 (13)	0.0017 (11)	0.0012 (11)	-0.0044 (10)
C11	0.0699 (18)	0.0465 (14)	0.0518 (14)	0.0044 (13)	0.0070 (13)	-0.0021 (11)
C12	0.0727 (19)	0.0635 (16)	0.0515 (15)	0.0114 (15)	-0.0028 (13)	0.0005 (13)
C13	0.078 (2)	0.0724 (18)	0.0620 (17)	0.0062 (16)	-0.0158 (15)	-0.0129 (14)
C14	0.086 (2)	0.0499 (14)	0.0707 (18)	-0.0079 (14)	-0.0158 (16)	-0.0095 (13)
C15	0.0698 (18)	0.0456 (13)	0.0601 (15)	0.0024 (13)	-0.0052 (13)	-0.0020 (11)
C16	0.112 (3)	0.092 (2)	0.0700 (19)	0.034 (2)	-0.0149 (17)	0.0079 (16)
O5	0.1172 (17)	0.0435 (9)	0.0748 (12)	0.0068 (10)	-0.0079 (11)	0.0063 (8)

Geometric parameters (\AA , $^\circ$)

O1—C2	1.365 (3)	C7—H7A	0.9600
O1—H1	0.8208	C7—H7B	0.9600
O2—C4	1.369 (3)	C7—H7C	0.9600
O2—C7	1.418 (3)	C8—H8	0.9300
O3—C9	1.252 (3)	C9—C10	1.480 (3)
O4—C11	1.354 (3)	C10—C15	1.398 (3)
O4—H4	0.8210	C10—C11	1.400 (3)
N1—C8	1.287 (3)	C11—C12	1.399 (3)
N1—N2	1.387 (2)	C12—C13	1.365 (3)
N2—C9	1.339 (3)	C12—C16	1.514 (3)
N2—H2	0.8961	C13—C14	1.375 (3)
C1—C6	1.400 (3)	C13—H13	0.9300
C1—C2	1.406 (3)	C14—C15	1.361 (3)
C1—C8	1.443 (3)	C14—H14	0.9300
C2—C3	1.373 (3)	C15—H15	0.9300
C3—C4	1.377 (3)	C16—H16A	0.9600

C3—H3	0.9300	C16—H16B	0.9600
C4—C5	1.387 (3)	C16—H16C	0.9600
C5—C6	1.364 (3)	O5—H5A	0.8417
C5—H5	0.9300	O5—H5B	0.8395
C6—H6	0.9300		
C2—O1—H1	109.3	N1—C8—C1	121.7 (2)
C4—O2—C7	117.94 (18)	N1—C8—H8	119.2
C11—O4—H4	109.6	C1—C8—H8	119.2
C8—N1—N2	115.52 (18)	O3—C9—N2	120.1 (2)
C9—N2—N1	119.55 (19)	O3—C9—C10	121.2 (2)
C9—N2—H2	124.6	N2—C9—C10	118.68 (19)
N1—N2—H2	115.8	C15—C10—C11	117.8 (2)
C6—C1—C2	116.7 (2)	C15—C10—C9	123.3 (2)
C6—C1—C8	120.1 (2)	C11—C10—C9	118.9 (2)
C2—C1—C8	123.2 (2)	O4—C11—C12	116.6 (2)
O1—C2—C3	117.17 (19)	O4—C11—C10	121.9 (2)
O1—C2—C1	121.21 (19)	C12—C11—C10	121.4 (2)
C3—C2—C1	121.6 (2)	C13—C12—C11	117.9 (2)
C2—C3—C4	119.7 (2)	C13—C12—C16	122.5 (2)
C2—C3—H3	120.2	C11—C12—C16	119.6 (2)
C4—C3—H3	120.2	C12—C13—C14	121.8 (2)
O2—C4—C3	124.5 (2)	C12—C13—H13	119.1
O2—C4—C5	115.0 (2)	C14—C13—H13	119.1
C3—C4—C5	120.5 (2)	C15—C14—C13	120.4 (2)
C6—C5—C4	119.4 (2)	C15—C14—H14	119.8
C6—C5—H5	120.3	C13—C14—H14	119.8
C4—C5—H5	120.3	C14—C15—C10	120.7 (2)
C5—C6—C1	122.2 (2)	C14—C15—H15	119.7
C5—C6—H6	118.9	C10—C15—H15	119.7
C1—C6—H6	118.9	C12—C16—H16A	109.5
O2—C7—H7A	109.5	C12—C16—H16B	109.5
O2—C7—H7B	109.5	H16A—C16—H16B	109.5
H7A—C7—H7B	109.5	C12—C16—H16C	109.5
O2—C7—H7C	109.5	H16A—C16—H16C	109.5
H7A—C7—H7C	109.5	H16B—C16—H16C	109.5
H7B—C7—H7C	109.5	H5A—O5—H5B	111.4
C8—N1—N2—C9	172.1 (2)	N1—N2—C9—C10	-179.92 (19)
C6—C1—C2—O1	179.1 (2)	O3—C9—C10—C15	178.3 (2)
C8—C1—C2—O1	0.3 (4)	N2—C9—C10—C15	-1.8 (4)
C6—C1—C2—C3	-0.4 (4)	O3—C9—C10—C11	-2.6 (4)
C8—C1—C2—C3	-179.2 (2)	N2—C9—C10—C11	177.2 (2)
O1—C2—C3—C4	-178.7 (2)	C15—C10—C11—O4	-178.4 (2)
C1—C2—C3—C4	0.8 (4)	C9—C10—C11—O4	2.5 (4)
C7—O2—C4—C3	-7.4 (4)	C15—C10—C11—C12	0.3 (4)
C7—O2—C4—C5	174.0 (2)	C9—C10—C11—C12	-178.9 (2)
C2—C3—C4—O2	179.9 (2)	O4—C11—C12—C13	178.5 (3)
C2—C3—C4—C5	-1.6 (4)	C10—C11—C12—C13	-0.2 (4)
O2—C4—C5—C6	-179.3 (2)	O4—C11—C12—C16	-0.6 (4)

supplementary materials


C3—C4—C5—C6	2.0 (4)	C10—C11—C12—C16	-179.3 (3)
C4—C5—C6—C1	-1.7 (4)	C11—C12—C13—C14	0.1 (4)
C2—C1—C6—C5	0.8 (4)	C16—C12—C13—C14	179.2 (3)
C8—C1—C6—C5	179.7 (2)	C12—C13—C14—C15	0.0 (5)
N2—N1—C8—C1	179.8 (2)	C13—C14—C15—C10	0.0 (4)
C6—C1—C8—N1	178.5 (2)	C11—C10—C15—C14	-0.2 (4)
C2—C1—C8—N1	-2.7 (4)	C9—C10—C15—C14	178.9 (2)
N1—N2—C9—O3	-0.1 (4)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	$H\cdots A$	$D\cdots A$	$D\cdots H\cdots A$
O4—H4 \cdots O3	0.82	1.80	2.528 (2)
O1—H1 \cdots N1	0.82	1.93	2.650 (2)
N2—H2 \cdots O5	0.90	2.01	2.899 (2)
O5—H5B \cdots O3 ⁱ	0.84	1.92	2.745 (2)
O5—H5A \cdots O1 ⁱⁱ	0.84	2.06	2.849 (3)

Symmetry codes: (i) $-x+1, y+1/2, -z+1/2$; (ii) $-x, y+1/2, -z+1/2$.

Fig. 1



supplementary materials

Fig. 2

