

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## (E)-N'-(4-Methoxybenzylidene)thio- phene-2-carbohydrazide

Yu-Feng Li<sup>a</sup> and Fang-Fang Jian<sup>b\*</sup>

<sup>a</sup>Microscale Science Institute, Department of Chemistry and Chemical Engineering, Weifang University, Weifang 261061, People's Republic of China, and <sup>b</sup>Microscale Science Institute, Weifang University, Weifang 261061, People's Republic of China  
Correspondence e-mail: liyufeng8111@163.com


Received 13 May 2010; accepted 14 May 2010

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.056;  $wR$  factor = 0.177; data-to-parameter ratio = 17.2.

In the title compound,  $\text{C}_{13}\text{H}_{12}\text{N}_2\text{O}_2\text{S}$ , the dihedral angle between the aromatic rings is  $15.20(11)^\circ$ . In the crystal, inversion dimers linked by pairs of  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds generate  $R_2^2(8)$  loops.

### Related literature

For a related structure, see: Li & Jian (2010).



### Experimental

#### Crystal data

$\text{C}_{13}\text{H}_{12}\text{N}_2\text{O}_2\text{S}$

$M_r = 260.31$

Monoclinic,  $P2_1/c$   
 $a = 16.106(3)$  Å  
 $b = 5.3292(11)$  Å  
 $c = 14.812(3)$  Å  
 $\beta = 104.91(3)^\circ$   
 $V = 1228.5(4)$  Å<sup>3</sup>

$Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.26$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.25 \times 0.22 \times 0.18$  mm

#### Data collection

Bruker SMART CCD  
diffractometer  
11253 measured reflections

2807 independent reflections  
2454 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.027$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$   
 $wR(F^2) = 0.177$   
 $S = 1.00$   
2807 reflections

163 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.74$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.53$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1A}\cdots\text{O2}^i$	0.86	2.12	2.963 (2)	167

Symmetry code: (i)  $-x, -y + 2, -z + 2$ .

Data collection: *SMART* (Bruker 1997); cell refinement: *SAINT* (Bruker 1997); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5449).

### References

- Bruker (1997). *SMART* and *SAINT*. Bruker AXS, Inc., Madison, Wisconsin, USA.  
Li, Y.-F. & Jian, F.-F. (2010). *Acta Cryst.* **E66**, o1398.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

**supplementary materials**

*Acta Cryst.* (2010). E66, o1400 [ doi:10.1107/S1600536810017836 ]

## (*E*)-*N'*-(4-Methoxybenzylidene)thiophene-2-carbohydrazide

Y.-F. Li and F.-F. Jian

### Experimental

A mixture of thiophene-2-carbohydrazide (0.10 mol), and 4-methoxybenzaldehyde (0.10 mol) was stirred in refluxing ethanol (10 ml) for 4 h to afford the title compound (0.079 mol, yield 79%). Colourless blocks of (I) were obtained by recrystallization from ethanol at room temperature.

### Refinement

H atoms were fixed geometrically and allowed to ride on their attached atoms, with C—H distances = 0.93–0.97 Å; N—H = 0.86 Å and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$  or  $1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$ .

### Figures




Fig. 1. The structure of (I) showing 30% probability displacement ellipsoids.

## (*E*)-*N'*-(4-Methoxybenzylidene)thiophene-2-carbohydrazide

### Crystal data

$\text{C}_{13}\text{H}_{12}\text{N}_2\text{O}_2\text{S}$

$M_r = 260.31$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 16.106$  (3) Å

$b = 5.3292$  (11) Å

$c = 14.812$  (3) Å

$\beta = 104.91$  (3)°

$V = 1228.5$  (4) Å<sup>3</sup>

$Z = 4$

$F(000) = 544$

$D_x = 1.407$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 2454 reflections

$\theta = 2.7$ – $25.3$ °

$\mu = 0.26$  mm<sup>-1</sup>

$T = 293$  K

Block, colorless

$0.25 \times 0.22 \times 0.18$  mm

### Data collection

Bruker SMART CCD  
diffractometer

Radiation source: fine-focus sealed tube  
graphite

2454 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.027$

$\theta_{\text{max}} = 27.5$ °,  $\theta_{\text{min}} = 3.3$ °

# supplementary materials

---

phi and  $\omega$  scans  $h = -20 \rightarrow 20$   
11253 measured reflections  $k = -6 \rightarrow 6$   
2807 independent reflections  $l = -19 \rightarrow 18$

## Refinement

Refinement on  $F^2$  Primary atom site location: structure-invariant direct methods  
Least-squares matrix: full Secondary atom site location: difference Fourier map  
 $R[F^2 > 2\sigma(F^2)] = 0.056$  Hydrogen site location: inferred from neighbouring sites  
 $wR(F^2) = 0.177$  H-atom parameters constrained  
 $S = 1.00$   $w = 1/[\sigma^2(F_o^2) + (0.1142P)^2 + 0.7797P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
2807 reflections  $(\Delta/\sigma)_{\max} = 0.001$   
163 parameters  $\Delta\rho_{\max} = 0.74 \text{ e } \text{Å}^{-3}$   
0 restraints  $\Delta\rho_{\min} = -0.53 \text{ e } \text{Å}^{-3}$

## Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{Å}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.19516 (4)	0.31168 (11)	0.98996 (4)	0.0456 (2)
N2	0.17079 (11)	0.7123 (3)	1.09840 (12)	0.0364 (4)
N1	0.09320 (10)	0.7857 (3)	1.04270 (12)	0.0372 (4)
H1A	0.0643	0.9002	1.0620	0.045*
O2	-0.00677 (9)	0.7645 (3)	0.90675 (11)	0.0437 (4)
C9	0.06111 (12)	0.6815 (4)	0.95773 (14)	0.0347 (4)
C8	0.19896 (13)	0.8393 (4)	1.17289 (15)	0.0381 (4)
H8A	0.1659	0.9700	1.1865	0.046*
C5	0.28237 (13)	0.7838 (4)	1.23759 (14)	0.0366 (4)
O1	0.52295 (11)	0.6947 (4)	1.42184 (13)	0.0608 (5)
C11	0.07350 (14)	0.3690 (4)	0.83349 (18)	0.0449 (5)
H11A	0.0269	0.4297	0.7878	0.054*
C10	0.10555 (11)	0.4692 (4)	0.92598 (13)	0.0344 (4)
C2	0.44404 (14)	0.7112 (4)	1.35954 (15)	0.0428 (5)
C3	0.41450 (15)	0.5483 (5)	1.28526 (16)	0.0480 (5)

H3A	0.4485	0.4149	1.2758	0.058*
C6	0.31271 (15)	0.9418 (5)	1.31370 (16)	0.0473 (5)
H6A	0.2785	1.0739	1.3239	0.057*
C7	0.39260 (16)	0.9065 (5)	1.37430 (17)	0.0520 (6)
H7A	0.4118	1.0136	1.4249	0.062*
C4	0.33415 (15)	0.5854 (4)	1.22531 (16)	0.0454 (5)
H4A	0.3144	0.4752	1.1757	0.054*
C13	0.19081 (14)	0.1074 (4)	0.90128 (16)	0.0447 (5)
H13A	0.2289	-0.0259	0.9053	0.054*
C12	0.12683 (14)	0.1562 (5)	0.82410 (17)	0.0457 (5)
H12A	0.1178	0.0615	0.7697	0.055*
C1	0.57849 (17)	0.5001 (7)	1.4092 (2)	0.0675 (8)
H1B	0.6315	0.5105	1.4569	0.101*
H1C	0.5898	0.5168	1.3489	0.101*
H1D	0.5518	0.3407	1.4131	0.101*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0484 (4)	0.0437 (4)	0.0437 (3)	0.0088 (2)	0.0098 (2)	0.0038 (2)
N2	0.0340 (8)	0.0363 (9)	0.0375 (9)	0.0022 (6)	0.0069 (7)	0.0034 (7)
N1	0.0319 (8)	0.0382 (9)	0.0407 (9)	0.0047 (6)	0.0079 (7)	0.0012 (7)
O2	0.0310 (7)	0.0480 (9)	0.0477 (9)	0.0055 (6)	0.0019 (6)	0.0001 (7)
C9	0.0301 (9)	0.0343 (10)	0.0404 (10)	-0.0017 (7)	0.0103 (7)	0.0048 (7)
C8	0.0390 (10)	0.0389 (10)	0.0375 (10)	0.0040 (8)	0.0122 (8)	0.0011 (8)
C5	0.0389 (10)	0.0369 (10)	0.0343 (10)	0.0005 (8)	0.0099 (8)	0.0016 (7)
O1	0.0453 (9)	0.0744 (13)	0.0518 (10)	0.0062 (8)	-0.0072 (7)	-0.0129 (9)
C11	0.0408 (11)	0.0411 (11)	0.0596 (13)	-0.0091 (9)	0.0253 (10)	-0.0152 (10)
C10	0.0287 (8)	0.0361 (10)	0.0380 (9)	-0.0016 (7)	0.0079 (7)	0.0030 (8)
C2	0.0385 (10)	0.0487 (12)	0.0376 (10)	-0.0005 (9)	0.0033 (8)	-0.0016 (9)
C3	0.0462 (12)	0.0447 (12)	0.0479 (12)	0.0092 (9)	0.0030 (9)	-0.0065 (10)
C6	0.0508 (12)	0.0478 (12)	0.0417 (11)	0.0087 (10)	0.0091 (9)	-0.0097 (9)
C7	0.0542 (13)	0.0543 (14)	0.0423 (11)	0.0024 (11)	0.0029 (10)	-0.0153 (10)
C4	0.0477 (11)	0.0401 (11)	0.0422 (11)	0.0043 (9)	0.0005 (9)	-0.0084 (9)
C13	0.0442 (11)	0.0382 (11)	0.0525 (12)	0.0007 (9)	0.0142 (9)	-0.0021 (9)
C12	0.0386 (11)	0.0480 (12)	0.0505 (12)	-0.0069 (9)	0.0116 (9)	-0.0118 (10)
C1	0.0423 (13)	0.086 (2)	0.0669 (17)	0.0125 (13)	0.0002 (11)	-0.0018 (15)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

S1—C13	1.694 (2)	C11—H11A	0.9300
S1—C10	1.727 (2)	C2—C7	1.382 (3)
N2—C8	1.274 (3)	C2—C3	1.386 (3)
N2—N1	1.366 (2)	C3—C4	1.382 (3)
N1—C9	1.351 (3)	C3—H3A	0.9300
N1—H1A	0.8600	C6—C7	1.379 (3)
O2—C9	1.239 (2)	C6—H6A	0.9300
C9—C10	1.479 (3)	C7—H7A	0.9300
C8—C5	1.466 (3)	C4—H4A	0.9300

## supplementary materials

C8—H8A	0.9300	C13—C12	1.353 (3)
C5—C4	1.387 (3)	C13—H13A	0.9300
C5—C6	1.391 (3)	C12—H12A	0.9300
O1—C2	1.368 (3)	C1—H1B	0.9600
O1—C1	1.413 (3)	C1—H1C	0.9600
C11—C10	1.437 (3)	C1—H1D	0.9600
C11—C12	1.451 (3)		
C13—S1—C10	91.43 (11)	C4—C3—C2	119.6 (2)
C8—N2—N1	115.96 (18)	C4—C3—H3A	120.2
C9—N1—N2	121.10 (17)	C2—C3—H3A	120.2
C9—N1—H1A	119.4	C7—C6—C5	121.3 (2)
N2—N1—H1A	119.4	C7—C6—H6A	119.4
O2—C9—N1	119.15 (19)	C5—C6—H6A	119.4
O2—C9—C10	120.17 (19)	C6—C7—C2	119.8 (2)
N1—C9—C10	120.68 (17)	C6—C7—H7A	120.1
N2—C8—C5	121.42 (19)	C2—C7—H7A	120.1
N2—C8—H8A	119.3	C3—C4—C5	121.3 (2)
C5—C8—H8A	119.3	C3—C4—H4A	119.3
C4—C5—C6	118.0 (2)	C5—C4—H4A	119.3
C4—C5—C8	123.29 (19)	C12—C13—S1	113.72 (18)
C6—C5—C8	118.65 (19)	C12—C13—H13A	123.1
C2—O1—C1	117.9 (2)	S1—C13—H13A	123.1
C10—C11—C12	107.8 (2)	C13—C12—C11	114.2 (2)
C10—C11—H11A	126.1	C13—C12—H12A	122.9
C12—C11—H11A	126.1	C11—C12—H12A	122.9
C11—C10—C9	120.20 (18)	O1—C1—H1B	109.5
C11—C10—S1	112.76 (16)	O1—C1—H1C	109.5
C9—C10—S1	127.02 (15)	H1B—C1—H1C	109.5
O1—C2—C7	115.6 (2)	O1—C1—H1D	109.5
O1—C2—C3	124.4 (2)	H1B—C1—H1D	109.5
C7—C2—C3	120.0 (2)	H1C—C1—H1D	109.5

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1A $\cdots$ O2 <sup>i</sup>	0.86	2.12	2.963 (2)	167

Symmetry codes: (i)  $-x, -y+2, -z+2$ .

Fig. 1

