

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-Chloro-*N*-(2-chlorobenzoyl)benzenesulfonamideB. Thimme Gowda,^{a*} Sabine Foro,^b P. A. Suchetan^a and Hartmut Fuess^b

^aDepartment of Chemistry, Mangalore University, Mangalagangothri 574 199, Mangalore, India, and ^bInstitute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany
Correspondence e-mail: gowdabt@yahoo.com


Received 19 May 2010; accepted 21 May 2010

Key indicators: single-crystal X-ray study; $T = 299$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.039; wR factor = 0.107; data-to-parameter ratio = 15.4.

In the structure of the title compound, $\text{C}_{13}\text{H}_9\text{Cl}_2\text{NO}_3\text{S}$, the conformation of the $\text{N}-\text{H}$ bond in the $\text{C}-\text{SO}_2-\text{NH}-\text{C}(\text{O})$ segment is *anti* to the $\text{C}=\text{O}$ bond. The molecule is twisted at the S atom with a torsion angle of 65.7 (2)°. The dihedral angle between the sulfonyl benzene ring and the $-\text{SO}_2-\text{NH}-\text{C}-\text{O}$ segment is 88.5 (1)°, and that between the sulfonyl and the benzoyl benzene rings is 58.0 (1)°. In the crystal, molecules are linked by pairs of $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, forming inversion dimers.

Related literature

For our study of the effect of ring and side-chain substituents on the crystal structures of *N*-aromatic sulfonamides and for related structures, see: Gowda *et al.* (2010); Suchetan *et al.* (2010*a,b,c*).



Experimental

Crystal data

$\text{C}_{13}\text{H}_9\text{Cl}_2\text{NO}_3\text{S}$
 $M_r = 330.17$
Triclinic, $P\bar{1}$
 $a = 6.3882$ (9) Å

$b = 10.311$ (1) Å
 $c = 11.171$ (1) Å
 $\alpha = 79.01$ (1)°
 $\beta = 74.47$ (1)°

$\gamma = 84.76$ (1)°
 $V = 695.32$ (13) Å³
 $Z = 2$
Mo $K\alpha$ radiation

$\mu = 0.62$ mm⁻¹
 $T = 299$ K
 $0.24 \times 0.20 \times 0.14$ mm

Data collection

Oxford Diffraction Xcalibur diffractometer with a Sapphire CCD detector
Absorption correction: multi-scan (*CrysAlis RED*; Oxford)

Diffraction, 2009)
 $T_{\min} = 0.865$, $T_{\max} = 0.918$
4417 measured reflections
2825 independent reflections
2378 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.016$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.107$
 $S = 1.06$
2825 reflections
184 parameters
1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.25$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.52$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1N}\cdots\text{O2}^i$	0.81 (2)	2.13 (2)	2.914 (2)	164 (2)

Symmetry code: (i) $-x + 1, -y, -z$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2009); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

PAS thanks the Council of Scientific and Industrial Research (CSIR), Government of India, New Delhi, for the award of a research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2765).

References

- Gowda, B. T., Foro, S., Suchetan, P. A. & Fuess, H. (2010). *Acta Cryst.* **E66**, o326.
Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Yarnton, England.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
Suchetan, P. A., Gowda, B. T., Foro, S. & Fuess, H. (2010*a*). *Acta Cryst.* **E66**, o766.
Suchetan, P. A., Gowda, B. T., Foro, S. & Fuess, H. (2010*b*). *Acta Cryst.* **E66**, o1040.
Suchetan, P. A., Gowda, B. T., Foro, S. & Fuess, H. (2010*c*). *Acta Cryst.* **E66**, o1253.

supplementary materials

Acta Cryst. (2010). E66, o1467 [doi:10.1107/S1600536810019057]

4-Chloro-*N*-(2-chlorobenzoyl)benzenesulfonamide

B. T. Gowda, S. Foro, P. A. Suchetan and H. Fuess

Comment

As a part of studying the effect of ring and the side chain substituents on the crystal structures of *N*-aromatic sulfonamides (Gowda *et al.*, 2010; Suchetan *et al.*, 2010*a,b,c*), the structure of *N*-(2-chlorobenzoyl)-4-chlorobenzenesulfonamide (I) has been determined. The conformation of the N—H bond in the C—SO₂—NH—C(O) segment is *anti* to the C=O bond (Fig. 1), similar to those observed in *N*-(2-chlorobenzoyl)-benzenesulfonamide (II) (Gowda *et al.*, 2010), *N*-(benzoyl)-4-chlorobenzenesulfonamide (III) (Suchetan *et al.*, 2010*a*), *N*-(2-chlorobenzoyl)-2-chlorobenzenesulfonamide (IV) (Suchetan *et al.*, 2010*b*) and *N*-(4-chlorobenzoyl)-4-chlorobenzenesulfonamide (V) (Suchetan *et al.*, 2010*c*).

Further, the conformation of the C=O bond in the C—SO₂—NH—C(O) segment of (I) is *syn* to the *ortho*-Cl in the benzoyl ring, similar to that observed in (II) and (IV).

The molecules are twisted at the *S* atom with the torsional angle of 65.7 (2)°, compared to those of -59.0 (2)° (molecule 1) and -67.3 (2)° (molecule 2) in (II), -70.0 (2)°, 61.3 (2)° in the two independent molecules of (III), 66.5 (2)° in (IV) and 67.5 (3)° in (V).

The dihedral angle between the sulfonyl benzene ring and the —SO₂—NH—C—O segment is 88.5 (1)°, compared to the values of 87.3 (1)° (molecule 1) and 73.3 (1)° (molecule 2) in (II), 72.0 (1)° & 77.3 (1)° in the two molecules of (III), 86.9 (1)° in (IV) and 79.0 (1)° in (V).

Furthermore, the dihedral angle between the sulfonyl and the benzoyl benzene rings is 58.0 (1)°, compared to the values of 69.8 (1)° (molecule 1) and 89.8 (1)° (molecule 2) in (II), 62.8 (1)° (molecule 1) and 78.6 (1)° (molecule 2) of (III), 76.9 (1)° in (IV) and 85.6 (1)° in (V).

The packing of molecules linked by of N—H···O(S) hydrogen bonds (Table 1) is shown in Fig. 2.

Experimental

The title compound was prepared by refluxing a mixture of 2-chlorobenzoic acid, 4-chlorobenzenesulfonamide and phosphorous oxy chloride for 3 h on a water bath. The resultant mixture was cooled and poured into ice cold water. The solid obtained was filtered, washed thoroughly with water and then dissolved in sodium bicarbonate solution. The compound was later reprecipitated by acidifying the filtered solution with dilute HCl. It was filtered, dried and recrystallized.

Prism like colourless single crystals of the title compound used in X-ray diffraction studies were obtained by slow evaporation of its toluene solution at room temperature.

Refinement

The H atoms of the NH groups were located in a difference map and later restrained to N—H = 0.86 (2) %Å. The other H atoms were positioned with idealized geometry using a riding model with C—H = 0.93 Å. All H atoms were refined with isotropic displacement parameters (set to 1.2 times of the U_{eq} of the parent atom).

Figures




Fig. 1. Molecular structure of the title compound, showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. Molecular packing in the title compound. Hydrogen bonds are shown as dashed lines.

4-Chloro-*N*-(2-chlorobenzoyl)benzenesulfonamide

Crystal data

$C_{13}H_9Cl_2NO_3S$

$M_r = 330.17$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 6.3882$ (9) Å

$b = 10.311$ (1) Å

$c = 11.171$ (1) Å

$\alpha = 79.01$ (1)°

$\beta = 74.47$ (1)°

$\gamma = 84.76$ (1)°

$V = 695.32$ (13) Å³

$Z = 2$

$F(000) = 336$

$D_x = 1.577$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2339 reflections

$\theta = 2.5$ – 27.8 °

$\mu = 0.62$ mm⁻¹

$T = 299$ K

Prism, colourless

$0.24 \times 0.20 \times 0.14$ mm

Data collection

Oxford Diffraction Xcalibur
diffractometer with a Sapphire CCD detector

Radiation source: fine-focus sealed tube
graphite

ω and ϕ scans

Absorption correction: multi-scan

2825 independent reflections

2378 reflections with $I > 2\sigma(I)$

$R_{int} = 0.016$

$\theta_{max} = 26.4$ °, $\theta_{min} = 2.5$ °

$h = -7 \rightarrow 7$

(CrysAlis RED; Oxford Diffraction, 2009)

$T_{\min} = 0.865$, $T_{\max} = 0.918$

$k = -12 \rightarrow 11$

4417 measured reflections

$l = -13 \rightarrow 13$

Refinement

Refinement on F^2

Primary atom site location: structure-invariant direct methods

Least-squares matrix: full

Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.039$

Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.107$

H atoms treated by a mixture of independent and constrained refinement

$S = 1.06$

$w = 1/[\sigma^2(F_o^2) + (0.0603P)^2 + 0.1863P]$

where $P = (F_o^2 + 2F_c^2)/3$

2825 reflections

$(\Delta/\sigma)_{\max} = 0.022$

184 parameters

$\Delta\rho_{\max} = 0.25 \text{ e } \text{\AA}^{-3}$

1 restraint

$\Delta\rho_{\min} = -0.52 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.3004 (3)	0.27841 (17)	0.07988 (17)	0.0364 (4)
C2	0.1714 (3)	0.39202 (19)	0.0968 (2)	0.0464 (5)
H2	0.0437	0.4059	0.0703	0.056*
C3	0.2338 (4)	0.4844 (2)	0.1532 (2)	0.0561 (6)
H3	0.1510	0.5624	0.1634	0.067*
C4	0.4198 (4)	0.4596 (2)	0.1942 (2)	0.0520 (5)
C5	0.5482 (4)	0.3465 (2)	0.1791 (2)	0.0543 (5)
H5	0.6730	0.3317	0.2085	0.065*
C6	0.4895 (3)	0.2553 (2)	0.1198 (2)	0.0478 (5)
H6	0.5760	0.1791	0.1068	0.057*
C7	0.0089 (3)	0.02215 (18)	0.22952 (18)	0.0384 (4)
C8	0.0163 (3)	-0.09961 (18)	0.32606 (17)	0.0396 (4)
C9	0.1632 (3)	-0.1155 (2)	0.39978 (18)	0.0451 (4)
C10	0.1579 (5)	-0.2246 (3)	0.4941 (2)	0.0641 (7)

supplementary materials

H10	0.2565	-0.2347	0.5436	0.077*
C11	0.0073 (6)	-0.3171 (3)	0.5141 (2)	0.0757 (8)
H11	0.0043	-0.3908	0.5771	0.091*
C12	-0.1402 (5)	-0.3025 (3)	0.4422 (3)	0.0730 (7)
H12	-0.2424	-0.3663	0.4568	0.088*
C13	-0.1370 (4)	-0.1940 (2)	0.3487 (2)	0.0554 (5)
H13	-0.2378	-0.1840	0.3007	0.066*
Cl1	0.34983 (10)	0.00381 (7)	0.37663 (6)	0.0664 (2)
Cl2	0.49683 (16)	0.57327 (7)	0.26873 (8)	0.0938 (3)
N1	0.1748 (3)	0.02815 (15)	0.12059 (15)	0.0398 (4)
H1N	0.276 (3)	-0.0255 (19)	0.114 (2)	0.048*
O1	0.0294 (3)	0.20637 (15)	-0.02823 (14)	0.0550 (4)
O2	0.4074 (3)	0.11865 (14)	-0.08329 (13)	0.0537 (4)
O3	-0.1307 (2)	0.10807 (16)	0.24443 (16)	0.0602 (4)
S1	0.22277 (8)	0.15972 (4)	0.00846 (4)	0.04047 (15)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0364 (9)	0.0325 (8)	0.0360 (9)	-0.0001 (7)	-0.0058 (7)	-0.0008 (7)
C2	0.0448 (11)	0.0390 (10)	0.0503 (11)	0.0069 (8)	-0.0122 (9)	0.0006 (8)
C3	0.0695 (14)	0.0344 (10)	0.0568 (13)	0.0062 (9)	-0.0075 (11)	-0.0055 (9)
C4	0.0665 (14)	0.0424 (11)	0.0432 (11)	-0.0177 (9)	-0.0044 (10)	-0.0045 (8)
C5	0.0468 (11)	0.0561 (13)	0.0610 (13)	-0.0108 (9)	-0.0159 (10)	-0.0051 (10)
C6	0.0382 (10)	0.0431 (10)	0.0617 (13)	0.0040 (8)	-0.0140 (9)	-0.0091 (9)
C7	0.0365 (9)	0.0413 (9)	0.0406 (9)	-0.0023 (7)	-0.0146 (7)	-0.0076 (7)
C8	0.0430 (10)	0.0399 (9)	0.0342 (9)	0.0004 (7)	-0.0076 (7)	-0.0064 (7)
C9	0.0464 (10)	0.0529 (11)	0.0364 (10)	0.0068 (8)	-0.0111 (8)	-0.0120 (8)
C10	0.0786 (16)	0.0707 (15)	0.0399 (11)	0.0199 (13)	-0.0188 (11)	-0.0076 (10)
C11	0.112 (2)	0.0556 (14)	0.0457 (13)	0.0047 (14)	-0.0087 (14)	0.0059 (11)
C12	0.095 (2)	0.0539 (14)	0.0628 (16)	-0.0230 (13)	-0.0095 (14)	0.0015 (12)
C13	0.0615 (13)	0.0548 (12)	0.0500 (12)	-0.0148 (10)	-0.0127 (10)	-0.0059 (10)
Cl1	0.0571 (4)	0.0895 (4)	0.0633 (4)	-0.0126 (3)	-0.0273 (3)	-0.0178 (3)
Cl2	0.1363 (7)	0.0706 (5)	0.0834 (5)	-0.0386 (4)	-0.0236 (5)	-0.0259 (4)
N1	0.0441 (9)	0.0340 (8)	0.0394 (8)	0.0013 (6)	-0.0095 (7)	-0.0049 (6)
O1	0.0577 (9)	0.0586 (9)	0.0546 (9)	0.0005 (7)	-0.0303 (7)	-0.0030 (7)
O2	0.0656 (9)	0.0503 (8)	0.0367 (7)	0.0079 (7)	-0.0037 (6)	-0.0050 (6)
O3	0.0483 (8)	0.0617 (9)	0.0603 (10)	0.0152 (7)	-0.0092 (7)	-0.0008 (7)
S1	0.0464 (3)	0.0387 (3)	0.0356 (3)	0.00192 (18)	-0.01273 (19)	-0.00320 (18)

Geometric parameters (\AA , $^\circ$)

C1—C6	1.382 (3)	C8—C13	1.385 (3)
C1—C2	1.383 (3)	C8—C9	1.386 (3)
C1—S1	1.7537 (19)	C9—C10	1.382 (3)
C2—C3	1.377 (3)	C9—C11	1.727 (2)
C2—H2	0.9300	C10—C11	1.363 (4)
C3—C4	1.370 (3)	C10—H10	0.9300
C3—H3	0.9300	C11—C12	1.375 (4)

C4—C5	1.374 (3)	C11—H11	0.9300
C4—C12	1.733 (2)	C12—C13	1.375 (3)
C5—C6	1.379 (3)	C12—H12	0.9300
C5—H5	0.9300	C13—H13	0.9300
C6—H6	0.9300	N1—S1	1.6510 (16)
C7—O3	1.200 (2)	N1—H1N	0.810 (15)
C7—N1	1.379 (3)	O1—S1	1.4185 (15)
C7—C8	1.497 (3)	O2—S1	1.4311 (15)
C6—C1—C2	121.20 (18)	C10—C9—C8	120.5 (2)
C6—C1—S1	118.92 (14)	C10—C9—C11	119.64 (18)
C2—C1—S1	119.87 (15)	C8—C9—C11	119.82 (16)
C3—C2—C1	119.35 (19)	C11—C10—C9	119.6 (2)
C3—C2—H2	120.3	C11—C10—H10	120.2
C1—C2—H2	120.3	C9—C10—H10	120.2
C4—C3—C2	119.03 (19)	C10—C11—C12	120.7 (2)
C4—C3—H3	120.5	C10—C11—H11	119.7
C2—C3—H3	120.5	C12—C11—H11	119.7
C3—C4—C5	122.2 (2)	C13—C12—C11	120.2 (2)
C3—C4—C12	119.51 (17)	C13—C12—H12	119.9
C5—C4—C12	118.31 (18)	C11—C12—H12	119.9
C4—C5—C6	119.0 (2)	C12—C13—C8	120.0 (2)
C4—C5—H5	120.5	C12—C13—H13	120.0
C6—C5—H5	120.5	C8—C13—H13	120.0
C5—C6—C1	119.17 (18)	C7—N1—S1	124.25 (13)
C5—C6—H6	120.4	C7—N1—H1N	121.9 (16)
C1—C6—H6	120.4	S1—N1—H1N	112.1 (16)
O3—C7—N1	122.05 (18)	O1—S1—O2	119.26 (9)
O3—C7—C8	123.12 (18)	O1—S1—N1	110.18 (9)
N1—C7—C8	114.82 (15)	O2—S1—N1	103.74 (8)
C13—C8—C9	119.07 (19)	O1—S1—C1	108.96 (9)
C13—C8—C7	119.35 (17)	O2—S1—C1	109.21 (9)
C9—C8—C7	121.40 (17)	N1—S1—C1	104.43 (8)
C6—C1—C2—C3	0.7 (3)	C8—C9—C10—C11	0.2 (3)
S1—C1—C2—C3	179.87 (16)	C11—C9—C10—C11	178.50 (19)
C1—C2—C3—C4	-1.7 (3)	C9—C10—C11—C12	-0.5 (4)
C2—C3—C4—C5	1.0 (3)	C10—C11—C12—C13	0.1 (4)
C2—C3—C4—C12	-178.50 (16)	C11—C12—C13—C8	0.6 (4)
C3—C4—C5—C6	0.6 (3)	C9—C8—C13—C12	-0.9 (3)
C12—C4—C5—C6	-179.85 (17)	C7—C8—C13—C12	-176.2 (2)
C4—C5—C6—C1	-1.6 (3)	O3—C7—N1—S1	9.1 (3)
C2—C1—C6—C5	0.9 (3)	C8—C7—N1—S1	-171.37 (13)
S1—C1—C6—C5	-178.23 (16)	C7—N1—S1—O1	-51.19 (18)
O3—C7—C8—C13	70.4 (3)	C7—N1—S1—O2	-179.94 (15)
N1—C7—C8—C13	-109.1 (2)	C7—N1—S1—C1	65.68 (17)
O3—C7—C8—C9	-104.8 (2)	C6—C1—S1—O1	-178.06 (15)
N1—C7—C8—C9	75.7 (2)	C2—C1—S1—O1	2.78 (18)
C13—C8—C9—C10	0.4 (3)	C6—C1—S1—O2	-46.22 (18)
C7—C8—C9—C10	175.66 (18)	C2—C1—S1—O2	134.61 (15)

supplementary materials

C13—C8—C9—C11	-177.81 (16)	C6—C1—S1—N1	64.23 (17)
C7—C8—C9—C11	-2.6 (3)	C2—C1—S1—N1	-114.93 (16)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1N \cdots O2 ⁱ	0.81 (2)	2.13 (2)	2.914 (2)	164 (2)

Symmetry codes: (i) $-x+1, -y, -z$.

Fig. 1




Fig. 2

