

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

[5,10,15,20-Tetrakis(4-tolyl)porphyrin]-zinc(II) dichloromethane solvateSean McGill,^a Vladimir N. Nesterov^b and Stephanie L. Gould^{a*}^aDepartment of Chemistry, Austin College, 900 North Grand, Sherman, TX 75090-4400, USA, and ^bDepartment of Chemistry, University of North Texas, Denton, TX 76203-5017, USA

Correspondence e-mail: sgould@austincollege.edu


Received 25 April 2010; accepted 26 May 2010

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; disorder in solvent or counterion; R factor = 0.042; wR factor = 0.094; data-to-parameter ratio = 14.9.

In the title complex, $[\text{Zn}(\text{C}_{48}\text{H}_{36}\text{N}_4)] \cdot \text{CH}_2\text{Cl}_2$, the Zn^{II} atom lies on an inversion center and the dichloromethane solvent molecule is disordered around an inversion center. The tolyl substituents are twisted compared to the central aromatic ring system of the porphyrin, similar to what is seen in previously published structures of this molecule [Dastidar & Goldberg (1996). *Acta Cryst.* **C52**, 1976–1980]. The dihedral angles between the mean planes of the tolyl rings and the central ring are 66.98 (6)° and 60.40 (6)°.

Related literature

For other solvates of this molecule see: Dastidar & Goldberg (1996). For similar structures of ligand-bridged porphyrin sandwich-type structures, see: Diskin-Posner *et al.* (2002); Mak *et al.* (1998); Kieran *et al.* (2005); Dastidar *et al.* (1996). For the synthesis of the title compound, see: Adler *et al.* (1967).



Experimental

Crystal data

$[\text{Zn}(\text{C}_{48}\text{H}_{36}\text{N}_4)] \cdot \text{CH}_2\text{Cl}_2$
 $M_r = 819.10$
 Monoclinic, $P2_1/c$
 $a = 14.349$ (2) Å
 $b = 8.5273$ (14) Å
 $c = 15.637$ (3) Å
 $\beta = 94.995$ (2)°

$V = 1906.1$ (5) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.83$ mm⁻¹
 $T = 100$ K
 $0.16 \times 0.13 \times 0.09$ mm

Data collection

Bruker SMART APEXII CCD diffractometer
 Absorption correction: numerical (*SADABS*; Bruker, 2001)
 $T_{\text{min}} = 0.880$, $T_{\text{max}} = 0.930$

16249 measured reflections
 3901 independent reflections
 3533 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.094$
 $S = 1.00$
 3901 reflections
 261 parameters

1 restraint
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.95$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.23$ e Å⁻³

Data collection: *APEX2* (Bruker, 2007); cell refinement: *S SAINT* (Bruker, 2007); data reduction: *S SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

This research was funded by in part by the National Science Foundation (CHE-0924153) and a chemistry department grant from the Welch Foundation (AD-0007). The X-ray data were collected at the University of North Texas.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NK2033).

References

- Adler, A. D., Longo, F. R., Finarelli, J. D., Goldmacher, J., Assour, J. & Korsakoff, L. (1967). *J. Org. Chem.* **32**, 476–477.
 Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2007). *APEX2* and *S SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Dastidar, P. & Goldberg, I. (1996). *Acta Cryst.* **C52**, 1976–1980.
 Dastidar, P., Krupitsky, H., Stein, Z. & Goldberg, I. (1996). *J. Incl. Phen. Mol. Recogn. Chem.* **24**, 241–262.
 Diskin-Posner, Y., Patra, G. K. & Goldberg, I. (2002). *CrystEngComm*, **4**, 296–301.
 Kieran, A. L., Pascu, S. I., Jarrosson, T. & Sanders, J. K. M. (2005). *Chem. Commun.* pp. 1276–1279.
 Mak, C. C., Bampos, N. & Sanders, J. K. M. (1998). *Angew. Chem. Int. Ed.* **37**, 3020–3023.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2010). E66, m723 [doi:10.1107/S1600536810019963]

[5,10,15,20-Tetrakis(4-tolyl)porphyrin]zinc(II) dichloromethane solvate

S. McGill, V. N. Nesterov and S. L. Gould

Comment

While pursuing ligand bridged porphyrin sandwich-type supramolecular structures, similar to those created by Diskin-Posner *et al.* (2002), Mak *et al.* (1998) and Kieran *et al.* (2005), we sought to crystallize zinc 5,10,15,20-tetrakis(4'-tolyl)porphyrin with pyrazine. The resulting deep red crystals that formed were found not to contain any pyrazine, but contained a well ordered porphyrin structure different than those previously published (Dastidar & Goldberg, 1996). The role of the pyrazine in the crystal deposition is unknown and will be explored further as the same crystals can not be obtained without the compound being present.

The title porphyrin (Figure 1) has a zinc atom located at the center of the porphyrin framework in near exact planarity to the porphyrin. The tolyl substituents are angled compared to the central aromatic ring of the porphyrin, similar to what is seen in previously published structures (Dastidar & Goldberg, 1996). The dihedral angles between the mean planes of the peripheral rings to the central ring are 66.98 (6) and 60.40 (6) °.

Experimental

The synthesis of zinc 5,10,15,20-tetrakis(4'-tolyl)porphyrin was carried out according literature procedures (Adler *et al.*, 1967). Dark red crystals were grown by liquid diffusion of methanol into a dichloromethane solution containing 20 mg of zinc 5,10,15,20-tetrakis(4'- tolyl)porphyrin and 2 mg of pyrazine.

Refinement

The dichloromethane solvent molecule is disordered around a center of inversion. Non-hydrogen atoms were refined anisotropically with an occupation factor of 0.5 for C and 1.0 for Cl. All C-bound H atoms were placed in idealized positions (C—H = 0.95–1.00 Å) and allowed to ride on their parent atoms. H atoms were constrained so that $U_{\text{iso}}(\text{H})$ were equal to 1.2U_{eq} or 1.5U_{eq} of their respective parent atoms.

Figures




Fig. 1. Ellipsoid plot of zinc 5,10,15,20-tetrakis(4'-tolyl)porphyrin (50% probability displacement ellipsoids). Unlabeled atoms are related to labeled atoms by inversion symmetry.

[5,10,15,20-Tetrakis(4-tolyl)porphyrin]zinc(II) dichloromethane solvate

Crystal data

$[\text{Zn}(\text{C}_{48}\text{H}_{36}\text{N}_4)] \cdot \text{CH}_2\text{Cl}_2$	$F(000) = 848$
$M_r = 819.10$	$D_x = 1.427 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 8831 reflections
$a = 14.349 (2) \text{ \AA}$	$\theta = 2.6\text{--}26.1^\circ$
$b = 8.5273 (14) \text{ \AA}$	$\mu = 0.83 \text{ mm}^{-1}$
$c = 15.637 (3) \text{ \AA}$	$T = 100 \text{ K}$
$\beta = 94.995 (2)^\circ$	Block, red
$V = 1906.1 (5) \text{ \AA}^3$	$0.16 \times 0.13 \times 0.09 \text{ mm}$
$Z = 2$	

Data collection

Bruker SMART APEXII CCD diffractometer	3901 independent reflections
Radiation source: fine-focus sealed tube graphite	3533 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.026$
Absorption correction: numerical (<i>SADABS</i> ; Bruker, 2001)	$\theta_{\text{max}} = 26.4^\circ$, $\theta_{\text{min}} = 2.6^\circ$
$T_{\text{min}} = 0.880$, $T_{\text{max}} = 0.930$	$h = -17 \rightarrow 17$
16249 measured reflections	$k = -10 \rightarrow 10$
	$l = -19 \rightarrow 19$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.042$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.094$	H-atom parameters constrained
$S = 1.00$	$w = 1/[\sigma^2(F_o^2) + (0.020P)^2 + 5.P]$
3901 reflections	where $P = (F_o^2 + 2F_c^2)/3$
261 parameters	$(\Delta/\sigma)_{\text{max}} = 0.002$
1 restraint	$\Delta\rho_{\text{max}} = 0.95 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -1.23 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations

between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Zn1	0.5000	0.5000	0.5000	0.01504 (11)	
N1	0.43558 (13)	0.5556 (2)	0.38246 (11)	0.0164 (4)	
N2	0.37660 (13)	0.5259 (2)	0.55317 (11)	0.0164 (4)	
C1	0.27573 (15)	0.6038 (3)	0.42370 (14)	0.0168 (4)	
C2	0.34261 (16)	0.5909 (3)	0.36365 (14)	0.0170 (4)	
C3	0.32403 (16)	0.6157 (3)	0.27249 (14)	0.0193 (5)	
H3A	0.2649	0.6380	0.2428	0.023*	
C4	0.40637 (16)	0.6012 (3)	0.23723 (14)	0.0187 (5)	
H4A	0.4163	0.6145	0.1784	0.022*	
C5	0.47622 (16)	0.5618 (3)	0.30559 (14)	0.0167 (4)	
C6	0.57093 (15)	0.5358 (3)	0.29533 (14)	0.0163 (4)	
C7	0.60335 (15)	0.5564 (3)	0.20727 (14)	0.0165 (4)	
C8	0.57431 (16)	0.4550 (3)	0.14006 (14)	0.0182 (5)	
H8A	0.5322	0.3721	0.1496	0.022*	
C9	0.60651 (16)	0.4740 (3)	0.05923 (14)	0.0197 (5)	
H9A	0.5861	0.4036	0.0143	0.024*	
C10	0.66818 (16)	0.5946 (3)	0.04310 (14)	0.0208 (5)	
C11	0.69649 (17)	0.6956 (3)	0.11008 (15)	0.0215 (5)	
H11A	0.7385	0.7785	0.1005	0.026*	
C12	0.66440 (16)	0.6775 (3)	0.19122 (14)	0.0194 (5)	
H12A	0.6844	0.7485	0.2359	0.023*	
C13	0.7035 (2)	0.6148 (3)	-0.04444 (15)	0.0301 (6)	
H13A	0.7502	0.6989	-0.0422	0.045*	
H13B	0.6511	0.6420	-0.0863	0.045*	
H13C	0.7321	0.5167	-0.0617	0.045*	
C14	0.29318 (16)	0.5754 (3)	0.51224 (14)	0.0171 (4)	
C15	0.22477 (16)	0.5933 (3)	0.57396 (14)	0.0189 (5)	
H15A	0.1622	0.6288	0.5625	0.023*	
C16	0.26662 (16)	0.5501 (3)	0.65116 (14)	0.0191 (5)	
H16A	0.2384	0.5474	0.7039	0.023*	
C17	0.36176 (15)	0.5088 (3)	0.63887 (14)	0.0167 (4)	
C18	0.17871 (15)	0.6502 (3)	0.39095 (14)	0.0181 (5)	
C19	0.16187 (16)	0.7945 (3)	0.35042 (14)	0.0207 (5)	
H19A	0.2122	0.8654	0.3455	0.025*	
C20	0.07234 (17)	0.8357 (3)	0.31716 (15)	0.0246 (5)	
H20A	0.0626	0.9335	0.2887	0.030*	
C21	-0.00341 (17)	0.7362 (3)	0.32472 (15)	0.0247 (5)	
C22	0.01316 (17)	0.5947 (3)	0.36693 (15)	0.0241 (5)	

supplementary materials

H22A	-0.0378	0.5260	0.3739	0.029*	
C23	0.10252 (16)	0.5513 (3)	0.39923 (15)	0.0208 (5)	
H23A	0.1120	0.4531	0.4273	0.025*	
C24	-0.10064 (18)	0.7807 (4)	0.28820 (18)	0.0340 (6)	
H24A	-0.1464	0.7423	0.3262	0.051*	
H24B	-0.1133	0.7336	0.2312	0.051*	
H24C	-0.1053	0.8951	0.2835	0.051*	
C11	-0.00291 (18)	1.13537 (18)	0.44337 (11)	0.1434 (9)	
C1A	0.0459 (5)	0.9997 (10)	0.5120 (5)	0.081 (3)	0.50
H1AA	0.0352	1.0697	0.5615	0.098*	0.50
H1AB	0.0987	0.9580	0.4859	0.098*	0.50

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.01529 (18)	0.01885 (19)	0.01128 (17)	0.00098 (14)	0.00281 (13)	0.00119 (14)
N1	0.0164 (9)	0.0193 (9)	0.0138 (9)	0.0001 (7)	0.0031 (7)	0.0008 (7)
N2	0.0166 (9)	0.0203 (10)	0.0125 (9)	0.0004 (7)	0.0022 (7)	0.0014 (7)
C1	0.0173 (11)	0.0175 (11)	0.0157 (10)	-0.0008 (9)	0.0020 (8)	0.0009 (9)
C2	0.0181 (11)	0.0176 (11)	0.0154 (10)	-0.0011 (9)	0.0014 (8)	0.0007 (8)
C3	0.0199 (11)	0.0226 (12)	0.0150 (11)	0.0003 (9)	-0.0001 (9)	0.0007 (9)
C4	0.0213 (11)	0.0217 (12)	0.0132 (10)	0.0009 (9)	0.0017 (8)	0.0010 (9)
C5	0.0203 (11)	0.0165 (11)	0.0136 (10)	-0.0008 (9)	0.0026 (8)	0.0001 (8)
C6	0.0192 (11)	0.0159 (11)	0.0139 (10)	-0.0013 (8)	0.0033 (8)	-0.0006 (8)
C7	0.0166 (10)	0.0198 (11)	0.0133 (10)	0.0038 (9)	0.0029 (8)	0.0020 (8)
C8	0.0173 (11)	0.0202 (11)	0.0170 (11)	0.0007 (9)	0.0007 (8)	0.0011 (9)
C9	0.0211 (11)	0.0234 (12)	0.0144 (10)	0.0034 (9)	0.0011 (8)	-0.0024 (9)
C10	0.0225 (12)	0.0255 (12)	0.0150 (11)	0.0058 (10)	0.0046 (9)	0.0034 (9)
C11	0.0235 (12)	0.0219 (12)	0.0200 (11)	-0.0018 (10)	0.0064 (9)	0.0034 (9)
C12	0.0218 (11)	0.0211 (12)	0.0156 (11)	-0.0004 (9)	0.0029 (9)	-0.0014 (9)
C13	0.0384 (15)	0.0363 (15)	0.0172 (12)	-0.0014 (12)	0.0106 (11)	0.0014 (11)
C14	0.0176 (11)	0.0179 (11)	0.0162 (11)	-0.0008 (9)	0.0034 (8)	0.0001 (9)
C15	0.0165 (11)	0.0232 (12)	0.0175 (11)	0.0014 (9)	0.0041 (8)	0.0006 (9)
C16	0.0198 (11)	0.0230 (12)	0.0151 (10)	0.0000 (9)	0.0048 (8)	-0.0008 (9)
C17	0.0197 (11)	0.0172 (11)	0.0137 (10)	-0.0007 (9)	0.0044 (8)	-0.0009 (8)
C18	0.0173 (11)	0.0252 (12)	0.0120 (10)	0.0014 (9)	0.0021 (8)	-0.0011 (9)
C19	0.0192 (11)	0.0253 (12)	0.0177 (11)	-0.0014 (9)	0.0027 (9)	0.0022 (9)
C20	0.0241 (12)	0.0301 (13)	0.0195 (11)	0.0039 (10)	0.0020 (9)	0.0051 (10)
C21	0.0183 (11)	0.0389 (15)	0.0170 (11)	0.0026 (10)	0.0018 (9)	0.0002 (10)
C22	0.0192 (11)	0.0347 (14)	0.0186 (11)	-0.0056 (10)	0.0030 (9)	-0.0002 (10)
C23	0.0208 (11)	0.0237 (12)	0.0182 (11)	-0.0015 (10)	0.0033 (9)	0.0009 (9)
C24	0.0191 (13)	0.0513 (18)	0.0311 (14)	0.0039 (12)	-0.0009 (10)	0.0072 (13)
C11	0.272 (3)	0.0665 (9)	0.1016 (11)	0.0678 (12)	0.0716 (14)	-0.0032 (8)
C1A	0.038 (4)	0.125 (8)	0.080 (6)	0.025 (5)	-0.005 (4)	-0.074 (6)

Geometric parameters (\AA , $^\circ$)

Zn1—N2	2.0326 (18)	C13—H13A	0.9800
--------	-------------	----------	--------

Zn1—N2 ⁱ	2.0327 (18)	C13—H13B	0.9800
Zn1—N1	2.0401 (18)	C13—H13C	0.9800
Zn1—N1 ⁱ	2.0402 (18)	C14—C15	1.443 (3)
N1—C2	1.374 (3)	C15—C16	1.352 (3)
N1—C5	1.382 (3)	C15—H15A	0.9500
N2—C14	1.374 (3)	C16—C17	1.439 (3)
N2—C17	1.383 (3)	C16—H16A	0.9500
C1—C2	1.404 (3)	C17—C6 ⁱ	1.401 (3)
C1—C14	1.406 (3)	C18—C23	1.396 (3)
C1—C18	1.494 (3)	C18—C19	1.395 (3)
C2—C3	1.443 (3)	C19—C20	1.389 (3)
C3—C4	1.352 (3)	C19—H19A	0.9500
C3—H3A	0.9500	C20—C21	1.392 (4)
C4—C5	1.440 (3)	C20—H20A	0.9500
C4—H4A	0.9500	C21—C22	1.386 (4)
C5—C6	1.400 (3)	C21—C24	1.509 (3)
C6—C17 ⁱ	1.401 (3)	C22—C23	1.387 (3)
C6—C7	1.502 (3)	C22—H22A	0.9500
C7—C12	1.391 (3)	C23—H23A	0.9500
C7—C8	1.396 (3)	C24—H24A	0.9800
C8—C9	1.392 (3)	C24—H24B	0.9800
C8—H8A	0.9500	C24—H24C	0.9800
C9—C10	1.394 (3)	C11—C1A ⁱⁱ	1.507 (9)
C9—H9A	0.9500	C11—C1A	1.688 (8)
C10—C11	1.389 (3)	C1A—C1A ⁱⁱ	1.339 (14)
C10—C13	1.510 (3)	C1A—C11 ⁱⁱ	1.507 (9)
C11—C12	1.396 (3)	C1A—H1AA	1.0000
C11—H11A	0.9500	C1A—H1AB	0.9592
C12—H12A	0.9500		
N2—Zn1—N2 ⁱ	180.00 (9)	C10—C13—H13C	109.5
N2—Zn1—N1	90.04 (7)	H13A—C13—H13C	109.5
N2 ⁱ —Zn1—N1	89.96 (7)	H13B—C13—H13C	109.5
N2—Zn1—N1 ⁱ	89.96 (7)	N2—C14—C1	125.8 (2)
N2 ⁱ —Zn1—N1 ⁱ	90.04 (7)	N2—C14—C15	109.68 (19)
N1—Zn1—N1 ⁱ	179.999 (1)	C1—C14—C15	124.5 (2)
C2—N1—C5	106.32 (18)	C16—C15—C14	107.1 (2)
C2—N1—Zn1	126.80 (14)	C16—C15—H15A	126.5
C5—N1—Zn1	126.87 (15)	C14—C15—H15A	126.5
C14—N2—C17	106.40 (18)	C15—C16—C17	107.5 (2)
C14—N2—Zn1	126.70 (14)	C15—C16—H16A	126.3
C17—N2—Zn1	126.74 (15)	C17—C16—H16A	126.3
C2—C1—C14	124.9 (2)	N2—C17—C6 ⁱ	125.9 (2)
C2—C1—C18	117.52 (19)	N2—C17—C16	109.37 (19)
C14—C1—C18	117.63 (19)	C6 ⁱ —C17—C16	124.7 (2)
N1—C2—C1	125.5 (2)	C23—C18—C19	118.0 (2)
N1—C2—C3	109.62 (19)	C23—C18—C1	121.5 (2)

supplementary materials

C1—C2—C3	124.8 (2)	C19—C18—C1	120.6 (2)
C4—C3—C2	107.3 (2)	C20—C19—C18	120.7 (2)
C4—C3—H3A	126.4	C20—C19—H19A	119.6
C2—C3—H3A	126.4	C18—C19—H19A	119.6
C3—C4—C5	107.2 (2)	C19—C20—C21	121.2 (2)
C3—C4—H4A	126.4	C19—C20—H20A	119.4
C5—C4—H4A	126.4	C21—C20—H20A	119.4
N1—C5—C6	125.5 (2)	C22—C21—C20	117.9 (2)
N1—C5—C4	109.59 (19)	C22—C21—C24	120.9 (2)
C6—C5—C4	124.9 (2)	C20—C21—C24	121.2 (2)
C5—C6—C17 ⁱ	124.9 (2)	C21—C22—C23	121.4 (2)
C5—C6—C7	117.89 (19)	C21—C22—H22A	119.3
C17 ⁱ —C6—C7	117.16 (19)	C23—C22—H22A	119.3
C12—C7—C8	118.4 (2)	C22—C23—C18	120.8 (2)
C12—C7—C6	120.2 (2)	C22—C23—H23A	119.6
C8—C7—C6	121.4 (2)	C18—C23—H23A	119.6
C9—C8—C7	120.7 (2)	C21—C24—H24A	109.5
C9—C8—H8A	119.7	C21—C24—H24B	109.5
C7—C8—H8A	119.7	H24A—C24—H24B	109.5
C8—C9—C10	121.1 (2)	C21—C24—H24C	109.5
C8—C9—H9A	119.5	H24A—C24—H24C	109.5
C10—C9—H9A	119.5	H24B—C24—H24C	109.5
C11—C10—C9	118.0 (2)	C1A ⁱⁱ —C11—C1A	49.1 (5)
C11—C10—C13	120.9 (2)	C1A ⁱⁱ —C1A—C11 ⁱⁱ	72.5 (8)
C9—C10—C13	121.1 (2)	C1A ⁱⁱ —C1A—C11	58.4 (6)
C10—C11—C12	121.3 (2)	C11 ⁱⁱ —C1A—C11	130.9 (5)
C10—C11—H11A	119.4	C1A ⁱⁱ —C1A—H1AA	90.0
C12—C11—H11A	119.4	C11 ⁱⁱ —C1A—H1AA	90.0
C7—C12—C11	120.6 (2)	C11—C1A—H1AA	90.0
C7—C12—H12A	119.7	C1A ⁱⁱ —C1A—H1AB	132.6
C11—C12—H12A	119.7	C11 ⁱⁱ —C1A—H1AB	106.5
C10—C13—H13A	109.5	C11—C1A—H1AB	106.4
C10—C13—H13B	109.5	H1AA—C1A—H1AB	137.0
H13A—C13—H13B	109.5		
N2—Zn1—N1—C2	-0.55 (19)	C9—C10—C11—C12	-0.1 (4)
N2 ⁱ —Zn1—N1—C2	179.45 (19)	C13—C10—C11—C12	-179.8 (2)
N2—Zn1—N1—C5	-178.92 (19)	C8—C7—C12—C11	-0.7 (3)
N2 ⁱ —Zn1—N1—C5	1.08 (19)	C6—C7—C12—C11	178.6 (2)
N1—Zn1—N2—C14	-4.16 (19)	C10—C11—C12—C7	0.5 (4)
N1 ⁱ —Zn1—N2—C14	175.84 (19)	C17—N2—C14—C1	-178.3 (2)
N1—Zn1—N2—C17	-178.82 (19)	Zn1—N2—C14—C1	6.2 (3)
N1 ⁱ —Zn1—N2—C17	1.18 (19)	C17—N2—C14—C15	1.3 (3)
C5—N1—C2—C1	-177.5 (2)	Zn1—N2—C14—C15	-174.23 (15)
Zn1—N1—C2—C1	3.9 (3)	C2—C1—C14—N2	-2.4 (4)
C5—N1—C2—C3	1.6 (3)	C18—C1—C14—N2	177.3 (2)
Zn1—N1—C2—C3	-177.08 (15)	C2—C1—C14—C15	178.1 (2)

C14—C1—C2—N1	-3.0 (4)	C18—C1—C14—C15	-2.2 (3)
C18—C1—C2—N1	177.3 (2)	N2—C14—C15—C16	-1.8 (3)
C14—C1—C2—C3	178.1 (2)	C1—C14—C15—C16	177.8 (2)
C18—C1—C2—C3	-1.5 (3)	C14—C15—C16—C17	1.4 (3)
N1—C2—C3—C4	-2.3 (3)	C14—N2—C17—C6 ⁱ	-178.2 (2)
C1—C2—C3—C4	176.7 (2)	Zn1—N2—C17—C6 ⁱ	-2.6 (3)
C2—C3—C4—C5	2.0 (3)	C14—N2—C17—C16	-0.4 (3)
C2—N1—C5—C6	179.0 (2)	Zn1—N2—C17—C16	175.12 (15)
Zn1—N1—C5—C6	-2.4 (3)	C15—C16—C17—N2	-0.7 (3)
C2—N1—C5—C4	-0.3 (3)	C15—C16—C17—C6 ⁱ	177.1 (2)
Zn1—N1—C5—C4	178.34 (15)	C2—C1—C18—C23	118.4 (2)
C3—C4—C5—N1	-1.1 (3)	C14—C1—C18—C23	-61.2 (3)
C3—C4—C5—C6	179.6 (2)	C2—C1—C18—C19	-61.1 (3)
N1—C5—C6—C17 ⁱ	3.6 (4)	C14—C1—C18—C19	119.2 (2)
C4—C5—C6—C17 ⁱ	-177.3 (2)	C23—C18—C19—C20	-1.9 (3)
N1—C5—C6—C7	-176.6 (2)	C1—C18—C19—C20	177.7 (2)
C4—C5—C6—C7	2.5 (3)	C18—C19—C20—C21	1.3 (4)
C5—C6—C7—C12	113.2 (2)	C19—C20—C21—C22	0.3 (4)
C17 ⁱ —C6—C7—C12	-67.0 (3)	C19—C20—C21—C24	-179.7 (2)
C5—C6—C7—C8	-67.5 (3)	C20—C21—C22—C23	-1.3 (4)
C17 ⁱ —C6—C7—C8	112.3 (2)	C24—C21—C22—C23	178.7 (2)
C12—C7—C8—C9	0.6 (3)	C21—C22—C23—C18	0.7 (4)
C6—C7—C8—C9	-178.7 (2)	C19—C18—C23—C22	0.9 (3)
C7—C8—C9—C10	-0.2 (3)	C1—C18—C23—C22	-178.7 (2)
C8—C9—C10—C11	-0.1 (3)	C1A ⁱⁱ —C11—C1A—C11 ⁱⁱ	0.0
C8—C9—C10—C13	179.7 (2)		

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x, -y+2, -z+1$.

Fig. 1

