

catena-Poly[[bis(μ -3-aminopyrazine-2-carboxylato)- $\kappa^3 N^1, O^1; \kappa^3 O^1, O^1$]-dilithium]-di- μ -aqua]

Wojciech Starosta and Janusz Leciejewicz*

Institute of Nuclear Chemistry and Technology, ul. Dorodna 16, 03-195 Warszawa, Poland

Correspondence e-mail: j.leciejewicz@ichcj.waw.pl


Received 21 May 2010; accepted 31 May 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.049; wR factor = 0.147; data-to-parameter ratio = 16.6.

The title compound, $[\text{Li}(\text{C}_5\text{H}_4\text{N}_3\text{O}_2)(\text{H}_2\text{O})]_n$, is composed of centrosymmetric dinuclear units, in which the Li^{I} ions are bridged by two carboxylate O atoms donated by two ligands. The dinuclear unit is nearly planar [r.m.s. deviation = 0.0125 (2) \AA]. The Li^{I} ion is coordinated by an N,O -chelating ligand, a bridging carboxylate O atom from another ligand and two bridging water O atoms in a distorted trigonal-bipyramidal geometry. The water O atoms bridge the dinuclear units into a polymeric molecular column along [010]. The columns are held together by $\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds. An intramolecular $\text{N}-\text{H}\cdots\text{O}$ interaction also occurs.

Related literature

For the structures of metal (M) complexes with the 3-amino-pyrazine-2-carboxylate ligand, see: Leciejewicz *et al.* (1997 [$M = \text{Ca}(\text{II})$], 1998 [$M = \text{Sr}(\text{II})$]); Ptasiewicz-Bąk & Leciejewicz (1997 [$M = \text{Mg}(\text{II})$], 1999 [$M = \text{Ni}(\text{II})$]); Tayebee *et al.* (2008) [$M = \text{Na}(\text{I})$]. For the structure of an $\text{Li}(\text{I})$ complex with pyrazine-2,3-dicarboxylate and aqua ligands, see: Tombul *et al.* (2008).



Experimental

Crystal data

$[\text{Li}(\text{C}_5\text{H}_4\text{N}_3\text{O}_2)(\text{H}_2\text{O})]$
 $M_r = 163.07$
Monoclinic, $P2_{1}/c$
 $a = 14.279 (3)\text{ \AA}$
 $b = 3.6000 (7)\text{ \AA}$
 $c = 13.300 (3)\text{ \AA}$
 $\beta = 106.43 (3)^{\circ}$

$V = 655.7 (2)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.13\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.26 \times 0.21 \times 0.04\text{ mm}$

Data collection

Kuma KM-4 four-circle diffractometer
Absorption correction: analytical (*CrysAlis RED*; Oxford Diffraction, 2006)
 $T_{\min} = 0.980$, $T_{\max} = 0.994$
1997 measured reflections

1913 independent reflections
1297 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.017$
3 standard reflections every 200 reflections
intensity decay: 7.3%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.147$
 $S = 1.04$
1913 reflections
115 parameters
3 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.51\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.39\text{ e \AA}^{-3}$

Table 1
Selected bond lengths (\AA).

$\text{Li1}-\text{N1}$	2.118 (3)	$\text{Li1}-\text{O3}$	2.065 (3)
$\text{Li1}-\text{O1}$	1.999 (3)	$\text{Li1}-\text{O3}^{\text{ii}}$	2.201 (3)
$\text{Li1}-\text{O1}^{\text{i}}$	1.995 (3)		

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $x, y - 1, z$.

Table 2Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O3—H31···O2 ⁱⁱⁱ	0.88 (1)	1.83 (1)	2.7028 (16)	175 (2)
O3—H32···O1 ^{iv}	0.84 (2)	2.54 (2)	2.9083 (17)	108 (2)
N3—H1···O2	0.86	2.08	2.7229 (17)	131
N3—H2···N2 ^v	0.86	2.30	3.1278 (19)	162

Symmetry codes: (iii) $x, -y + \frac{3}{2}, z + \frac{1}{2}$; (iv) $-x + 1, -y + 2, -z + 1$; (v) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *KM-4 Software* (Kuma, 1996); cell refinement: *KM-4 Software*; data reduction: *DATAPROC* (Kuma, 2001); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2312).

References

- Kuma (1996). *KM-4 Software*. Kuma Diffraction Ltd, Wrocław, Poland.
- Kuma (2001). *DATAPROC*. Kuma Diffraction Ltd, Wrocław, Poland.
- Leciejewicz, J., Ptasiwicz-Bąk, H. & Paluchowska, B. (1997). *Pol. J. Chem.* **71**, 1339–1364.
- Leciejewicz, J., Ptasiwicz-Bąk, H. & Zachara, J. (1998). *Pol. J. Chem.* **72**, 1994–1998.
- Oxford Diffraction (2006). *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, England.
- Ptasiwicz-Bąk, H. & Leciejewicz, J. (1997). *Pol. J. Chem.* **71**, 1350–1358.
- Ptasiwicz-Bąk, H. & Leciejewicz, J. (1999). *Pol. J. Chem.* **73**, 717–725.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Tayebec, R., Amani, V. & Khavasi, H. P. (2008). *Chin. J. Chem.* **26**, 500–504.
- Tombul, M., Güven, K. & Büyükgüngör, O. (2008). *Acta Cryst. E* **64**, m491–m492.

supplementary materials

Acta Cryst. (2010). E66, m744-m745 [doi:10.1107/S1600536810020647]

catena-Poly[[bis(μ -3-aminopyrazine-2-carboxylato)- $\kappa^3N^1,O;O;\kappa^3O:N^1,O$)dilithium]-di- μ -aqua]

W. Starosta and J. Leciejewicz

Comment

Structural studies of divalent metal ion complexes with 3-aminopyrazine-2-carboxylate ligand have shown that the structures of Mg(II) and Ni(II) complexes consist of $ML_2(H_2O)_2$ monomers. In the Mg(II) complex, the ligand adopts a *cis* configuration (Ptasiewicz-Bąk & Leciejewicz, 1997), while in the Ni(II) complex, a *trans* configuration (Ptasiewicz-Bąk & Leciejewicz, 1999). Catenated polymeric molecular patterns have been reported in the structures of a Ca(II) complex (Leciejewicz *et al.*, 1997) and a Sr(II) complex (Leciejewicz *et al.*, 1998), in which metal ions are bridged by ligand carboxylate groups acting as bidentate. On the other hand, the structure of a Na(I) complex with the title ligand (Tayebee *et al.*, 2008) is three-dimensional polymeric with Na(I) ions linked by an extended bridging system formed mainly by coordinated water O atoms.

The title compound is composed of centrosymmetric dinuclear units, in which each of the two Li^I ions is chelated by a ligand *via* an N,O-bonding group. Its O atom acts as bidentate and bridges the other Li^I ion (Fig. 1). The dinuclear unit is nearly planar with r.m.s. of 0.0125 (2) Å. The Li^I ion is also coordinated by two water O atoms, which bridge the dinuclear units into molecular columns along two bridging pathways propagating in the *b*-axis direction (Fig. 2). The coordination geometry of the Li^I ion is trigonal bipyramidal, with the equatorial plane formed by O1, O3, O3ⁱⁱ and with N1 and O1ⁱ at the axial positions [symmetry codes: (i) 1-x, 1-y, 1-z; (ii) x, y-1, z]. The Li—O and Li—N bond distances (Table 1) and bond angles are typical for Li(I) complexes with carboxylate ligands (see, for example, Tombul *et al.*, 2008). The columns are linked by a network of hydrogen bonds, in which water O atoms are donors and the non-bonded carboxylate O atoms in adjacent columns act as acceptors. A weak hydrogen bond links an amino N atom with a hetero-ring N atom in the adjacent column. An intramolecular hydrogen bond which operates between the amino N3 atom and the non-bonding carboxylate O2 atom is also observed (Table 2).

Experimental

The title compound was synthesized by reacting 50 ml of boiling aqueous solutions, one containing 1 mmol of 3-aminopyrazine-2-carboxylic acid (Aldrich), the other containing 1 mmol of lithium hydroxide (Aldrich). The mixture was boiled under reflux for 3 h and after cooling to room temperature, filtered and left to crystallize. A few days later, colourless single crystals in the form of flat needles were found after evaporation to dryness. They were extracted, washed with cold ethanol and dried in air. A crystal used for X-ray data collection was cut to adopt the shape of a flat plate.

Refinement

Water H atoms were found from difference Fourier maps and their coordinates were refined with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{O})$. H atoms attached to C and N atoms were positioned geometrically and refined as riding, with C—H = 0.93 and N—H = 0.86 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C,N})$.

supplementary materials

Figures




Fig. 1. The dinuclear unit of the title compound. Displacement ellipsoids are drawn at the 50% probability level. [Symmetry codes: (i) 1-x, 1-y, 1-z; (ii) x, -1+y, z; (iii) 1-x, 2-y, 1-z.]




Fig. 2. Packing diagram of the title compound.

catena-Poly[[bis(μ -3-aminopyrazine-2-carboxylato)- $\kappa^3N^1,O;O;\kappa^3O:N^1,O$)dilithium]-di- μ -aqua]

Crystal data

[Li(C ₅ H ₄ N ₃ O ₂)(H ₂ O)]	$F(000) = 336$
$M_r = 163.07$	$D_x = 1.652 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 25 reflections
$a = 14.279 (3) \text{ \AA}$	$\theta = 6\text{--}15^\circ$
$b = 3.6000 (7) \text{ \AA}$	$\mu = 0.13 \text{ mm}^{-1}$
$c = 13.300 (3) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 106.43 (3)^\circ$	Plate, colourless
$V = 655.7 (2) \text{ \AA}^3$	$0.26 \times 0.21 \times 0.04 \text{ mm}$
$Z = 4$	

Data collection

Kuma KM-4 four-circle diffractometer	1297 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube graphite	$R_{\text{int}} = 0.017$
profile data from ω -20 scans	$\theta_{\text{max}} = 30.1^\circ, \theta_{\text{min}} = 1.5^\circ$
Absorption correction: analytical (<i>CrysAlis RED</i> ; Oxford Diffraction, 2006)	$h = -19 \rightarrow 19$
	$k = -5 \rightarrow 0$

$T_{\min} = 0.980$, $T_{\max} = 0.994$
 1997 measured reflections
 1913 independent reflections

$l = 0 \rightarrow 18$
 3 standard reflections every 200 reflections
 intensity decay: 7.3%

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.147$

Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.1051P)^2 + 0.022P]$
 where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$
 1913 reflections
 115 parameters
 3 restraints

$\Delta\rho_{\max} = 0.51 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.39 \text{ e \AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C2	0.25443 (8)	0.4711 (3)	0.39664 (9)	0.0184 (3)
O2	0.32090 (8)	0.2294 (4)	0.26435 (8)	0.0336 (3)
N1	0.27984 (8)	0.6065 (3)	0.49343 (8)	0.0216 (3)
N2	0.08397 (8)	0.5364 (4)	0.38343 (10)	0.0285 (3)
N3	0.12597 (9)	0.2931 (4)	0.24154 (10)	0.0320 (3)
H2	0.0649	0.2745	0.2090	0.038*
H1	0.1690	0.2235	0.2115	0.038*
O1	0.42229 (7)	0.4275 (4)	0.41421 (8)	0.0352 (3)
C7	0.33887 (9)	0.3657 (4)	0.35429 (10)	0.0218 (3)
C3	0.15415 (9)	0.4307 (4)	0.33936 (10)	0.0217 (3)
C6	0.20958 (10)	0.7070 (4)	0.53653 (11)	0.0254 (3)
H6	0.2262	0.8010	0.6044	0.030*
C5	0.11268 (10)	0.6720 (4)	0.48077 (12)	0.0284 (3)
H5	0.0653	0.7459	0.5123	0.034*
Li1	0.43334 (19)	0.6091 (9)	0.5591 (2)	0.0370 (6)
O3	0.44058 (8)	1.0866 (3)	0.64699 (9)	0.0338 (3)
H32	0.4986 (11)	1.117 (6)	0.6825 (15)	0.041*
H31	0.4047 (13)	1.146 (6)	0.6882 (14)	0.041*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C2	0.0170 (5)	0.0153 (5)	0.0223 (6)	-0.0007 (4)	0.0048 (4)	0.0018 (4)
O2	0.0301 (5)	0.0450 (7)	0.0277 (5)	-0.0089 (5)	0.0114 (4)	-0.0120 (5)
N1	0.0209 (5)	0.0192 (5)	0.0242 (5)	0.0012 (4)	0.0057 (4)	-0.0006 (4)
N2	0.0194 (5)	0.0264 (6)	0.0390 (6)	0.0012 (4)	0.0071 (4)	0.0025 (5)

supplementary materials

N3	0.0227 (5)	0.0401 (7)	0.0290 (6)	-0.0056 (5)	0.0002 (4)	-0.0052 (5)
O1	0.0183 (5)	0.0522 (7)	0.0338 (5)	0.0004 (5)	0.0052 (4)	-0.0142 (5)
C7	0.0202 (6)	0.0209 (6)	0.0242 (5)	-0.0021 (4)	0.0064 (4)	-0.0015 (5)
C3	0.0199 (5)	0.0171 (5)	0.0262 (6)	-0.0019 (4)	0.0033 (4)	0.0034 (5)
C6	0.0282 (6)	0.0225 (7)	0.0273 (6)	0.0020 (5)	0.0108 (5)	-0.0020 (5)
C5	0.0238 (6)	0.0237 (7)	0.0408 (8)	0.0031 (5)	0.0140 (5)	0.0006 (6)
Li1	0.0256 (12)	0.0484 (17)	0.0355 (13)	0.0016 (11)	0.0060 (10)	-0.0120 (12)
O3	0.0256 (5)	0.0411 (7)	0.0352 (6)	-0.0012 (5)	0.0092 (4)	-0.0069 (5)

Geometric parameters (\AA , $^\circ$)

C2—N1	1.3274 (16)	C6—C5	1.378 (2)
C2—C3	1.4263 (17)	C6—H6	0.9300
C2—C7	1.5164 (17)	C5—H5	0.9300
O2—C7	1.2505 (17)	Li1—N1	2.118 (3)
N1—C6	1.3385 (17)	Li1—O1	1.999 (3)
N2—C5	1.335 (2)	Li1—O1 ⁱ	1.995 (3)
N2—C3	1.3510 (18)	Li1—O3	2.065 (3)
N3—C3	1.3431 (18)	Li1—O3 ⁱⁱ	2.201 (3)
N3—H2	0.8600	Li1—Li1 ⁱ	2.900 (5)
N3—H1	0.8600	O3—H32	0.837 (15)
O1—C7	1.2515 (17)	O3—H31	0.875 (14)
N1—C2—C3	120.87 (11)	N2—C5—H5	118.6
N1—C2—C7	115.10 (11)	C6—C5—H5	118.6
C3—C2—C7	124.03 (11)	O1 ⁱ —Li1—O1	86.88 (11)
C2—N1—C6	118.83 (11)	O1 ⁱ —Li1—O3	94.05 (12)
C2—N1—Li1	111.64 (11)	O1—Li1—O3	142.73 (18)
C6—N1—Li1	129.48 (11)	O1 ⁱ —Li1—N1	165.94 (15)
C5—N2—C3	117.50 (11)	O1—Li1—N1	79.08 (10)
C3—N3—H2	120.0	O3—Li1—N1	96.74 (12)
C3—N3—H1	120.0	O1 ⁱ —Li1—O3 ⁱⁱ	87.61 (12)
H2—N3—H1	120.0	O1—Li1—O3 ⁱⁱ	102.21 (14)
C7—O1—Li1 ⁱ	148.32 (12)	O3—Li1—O3 ⁱⁱ	115.07 (14)
C7—O1—Li1	118.26 (11)	N1—Li1—O3 ⁱⁱ	95.90 (13)
Li1 ⁱ —O1—Li1	93.13 (11)	O1 ⁱ —Li1—Li1 ⁱ	43.49 (8)
O2—C7—O1	125.43 (12)	O1—Li1—Li1 ⁱ	43.38 (8)
O2—C7—C2	118.94 (12)	O3—Li1—Li1 ⁱ	126.66 (18)
O1—C7—C2	115.63 (11)	N1—Li1—Li1 ⁱ	122.46 (16)
N3—C3—N2	117.93 (12)	O3 ⁱⁱ —Li1—Li1 ⁱ	96.72 (16)
N3—C3—C2	122.37 (12)	Li1—O3—Li1 ⁱⁱⁱ	115.07 (14)
N2—C3—C2	119.69 (12)	Li1—O3—H32	108.2 (15)
N1—C6—C5	120.32 (12)	Li1 ⁱⁱⁱ —O3—H32	94.4 (16)
N1—C6—H6	119.8	Li1—O3—H31	127.7 (15)
C5—C6—H6	119.8	Li1 ⁱⁱⁱ —O3—H31	100.2 (15)
N2—C5—C6	122.78 (13)	H32—O3—H31	106.0 (15)

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x, y-1, z$; (iii) $x, y+1, z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O3—H31…O2 ^{iv}	0.88 (1)	1.83 (1)	2.7028 (16)	175.(2)
O3—H32…O1 ^v	0.84 (2)	2.54 (2)	2.9083 (17)	108.(2)
N3—H1…O2	0.86	2.08	2.7229 (17)	131
N3—H2…N2 ^{vi}	0.86	2.30	3.1278 (19)	162

Symmetry codes: (iv) $x, -y+3/2, z+1/2$; (v) $-x+1, -y+2, -z+1$; (vi) $-x, y-1/2, -z+1/2$.

supplementary materials

Fig. 1




Fig. 2

