

{*N,N'*-Bis[1-(2-pyridyl)ethylidene]ethane-1,2-diamine- κ^4N,N',N'',N''' (thiocyanato- κN)zinc(II) perchlorate}

Fu-Ming Wang

Department of Chemistry, Dezhou University, Dezhou Shandong 253023, People's Republic of China

Correspondence e-mail: wfm99999@126.com


Received 6 June 2010; accepted 7 June 2010

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(C-C) = 0.008$ Å; disorder in main residue; R factor = 0.056; wR factor = 0.170; data-to-parameter ratio = 16.1.

In the title compound, $[Zn(NCS)(C_{16}H_{18}N_4)]ClO_4$, the Zn^{II} atom is five-coordinated by four N atoms of the Schiff base ligand *N,N'*-bis[1-(2-pyridyl)ethylidene]ethane-1,2-diamine in the basal plane, and by the N atom of a thiocyanate ligand at the apical position, forming a distorted square-pyramidal geometry. The r.m.s. deviation from a plane through the four N atoms of the Schiff base is 0.015 (3) Å, and the deviation of the Ni atom from that plane is 0.591 (2) Å. Bond lengths are comparable with those observed in similar zinc(II) complexes with Schiff bases. The two methylene C atoms of the ethane-1,2-diamine bridge of the Schiff base ligand are disordered over two sites with occupancies of 0.587 (3) and 0.413 (3).

Related literature

For background to Schiff base compounds and their applications, see: Ruck & Jacobsen (2002); Mukhopadhyay *et al.* (2003); Polt *et al.* (2003); Mukherjee *et al.* (2001). For complexes derived from *N,N'*-bis(1-(pyridin-2-yl)ethylidene)-ethane-1,2-diamine, see: Gourbatsis *et al.* (1998); Louloudi *et al.* (1999); Karmakar *et al.* (2002); Banerjee *et al.* (2004). For bond lengths in similar zinc(II) complexes with Schiff bases, see: Ghosh *et al.* (2006); Chen *et al.* (2005). For the synthesis of the Schiff base ligand, see: Gourbatsis *et al.* (1990).



Experimental

Crystal data

$[Zn(NCS)(C_{16}H_{18}N_4)]ClO_4$	$V = 2076.9$ (7) Å ³
$M_r = 489.24$	$Z = 4$
Monoclinic, $P_{\bar{2}1}/c$	Mo $K\alpha$ radiation
$a = 8.685$ (2) Å	$\mu = 1.45$ mm ⁻¹
$b = 13.963$ (3) Å	$T = 298$ K
$c = 17.374$ (2) Å	$0.32 \times 0.30 \times 0.30$ mm
$\beta = 99.690$ (3)°	

Data collection

Bruker SMART CCD area-detector diffractometer	16681 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	4529 independent reflections
$R_{\text{int}} = 0.056$	2534 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.655$, $T_{\max} = 0.671$	

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$	48 restraints
$wR(F^2) = 0.170$	H-atom parameters constrained
$S = 1.02$	$\Delta\rho_{\max} = 0.50$ e Å ⁻³
4529 reflections	$\Delta\rho_{\min} = -0.66$ e Å ⁻³
282 parameters	

Table 1
Selected geometric parameters (Å, °).

Zn1—N5	1.982 (5)	Zn1—N2	2.102 (4)
Zn1—N3	2.099 (4)	Zn1—N4	2.115 (4)
Zn1—N1	2.100 (4)		
N5—Zn1—N3	109.08 (17)	N1—Zn1—N2	77.12 (15)
N5—Zn1—N1	105.95 (16)	N5—Zn1—N4	101.91 (17)
N3—Zn1—N1	141.17 (14)	N3—Zn1—N4	77.11 (14)
N5—Zn1—N2	110.56 (19)	N1—Zn1—N4	111.48 (15)
N3—Zn1—N2	75.37 (15)	N2—Zn1—N4	142.71 (17)

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1998); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported financially by Dezhou University, People's Republic of China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5017).

References

- Banerjee, S., Gangopadhyay, J., Lu, C.-Z., Chen, J.-T. & Ghosh, A. (2004). *Eur. J. Inorg. Chem.* pp. 2533–2541.
- Bruker (1998). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, G., Bai, Z.-P. & Qu, S.-J. (2005). *Acta Cryst. E* **61**, m2483–m2484.
- Ghosh, R., Rahaman, S. H., Lin, C.-N., Lu, T.-H. & Ghosh, B. K. (2006). *Polyhedron*, **25**, 3104–3112.
- Gourbatsis, S., Hadjiliadis, N., Perlepes, S. P., Garoufis, A. & Butler, I. S. (1998). *Transition Met. Chem.* **23**, 599–604.
- Gourbatsis, S., Perlepes, S. P., Hadjiliadis, N. & Kalkanis, G. (1990). *Transition Met. Chem.* **15**, 300–308.
- Karmakar, T. K., Chandra, S. K., Ribas, J., Mostafa, G., Lu, T. H. & Ghosh, B. K. (2002). *Chem. Commun.* pp. 2364–2365.
- Louloudi, M., Nastopoulos, V., Gourbatsis, S., Perlepes, S. P. & Hadjiliadis, N. (1999). *Inorg. Chem. Commun.* **2**, 479–483.
- Mukherjee, P. S., Dalai, S., Mostafa, G., Lu, T.-H., Rentschler, E. & Chaudhuri, N. R. (2001). *New J. Chem.* **25**, 1203–1207.
- Mukhopadhyay, S., Mandal, D., Ghosh, D., Goldberg, I. & Chaudhury, M. (2003). *Inorg. Chem.* **42**, 8439–8445.
- Polt, R., Kelly, B. D., Dangel, B. D., Tadiikonda, U. B., Ross, R. E., Raitsimring, A. M. & Astashkin, A. V. (2003). *Inorg. Chem.* **42**, 566–574.
- Ruck, R. T. & Jacobsen, E. N. (2002). *J. Am. Chem. Soc.* **124**, 2882–2883.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2010). E66, m778-m779 [doi:10.1107/S1600536810021628]

{*N,N'*-Bis[1-(2-pyridyl)ethylidene]ethane-1,2-diamine- κ^4N,N',N'',N''' }(thiocyanato- κN)zinc(II) perchlorate

F.-M. Wang

Comment

Metal complexes with Schiff bases have been known since 1840. The Schiff bases and their complexes have played an important role in the development of coordination chemistry, biological and material sciences (Ruck & Jacobsen, 2002; Mukhopadhyay *et al.*, 2003; Polt *et al.*, 2003; Mukherjee *et al.*, 2001). Several complexes derived from *N,N'*-bis(1-(pyridin-2-yl)ethylidene)ethane-1,2-diamine have been reported (Gourbatsis *et al.*, 1998; Louloudi *et al.*, 1999; Karmakar *et al.*, 2002; Banerjee *et al.*, 2004). In this paper, the title new zinc(II) complex is reported.

The title compound consists of a mononuclear zinc(II) complex cation and a perchlorate anion, Fig. 1. The Zn^{II} atom is five-coordinated by four N atoms of the Schiff base ligand *N,N'*-bis(1-(pyridin-2-yl)ethylidene)ethane-1,2-diamine, and by one N atom of a thiocyanate ligand, forming a square pyramidal geometry. The coordinate bond lengths (Table 1) are comparable to those observed in other similar zinc(II) complexes with Schiff bases (Ghosh *et al.*, 2006; Chen *et al.*, 2005).

Experimental

The Schiff base ligand *N,N'*-bis(1-(pyridin-2-yl)ethylidene)ethane-1,2-diamine was synthesized according to the literature method (Gourbatsis *et al.*, 1990). To a stirred methanol solution of the Schiff base ligand (1.0 mmol, 0.266 g) was added a methanol solution of zinc(II) perchlorate (1.0 mmol, 0.390 g) and ammonium thiocyanate (1.0 mmol, 0.076 g). The mixture was boiled under reflux for 2 h, then cooled to room temperature. Colourless block-like single crystals, suitable for X-ray diffraction, were formed after slow evaporation of the solution in air for a few days.

Refinement

Hydrogen atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with C–H distances of 0.93–0.97 Å, and with $U_{\text{iso}}(\text{H})$ set at $1.2U_{\text{eq}}(\text{C})$ and $1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$. The C8 and C9 atoms are disordered over two sites with occupancies of 0.587 (3) and 0.413 (3).

Figures




Fig. 1. The asymmetric unit of the title compound, showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. Only the major component of the disordered group is shown.

supplementary materials

{N,N'-Bis[1-(2-pyridyl)ethylidene]ethane-1,2-diamine- κ⁴N,N',N'',N'''}(thiocyanato- κN)zinc(II) perchlorate

Crystal data

[Zn(NCS)(C ₁₆ H ₁₈ N ₄)]ClO ₄	$F(000) = 1000$
$M_r = 489.24$	$D_x = 1.565 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 2719 reflections
$a = 8.685 (2) \text{ \AA}$	$\theta = 2.4\text{--}25.0^\circ$
$b = 13.963 (3) \text{ \AA}$	$\mu = 1.45 \text{ mm}^{-1}$
$c = 17.374 (2) \text{ \AA}$	$T = 298 \text{ K}$
$\beta = 99.690 (3)^\circ$	Block, colourless
$V = 2076.9 (7) \text{ \AA}^3$	$0.32 \times 0.30 \times 0.30 \text{ mm}$
$Z = 4$	

Data collection

Bruker SMART CCD area-detector diffractometer	4529 independent reflections
Radiation source: fine-focus sealed tube graphite	2534 reflections with $I > 2\sigma(I)$
ω scan	$R_{\text{int}} = 0.056$
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	$\theta_{\text{max}} = 27.0^\circ, \theta_{\text{min}} = 2.4^\circ$
$T_{\text{min}} = 0.655, T_{\text{max}} = 0.671$	$h = -11 \rightarrow 10$
16681 measured reflections	$k = -17 \rightarrow 16$
	$l = -22 \rightarrow 22$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.056$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.170$	H-atom parameters constrained
$S = 1.02$	$w = 1/[\sigma^2(F_o^2) + (0.0792P)^2 + 0.7526P]$
4529 reflections	where $P = (F_o^2 + 2F_c^2)/3$
282 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
48 restraints	$\Delta\rho_{\text{max}} = 0.50 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.66 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations

between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Zn1	0.40268 (7)	0.18336 (3)	0.23821 (3)	0.0532 (2)	
Cl1	0.13446 (17)	0.32198 (10)	0.93392 (9)	0.0761 (4)	
S1	0.80304 (19)	0.18105 (12)	0.08648 (10)	0.0888 (5)	
O1	0.1863 (9)	0.2376 (4)	0.9693 (3)	0.170 (3)	
O2	0.2281 (6)	0.3965 (4)	0.9651 (3)	0.145 (2)	
O3	0.1346 (6)	0.3155 (3)	0.8524 (2)	0.0975 (14)	
O4	-0.0172 (5)	0.3397 (4)	0.9480 (3)	0.1246 (19)	
N1	0.2700 (4)	0.3093 (2)	0.2175 (2)	0.0516 (9)	
N4	0.2661 (4)	0.0604 (3)	0.2040 (2)	0.0547 (10)	
N5	0.5483 (6)	0.1840 (3)	0.1618 (3)	0.0725 (13)	
C1	0.1666 (6)	0.3311 (4)	0.1535 (3)	0.0640 (13)	
H1	0.1435	0.2853	0.1144	0.077*	
C2	0.0928 (6)	0.4194 (4)	0.1434 (3)	0.0724 (15)	
H2	0.0234	0.4328	0.0978	0.087*	
C3	0.1231 (7)	0.4857 (4)	0.2009 (4)	0.0762 (16)	
H3	0.0728	0.5447	0.1959	0.091*	
C4	0.2296 (6)	0.4645 (3)	0.2672 (3)	0.0673 (14)	
H4	0.2531	0.5097	0.3067	0.081*	
C5	0.3015 (6)	0.3754 (3)	0.2744 (3)	0.0524 (11)	
C6	0.4210 (6)	0.3474 (3)	0.3435 (3)	0.0608 (13)	
C7	0.4484 (8)	0.4109 (4)	0.4138 (3)	0.0881 (19)	
H7A	0.3514	0.4216	0.4320	0.132*	
H7B	0.4902	0.4710	0.4003	0.132*	
H7C	0.5212	0.3808	0.4543	0.132*	
N2	0.4901 (5)	0.2687 (3)	0.3355 (3)	0.0713 (12)	0.59 (3)
N3	0.4830 (4)	0.0850 (3)	0.3273 (2)	0.0537 (10)	0.59 (3)
C8	0.573 (2)	0.2217 (8)	0.4079 (7)	0.076 (4)	0.59 (3)
H8A	0.5077	0.2231	0.4481	0.092*	0.59 (3)
H8B	0.6691	0.2557	0.4272	0.092*	0.59 (3)
C9	0.6085 (18)	0.1192 (11)	0.3889 (10)	0.073 (6)	0.59 (3)
H9A	0.7083	0.1159	0.3709	0.088*	0.59 (3)
H9B	0.6141	0.0795	0.4351	0.088*	0.59 (3)
N2'	0.4901 (5)	0.2687 (3)	0.3355 (3)	0.0713 (12)	0.41 (3)
N3'	0.4830 (4)	0.0850 (3)	0.3273 (2)	0.0537 (10)	0.41 (3)
C8'	0.6313 (16)	0.2226 (10)	0.3782 (13)	0.058 (5)	0.41 (3)
H8'A	0.6739	0.2587	0.4247	0.070*	0.41 (3)
H8'B	0.7109	0.2158	0.3456	0.070*	0.41 (3)
C9'	0.571 (3)	0.1262 (15)	0.3990 (8)	0.058 (6)	0.41 (3)

supplementary materials

H9'A	0.6570	0.0846	0.4201	0.070*	0.41 (3)
H9'B	0.5035	0.1335	0.4379	0.070*	0.41 (3)
C10	0.4124 (6)	0.0058 (3)	0.3271 (3)	0.0531 (12)	
C11	0.4348 (8)	-0.0665 (4)	0.3924 (3)	0.0878 (19)	
H11A	0.5251	-0.0497	0.4299	0.132*	
H11B	0.4493	-0.1290	0.3716	0.132*	
H11C	0.3442	-0.0670	0.4174	0.132*	
C12	0.2956 (5)	-0.0134 (3)	0.2555 (3)	0.0518 (11)	
C13	0.2222 (6)	-0.1010 (4)	0.2399 (3)	0.0714 (15)	
H13	0.2424	-0.1509	0.2756	0.086*	
C14	0.1198 (7)	-0.1136 (4)	0.1718 (4)	0.089 (2)	
H14	0.0677	-0.1716	0.1614	0.107*	
C15	0.0947 (7)	-0.0406 (5)	0.1193 (4)	0.093 (2)	
H15	0.0289	-0.0491	0.0717	0.112*	
C16	0.1684 (6)	0.0465 (4)	0.1376 (3)	0.0775 (16)	
H16	0.1487	0.0968	0.1022	0.093*	
C17	0.6548 (7)	0.1821 (3)	0.1302 (3)	0.0544 (12)	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0591 (4)	0.0415 (3)	0.0582 (4)	-0.0002 (3)	0.0079 (3)	0.0004 (2)
Cl1	0.0691 (9)	0.0799 (10)	0.0767 (9)	-0.0045 (8)	0.0053 (7)	-0.0098 (8)
S1	0.0678 (10)	0.1090 (13)	0.0936 (12)	0.0072 (9)	0.0252 (9)	0.0140 (9)
O1	0.247 (6)	0.129 (4)	0.128 (4)	0.090 (4)	0.019 (4)	0.038 (4)
O2	0.128 (4)	0.170 (5)	0.139 (4)	-0.084 (4)	0.031 (3)	-0.071 (4)
O3	0.117 (3)	0.115 (3)	0.063 (2)	-0.004 (3)	0.023 (2)	-0.015 (2)
O4	0.054 (3)	0.207 (6)	0.119 (4)	0.000 (3)	0.031 (3)	-0.013 (3)
N1	0.058 (2)	0.045 (2)	0.052 (2)	0.0013 (18)	0.0111 (18)	0.0031 (17)
N4	0.050 (2)	0.050 (2)	0.062 (2)	0.0005 (18)	0.0044 (19)	-0.0043 (19)
N5	0.088 (3)	0.055 (3)	0.080 (3)	0.003 (2)	0.032 (3)	0.009 (2)
C1	0.072 (3)	0.062 (3)	0.057 (3)	0.003 (3)	0.009 (3)	0.001 (2)
C2	0.077 (4)	0.062 (3)	0.077 (4)	0.018 (3)	0.008 (3)	0.017 (3)
C3	0.085 (4)	0.058 (3)	0.088 (4)	0.022 (3)	0.017 (3)	0.013 (3)
C4	0.076 (4)	0.043 (3)	0.086 (4)	0.001 (3)	0.024 (3)	-0.005 (3)
C5	0.057 (3)	0.045 (3)	0.058 (3)	-0.009 (2)	0.018 (2)	0.001 (2)
C6	0.068 (3)	0.043 (3)	0.069 (3)	-0.012 (2)	0.002 (3)	-0.001 (2)
C7	0.115 (5)	0.061 (3)	0.079 (4)	-0.008 (3)	-0.009 (3)	-0.017 (3)
N2	0.064 (3)	0.053 (3)	0.088 (3)	-0.001 (2)	-0.015 (2)	-0.004 (2)
N3	0.055 (2)	0.049 (2)	0.055 (2)	-0.0005 (19)	0.0049 (18)	-0.0021 (18)
C8	0.073 (8)	0.071 (6)	0.080 (7)	-0.002 (5)	-0.002 (6)	-0.008 (5)
C9	0.080 (9)	0.063 (8)	0.071 (8)	0.004 (6)	-0.003 (6)	-0.007 (5)
N2'	0.064 (3)	0.053 (3)	0.088 (3)	-0.001 (2)	-0.015 (2)	-0.004 (2)
N3'	0.055 (2)	0.049 (2)	0.055 (2)	-0.0005 (19)	0.0049 (18)	-0.0021 (18)
C8'	0.044 (7)	0.058 (7)	0.069 (8)	-0.001 (6)	0.001 (6)	-0.010 (6)
C9'	0.068 (9)	0.051 (9)	0.051 (8)	0.005 (7)	-0.003 (7)	0.004 (6)
C10	0.065 (3)	0.039 (2)	0.057 (3)	0.008 (2)	0.017 (2)	0.000 (2)
C11	0.129 (6)	0.055 (3)	0.076 (4)	-0.006 (3)	0.008 (4)	0.009 (3)

C12	0.048 (3)	0.043 (3)	0.068 (3)	0.004 (2)	0.020 (2)	-0.007 (2)
C13	0.065 (4)	0.050 (3)	0.098 (4)	-0.007 (3)	0.013 (3)	-0.006 (3)
C14	0.076 (4)	0.058 (4)	0.129 (6)	-0.015 (3)	0.001 (4)	-0.021 (4)
C15	0.077 (4)	0.085 (5)	0.104 (5)	-0.011 (4)	-0.023 (4)	-0.023 (4)
C16	0.071 (4)	0.069 (4)	0.084 (4)	0.001 (3)	-0.011 (3)	-0.004 (3)
C17	0.065 (3)	0.038 (2)	0.058 (3)	0.001 (2)	0.002 (3)	0.007 (2)

Geometric parameters (Å, °)

Zn1—N5	1.982 (5)	C7—H7C	0.9600
Zn1—N3	2.099 (4)	N2—C8	1.492 (8)
Zn1—N1	2.100 (4)	N3—C10	1.264 (5)
Zn1—N2	2.102 (4)	N3—C9	1.472 (8)
Zn1—N4	2.115 (4)	C8—C9	1.513 (10)
Cl1—O1	1.370 (5)	C8—H8A	0.9700
Cl1—O2	1.374 (4)	C8—H8B	0.9700
Cl1—O4	1.401 (4)	C9—H9A	0.9700
Cl1—O3	1.419 (4)	C9—H9B	0.9700
S1—C17	1.601 (6)	C8'—C9'	1.512 (10)
N1—C1	1.340 (6)	C8'—H8'A	0.9700
N1—C5	1.347 (5)	C8'—H8'B	0.9700
N4—C16	1.327 (6)	C9'—H9'A	0.9700
N4—C12	1.359 (6)	C9'—H9'B	0.9700
N5—C17	1.153 (7)	C10—C12	1.492 (6)
C1—C2	1.387 (7)	C10—C11	1.508 (7)
C1—H1	0.9300	C11—H11A	0.9600
C2—C3	1.355 (7)	C11—H11B	0.9600
C2—H2	0.9300	C11—H11C	0.9600
C3—C4	1.382 (7)	C12—C13	1.385 (6)
C3—H3	0.9300	C13—C14	1.368 (8)
C4—C5	1.388 (6)	C13—H13	0.9300
C4—H4	0.9300	C14—C15	1.360 (8)
C5—C6	1.501 (7)	C14—H14	0.9300
C6—N2	1.271 (6)	C15—C16	1.387 (7)
C6—C7	1.495 (7)	C15—H15	0.9300
C7—H7A	0.9600	C16—H16	0.9300
C7—H7B	0.9600		
N5—Zn1—N3	109.08 (17)	C6—N2—Zn1	117.7 (3)
N5—Zn1—N1	105.95 (16)	C8—N2—Zn1	119.2 (5)
N3—Zn1—N1	141.17 (14)	C10—N3—C9	125.9 (9)
N5—Zn1—N2	110.56 (19)	C10—N3—Zn1	118.0 (3)
N3—Zn1—N2	75.37 (15)	C9—N3—Zn1	115.9 (8)
N1—Zn1—N2	77.12 (15)	N2—C8—C9	108.5 (11)
N5—Zn1—N4	101.91 (17)	N2—C8—H8A	110.0
N3—Zn1—N4	77.11 (14)	C9—C8—H8A	110.0
N1—Zn1—N4	111.48 (15)	N2—C8—H8B	110.0
N2—Zn1—N4	142.71 (17)	C9—C8—H8B	110.0
O1—Cl1—O2	110.3 (5)	H8A—C8—H8B	108.4
O1—Cl1—O4	108.8 (4)	N3—C9—C8	108.2 (9)

supplementary materials

O2—Cl1—O4	108.0 (3)	N3—C9—H9A	110.1
O1—Cl1—O3	109.7 (3)	C8—C9—H9A	110.1
O2—Cl1—O3	110.0 (3)	N3—C9—H9B	110.1
O4—Cl1—O3	110.1 (3)	C8—C9—H9B	110.1
C1—N1—C5	118.6 (4)	H9A—C9—H9B	108.4
C1—N1—Zn1	127.1 (3)	C9'—C8'—H8'A	111.3
C5—N1—Zn1	114.2 (3)	C9'—C8'—H8'B	111.3
C16—N4—C12	118.9 (4)	H8'A—C8'—H8'B	109.2
C16—N4—Zn1	127.5 (4)	C8'—C9'—H9'A	110.2
C12—N4—Zn1	113.3 (3)	C8'—C9'—H9'B	110.2
C17—N5—Zn1	166.6 (5)	H9'A—C9'—H9'B	108.5
N1—C1—C2	122.5 (5)	N3—C10—C12	114.9 (4)
N1—C1—H1	118.7	N3—C10—C11	125.6 (5)
C2—C1—H1	118.7	C12—C10—C11	119.4 (4)
C3—C2—C1	119.1 (5)	C10—C11—H11A	109.5
C3—C2—H2	120.5	C10—C11—H11B	109.5
C1—C2—H2	120.5	H11A—C11—H11B	109.5
C2—C3—C4	119.2 (5)	C10—C11—H11C	109.5
C2—C3—H3	120.4	H11A—C11—H11C	109.5
C4—C3—H3	120.4	H11B—C11—H11C	109.5
C3—C4—C5	119.7 (5)	N4—C12—C13	120.8 (4)
C3—C4—H4	120.1	N4—C12—C10	116.0 (4)
C5—C4—H4	120.1	C13—C12—C10	123.2 (4)
N1—C5—C4	121.0 (5)	C14—C13—C12	119.6 (5)
N1—C5—C6	115.9 (4)	C14—C13—H13	120.2
C4—C5—C6	123.1 (5)	C12—C13—H13	120.2
N2—C6—C7	126.1 (5)	C15—C14—C13	119.4 (5)
N2—C6—C5	114.4 (4)	C15—C14—H14	120.3
C7—C6—C5	119.5 (5)	C13—C14—H14	120.3
C6—C7—H7A	109.5	C14—C15—C16	119.2 (5)
C6—C7—H7B	109.5	C14—C15—H15	120.4
H7A—C7—H7B	109.5	C16—C15—H15	120.4
C6—C7—H7C	109.5	N4—C16—C15	122.0 (5)
H7A—C7—H7C	109.5	N4—C16—H16	119.0
H7B—C7—H7C	109.5	C15—C16—H16	119.0
C6—N2—C8	117.1 (7)	N5—C17—S1	179.2 (5)

Fig. 1

