

catena-Poly[[aqua(imidazole)-cadmium(II)]- μ_3 -benzene-1,3-dicarboxylato]

 Zhengfang Zeng^a and Hongyan Xu^{b*}

^aSchool of Sciences, Nanchang University, Nanchang, Jiangxi 330031, People's Republic of China, and ^bFujian Institute of Research on the Structure of Matter, Chinese Academy of Sciences, Fuzhou, Fujian 350002, People's Republic of China
Correspondence e-mail: hongyan@fjirsm.ac.cn


Received 21 April 2010; accepted 9 June 2010

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.028; wR factor = 0.072; data-to-parameter ratio = 11.3.

In the title compound, $[\text{Cd}(\text{C}_8\text{H}_4\text{O}_4)(\text{C}_3\text{H}_4\text{N}_2)(\text{H}_2\text{O})]_n$, the Cd^{II} ion is seven-coordinated by five O atoms from three crystallographically independent benzene-1,3-carboxylate ligands, one N atom from the imidazole ligand and one coordinated water molecule. Neighboring Cd^{II} ions are bridged by the benzene-1,3-dicarboxylate ligands, forming a zigzag polymeric chain structure. These chains are further extended into a three-dimensional supramolecular structure through $\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For the synthesis, see: Yaghi *et al.* (1998). For related structures, see: Ma *et al.* (2008); Wang *et al.* (2008).



Experimental

Crystal data

$[\text{Cd}(\text{C}_8\text{H}_4\text{O}_4)(\text{C}_3\text{H}_4\text{N}_2)(\text{H}_2\text{O})]$

$M_r = 362.61$

Triclinic, $P\bar{1}$

$a = 8.2616(8)\text{ \AA}$

$b = 8.3138(8)\text{ \AA}$

$c = 10.235(1)\text{ \AA}$

$\alpha = 67.017(2)^\circ$

$\beta = 68.176(2)^\circ$

$\gamma = 81.054(2)^\circ$

$V = 600.76(10)\text{ \AA}^3$

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 1.84\text{ mm}^{-1}$

$T = 293\text{ K}$
 $0.27 \times 0.18 \times 0.06\text{ mm}$

Data collection

Bruker APEX CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.536$, $T_{\max} = 1.000$

3166 measured reflections
2086 independent reflections
2049 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.072$
 $S = 1.15$
2086 reflections
184 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.41\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.79\text{ e \AA}^{-3}$

Table 1
Selected bond lengths (Å).

Cd1–N1	2.216 (3)	Cd1–O1	2.413 (3)
Cd1–O3 ⁱ	2.251 (3)	Cd1–O1 ⁱⁱ	2.626 (3)
Cd1–O2 ⁱⁱ	2.311 (3)	Cd1–O4 ⁱ	2.663 (3)
Cd1–O5	2.394 (3)		

Symmetry codes: (i) $-x, -y + 1, -z + 1$; (ii) $-x, -y + 1, -z + 2$.

Table 2
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O5–H5A \cdots O2 ⁱⁱⁱ	0.84 (8)	2.07 (8)	2.809 (4)	146 (7)
O5–H5B \cdots O3 ^{iv}	0.87 (8)	1.96 (8)	2.786 (4)	158 (7)
N2–H2 \cdots O4 ^v	0.85 (7)	2.00 (7)	2.854 (5)	175 (6)

Symmetry codes: (iii) $x, y - 1, z$; (iv) $x, y - 1, z + 1$; (v) $x + 1, y, z$.

Data collection: *SMART* (Siemens, 1996); cell refinement: *SAINT* (Siemens, 1996); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the National Natural Science Foundation of China (20971122) and the State Key Laboratory of Structural Chemistry for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LX2148).

References

- Ma, S., Sun, D., Simmons, J. M., Collier, C. D., Yuan, D. & Zhou, H. C. (2008). *J. Am. Chem. Soc.* **130**, 1012–1016.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Siemens (1996). *SMART* and *SAINT*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Wang, J., Lin, Z., Ou, Y. C., Yang, N. L., Zhang, Y. H. & Tong, M. L. (2008). *Inorg. Chem.* **47**, 190–199.
- Yaghi, O. M., Li, H., Davis, C., Richardson, D. & Groy, T. L. (1998). *Acc. Chem. Res.* **31**, 474–484.

supplementary materials

Acta Cryst. (2010). E66, m794 [doi:10.1107/S1600536810022117]

catena-Poly[[aqua(imidazole)cadmium(II)]- μ_3 -benzene-1,3-dicarboxylato]

Z. Zeng and H. Xu

Comment

In recent years, considerable effort was paid in the study of metal-organic hybrid materials with the extended supermolecular framework structures, owing to their intriguing network topologies and potential application in adsorption, molecular recognition, catalysis and magnetism (Yaghi *et al.*, 1998). Currently, carboxylic acids have been widely used as polydentate and bridging ligands since they can bind metal ions in diverse modes and play an important role in adjusting various topologies structures. A number of promising supramolecular complexes have been designed and constructed by using carboxylic acids as favorable conditions. (Ma *et al.*, 2008; Wang *et al.*, 2008). In this paper, we report the synthesis and structural characterization of the title compound in which the isophthalic acid ligand displayed its good coordination ability and diverse coordination modes.

The Cd^{II} ion is six-coordinated by four O atoms from three crystallographically independent benzene-1,3-dicarboxylate ligands and one N atom from the chelating imidazole ligand and one coordinated water molecule (Fig. 1 & Table 1). The neighboring Cd^I ion is bridged by the benzene-1,3-dicarboxylate ligands to form a one-dimensional chain structure. In the crystal structure, the adjacent chains are linked via O–H···O and N–H···O hydrogen bonds (Fig. 2 & Table 2) resulting in the formation of a three dimensional supramolecular structure.

Experimental

A 10 ml aqueous solution of imidazole (0.014 g, 0.20 mmol) and isophthalic acid (0.034 g, 0.020 mmol) was slowly added into cadmium nitrate (0.062 g, 0.20 mmol) solution in methanol (10 ml), and the mixed solution was stirred for 20 min and then was heated in a 30 ml Teflon-line autoclave under autogeneous pressure at 423 K for 3 d. After cooling to room temperature, colorless block crystals were obtained. (yield 38%).

Refinement

H atoms bound to H₂O and NH atoms were located in a difference map and refined freely. Other H atoms were positioned geometrically and refined using a riding model, with C–H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of (I), with atom labels and 50% probability displacement ellipsoids for non-H atoms. [Symmetry codes: (i) $-x, -y+1, -z+2$; (ii) $-x, -y+1, -z+1$; (iii) $x, y, z+1$.]

supplementary materials




Fig. 2. The crystal packing of (I), showing one layer of molecules connected by O–H···O hydrogen bonds (dashed lines). H atoms not involved in hydrogen bonding have been omitted. [Symmetry codes: (i) $x, y-1, z$; (ii) $x, y-1, z+1$; (iii) $x+1, y, z$; (iv) $-x, -y+1, -z+1$; (v) $-x, -y+1, -z+2$.]

catena-Poly[[aqua(imidazole)cadmium(II)]- μ_3 -benzene-1,3-dicarboxylato]

Crystal data

[Cd(C₈H₄O₄)(C₃H₄N₂)(H₂O)]

$Z = 2$

$M_r = 362.61$

$F(000) = 356$

Triclinic, $P\bar{1}$

$D_x = 2.005 \text{ Mg m}^{-3}$

Hall symbol: -P 1

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

$a = 8.2616 (8) \text{ \AA}$

Cell parameters from 3126 reflections

$b = 8.3138 (8) \text{ \AA}$

$\theta = 3.3\text{--}27.5^\circ$

$c = 10.235 (1) \text{ \AA}$

$\mu = 1.84 \text{ mm}^{-1}$

$\alpha = 67.017 (2)^\circ$

$T = 293 \text{ K}$

$\beta = 68.176 (2)^\circ$

Block, yellow

$\gamma = 81.054 (2)^\circ$

$0.27 \times 0.18 \times 0.06 \text{ mm}$

$V = 600.76 (10) \text{ \AA}^3$

Data collection

Bruker APEX CCD
diffractometer

2086 independent reflections

Radiation source: fine-focus sealed tube
graphite

2049 reflections with $I > 2\sigma(I)$

Detector resolution: 10 pixels mm^{-1}

$R_{\text{int}} = 0.019$

ω scans

$\theta_{\text{max}} = 25.0^\circ, \theta_{\text{min}} = 2.3^\circ$

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)

$h = -9 \rightarrow 9$

$T_{\text{min}} = 0.536, T_{\text{max}} = 1.000$

$k = -9 \rightarrow 4$

3166 measured reflections

$l = -12 \rightarrow 11$

Refinement

Refinement on F^2

Primary atom site location: structure-invariant direct
methods

Least-squares matrix: full

Secondary atom site location: difference Fourier map

$R[F^2 > 2\sigma(F^2)] = 0.028$

Hydrogen site location: difference Fourier map

$wR(F^2) = 0.072$

H atoms treated by a mixture of independent and
constrained refinement

$S = 1.15$

$w = 1/[\sigma^2(F_o^2) + (0.0315P)^2 + 1.1125P]$

where $P = (F_o^2 + 2F_c^2)/3$

2086 reflections	$(\Delta/\sigma)_{\max} < 0.001$
184 parameters	$\Delta\rho_{\max} = 0.41 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.79 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	0.08672 (3)	0.32297 (3)	0.90289 (3)	0.02898 (11)
O1	-0.0532 (4)	0.6070 (4)	0.8593 (3)	0.0337 (6)
O2	0.0672 (4)	0.8334 (4)	0.8475 (3)	0.0382 (6)
O3	0.1082 (4)	0.7617 (4)	0.1628 (3)	0.0347 (6)
O4	-0.1232 (4)	0.6718 (4)	0.3685 (3)	0.0405 (7)
O5	0.2389 (4)	0.0533 (4)	0.9029 (4)	0.0414 (7)
H5A	0.209 (10)	0.017 (10)	0.849 (9)	0.10 (2)*
H5B	0.211 (10)	-0.024 (10)	0.994 (9)	0.10 (3)*
N1	0.3422 (4)	0.4507 (4)	0.8114 (4)	0.0340 (7)
N2	0.5999 (5)	0.5524 (5)	0.6516 (4)	0.0453 (9)
H2	0.687 (9)	0.587 (9)	0.570 (8)	0.08 (2)*
C1	0.0393 (5)	0.7429 (5)	0.7840 (4)	0.0272 (7)
C2	0.1174 (5)	0.7962 (5)	0.6165 (4)	0.0255 (7)
C3	0.2660 (5)	0.8979 (5)	0.5366 (4)	0.0325 (8)
H3	0.3153	0.9372	0.5868	0.039*
C4	0.3406 (5)	0.9406 (6)	0.3814 (5)	0.0398 (9)
H4	0.4428	1.0048	0.3284	0.048*
C5	0.2642 (5)	0.8885 (5)	0.3047 (4)	0.0333 (8)
H5	0.3145	0.9184	0.2006	0.040*
C8	0.0251 (5)	0.7384 (5)	0.3016 (4)	0.0293 (8)
C7	0.0420 (4)	0.7439 (4)	0.5394 (4)	0.0248 (7)
H7	-0.0575	0.6757	0.5930	0.030*
C11	0.4652 (6)	0.4570 (6)	0.6844 (5)	0.0425 (10)
H11	0.4592	0.4010	0.6235	0.051*
C10	0.4032 (6)	0.5481 (6)	0.8641 (5)	0.0436 (10)
H10	0.3441	0.5688	0.9532	0.052*
C9	0.5632 (6)	0.6097 (7)	0.7663 (6)	0.0488 (11)
H9	0.6338	0.6779	0.7762	0.059*
C6	0.1121 (5)	0.7914 (5)	0.3837 (4)	0.0261 (7)

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.03179 (17)	0.03293 (17)	0.02429 (16)	-0.01047 (11)	-0.00754 (11)	-0.01069 (12)
O1	0.0381 (15)	0.0359 (15)	0.0243 (13)	-0.0070 (12)	-0.0089 (11)	-0.0074 (11)
O2	0.0528 (17)	0.0447 (16)	0.0260 (13)	-0.0095 (13)	-0.0144 (12)	-0.0175 (12)
O3	0.0442 (15)	0.0420 (15)	0.0236 (13)	-0.0069 (12)	-0.0114 (11)	-0.0154 (11)
O4	0.0422 (16)	0.0530 (18)	0.0302 (14)	-0.0195 (13)	-0.0118 (12)	-0.0125 (13)
O5	0.0440 (17)	0.0385 (16)	0.0405 (17)	-0.0083 (13)	-0.0105 (14)	-0.0137 (15)
N1	0.0310 (16)	0.0374 (18)	0.0328 (17)	-0.0080 (13)	-0.0085 (13)	-0.0114 (14)
N2	0.0336 (19)	0.055 (2)	0.039 (2)	-0.0110 (17)	-0.0046 (16)	-0.0120 (18)
C1	0.0274 (18)	0.0310 (19)	0.0236 (17)	-0.0004 (14)	-0.0103 (14)	-0.0091 (15)
C2	0.0289 (18)	0.0254 (17)	0.0223 (17)	-0.0011 (14)	-0.0100 (14)	-0.0073 (14)
C3	0.036 (2)	0.042 (2)	0.0276 (19)	-0.0122 (16)	-0.0133 (16)	-0.0137 (16)
C4	0.033 (2)	0.053 (3)	0.032 (2)	-0.0214 (18)	-0.0048 (16)	-0.0118 (18)
C5	0.035 (2)	0.042 (2)	0.0202 (17)	-0.0076 (16)	-0.0060 (15)	-0.0092 (16)
C8	0.038 (2)	0.0273 (18)	0.0267 (18)	-0.0028 (15)	-0.0152 (16)	-0.0089 (15)
C7	0.0250 (17)	0.0278 (17)	0.0220 (17)	-0.0045 (13)	-0.0084 (13)	-0.0075 (14)
C11	0.040 (2)	0.050 (3)	0.040 (2)	-0.0124 (19)	-0.0100 (18)	-0.017 (2)
C10	0.036 (2)	0.057 (3)	0.042 (2)	-0.0128 (19)	-0.0100 (18)	-0.021 (2)
C9	0.040 (2)	0.057 (3)	0.054 (3)	-0.017 (2)	-0.015 (2)	-0.020 (2)
C6	0.0285 (18)	0.0281 (18)	0.0248 (17)	-0.0013 (14)	-0.0113 (14)	-0.0105 (14)

Geometric parameters (\AA , $^\circ$)

Cd1—N1	2.216 (3)	N2—H2	0.85 (7)
Cd1—O3 ⁱ	2.251 (3)	C1—C2	1.492 (5)
Cd1—O2 ⁱⁱ	2.311 (3)	C2—C7	1.384 (5)
Cd1—O5	2.394 (3)	C2—C3	1.388 (5)
Cd1—O1	2.413 (3)	C3—C4	1.389 (5)
Cd1—O1 ⁱⁱ	2.626 (3)	C3—H3	0.9300
Cd1—O4 ⁱ	2.663 (3)	C4—C5	1.387 (6)
O1—C1	1.266 (5)	C4—H4	0.9300
O2—C1	1.260 (4)	C5—C6	1.391 (5)
O3—C8	1.276 (4)	C5—H5	0.9300
O4—C8	1.252 (5)	C8—C6	1.501 (5)
O5—H5A	0.84 (8)	C7—C6	1.387 (5)
O5—H5B	0.87 (8)	C7—H7	0.9300
N1—C11	1.306 (5)	C11—H11	0.9300
N1—C10	1.368 (5)	C10—C9	1.356 (6)
N2—C11	1.330 (6)	C10—H10	0.9300
N2—C9	1.351 (6)	C9—H9	0.9300
N1—Cd1—O3 ⁱ	143.34 (11)	C9—N2—H2	124 (4)
N1—Cd1—O2 ⁱⁱ	127.43 (11)	O2—C1—O1	121.5 (3)
O3 ⁱ —Cd1—O2 ⁱⁱ	88.49 (10)	O2—C1—C2	119.4 (3)
N1—Cd1—O5	88.63 (11)	O1—C1—C2	119.1 (3)

O3 ⁱ —Cd1—O5	87.33 (11)	C7—C2—C3	119.4 (3)
O2 ⁱⁱ —Cd1—O5	85.20 (11)	C7—C2—C1	120.0 (3)
N1—Cd1—O1	89.20 (11)	C3—C2—C1	120.5 (3)
O3 ⁱ —Cd1—O1	89.03 (10)	C2—C3—C4	119.7 (3)
O2 ⁱⁱ —Cd1—O1	103.25 (10)	C2—C3—H3	120.2
O5—Cd1—O1	170.71 (11)	C4—C3—H3	120.2
N1—Cd1—O1 ⁱⁱ	83.08 (10)	C5—C4—C3	120.6 (4)
O3 ⁱ —Cd1—O1 ⁱⁱ	131.61 (9)	C5—C4—H4	119.7
O2 ⁱⁱ —Cd1—O1 ⁱⁱ	52.58 (9)	C3—C4—H4	119.7
O5—Cd1—O1 ⁱⁱ	112.79 (11)	C4—C5—C6	119.8 (3)
O1—Cd1—O1 ⁱⁱ	75.89 (9)	C4—C5—H5	120.1
N1—Cd1—O4 ⁱ	91.00 (10)	C6—C5—H5	120.1
O3 ⁱ —Cd1—O4 ⁱ	52.54 (9)	O4—C8—O3	121.5 (3)
O2 ⁱⁱ —Cd1—O4 ⁱ	137.72 (9)	O4—C8—C6	120.7 (3)
O5—Cd1—O4 ⁱ	78.09 (11)	O3—C8—C6	117.8 (3)
O1—Cd1—O4 ⁱ	92.92 (9)	C2—C7—C6	121.2 (3)
O1 ⁱⁱ —Cd1—O4 ⁱ	167.35 (9)	C2—C7—H7	119.4
C1—O1—Cd1	119.5 (2)	C6—C7—H7	119.4
C1—O1—Cd1 ⁱⁱ	85.5 (2)	N1—C11—N2	112.7 (4)
Cd1—O1—Cd1 ⁱⁱ	104.11 (9)	N1—C11—H11	123.7
C1—O2—Cd1 ⁱⁱ	100.4 (2)	N2—C11—H11	123.7
C8—O3—Cd1 ⁱ	102.2 (2)	C9—C10—N1	109.4 (4)
Cd1—O5—H5A	108 (5)	C9—C10—H10	125.3
Cd1—O5—H5B	110 (5)	N1—C10—H10	125.3
H5A—O5—H5B	109 (7)	N2—C9—C10	106.5 (4)
C11—N1—C10	104.7 (3)	N2—C9—H9	126.7
C11—N1—Cd1	125.3 (3)	C10—C9—H9	126.7
C10—N1—Cd1	129.8 (3)	C7—C6—C5	119.1 (3)
C11—N2—C9	106.7 (4)	C7—C6—C8	120.4 (3)
C11—N2—H2	129 (4)	C5—C6—C8	120.4 (3)
N1—Cd1—O1—C1	9.8 (3)	O2—C1—C2—C7	-156.1 (3)
O3 ⁱ —Cd1—O1—C1	-133.6 (3)	O1—C1—C2—C7	24.1 (5)
O2 ⁱⁱ —Cd1—O1—C1	138.2 (3)	O2—C1—C2—C3	24.1 (5)
O1 ⁱⁱ —Cd1—O1—C1	92.8 (3)	O1—C1—C2—C3	-155.6 (4)
O4 ⁱ —Cd1—O1—C1	-81.2 (3)	C7—C2—C3—C4	-2.3 (6)
N1—Cd1—O1—Cd1 ⁱⁱ	-83.06 (11)	C1—C2—C3—C4	177.5 (4)
O3 ⁱ —Cd1—O1—Cd1 ⁱⁱ	133.56 (10)	C2—C3—C4—C5	2.6 (7)
O2 ⁱⁱ —Cd1—O1—Cd1 ⁱⁱ	45.34 (11)	C3—C4—C5—C6	-0.5 (7)
O1 ⁱⁱ —Cd1—O1—Cd1 ⁱⁱ	0.0	Cd1 ⁱ —O3—C8—O4	4.2 (4)
O4 ⁱ —Cd1—O1—Cd1 ⁱⁱ	-174.03 (9)	Cd1 ⁱ —O3—C8—C6	-176.6 (3)
O3 ⁱ —Cd1—N1—C11	-24.3 (5)	C3—C2—C7—C6	-0.1 (5)
O2 ⁱⁱ —Cd1—N1—C11	142.2 (3)	C1—C2—C7—C6	-179.9 (3)
O5—Cd1—N1—C11	59.3 (4)	C10—N1—C11—N2	-0.4 (5)

supplementary materials

O1—Cd1—N1—C11	−111.6 (4)	Cd1—N1—C11—N2	175.6 (3)
O1 ⁱⁱ —Cd1—N1—C11	172.5 (4)	C9—N2—C11—N1	1.0 (6)
O4 ⁱ —Cd1—N1—C11	−18.7 (4)	C11—N1—C10—C9	−0.4 (5)
O3 ⁱ —Cd1—N1—C10	150.6 (3)	Cd1—N1—C10—C9	−176.1 (3)
O2 ⁱⁱ —Cd1—N1—C10	−42.9 (4)	C11—N2—C9—C10	−1.2 (5)
O5—Cd1—N1—C10	−125.8 (4)	N1—C10—C9—N2	0.9 (6)
O1—Cd1—N1—C10	63.3 (4)	C2—C7—C6—C5	2.1 (5)
O1 ⁱⁱ —Cd1—N1—C10	−12.6 (4)	C2—C7—C6—C8	−178.0 (3)
O4 ⁱ —Cd1—N1—C10	156.2 (4)	C4—C5—C6—C7	−1.8 (6)
Cd1 ⁱⁱ —O2—C1—O1	0.1 (4)	C4—C5—C6—C8	178.3 (4)
Cd1 ⁱⁱ —O2—C1—C2	−179.6 (3)	O4—C8—C6—C7	9.7 (5)
Cd1—O1—C1—O2	−103.8 (4)	O3—C8—C6—C7	−169.4 (3)
Cd1 ⁱⁱ —O1—C1—O2	−0.1 (3)	O4—C8—C6—C5	−170.4 (4)
Cd1—O1—C1—C2	76.0 (4)	O3—C8—C6—C5	10.4 (5)
Cd1 ⁱⁱ —O1—C1—C2	179.6 (3)		


Symmetry codes: (i) $-x, -y+1, -z+1$; (ii) $-x, -y+1, -z+2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O5—H5A···O2 ⁱⁱⁱ	0.84 (8)	2.07 (8)	2.809 (4)	146 (7)
O5—H5B···O3 ^{iv}	0.87 (8)	1.96 (8)	2.786 (4)	158 (7)
N2—H2···O4 ^v	0.85 (7)	2.00 (7)	2.854 (5)	175 (6)

Symmetry codes: (iii) $x, y-1, z$; (iv) $x, y-1, z+1$; (v) $x+1, y, z$.

Fig. 1



supplementary materials

Fig. 2

