

# Chloridotris[ $\mu_2$ -2-(dimethylamino)-ethanolato]- $\mu_3$ -hydroxido-tri- $\mu_2$ -trifluoroacetato-tetracopper(II) tetrahydrofuran solvate

S. Tajammul Hussain,<sup>a\*</sup> Shahzad Abu Bakar,<sup>a</sup> Mohammad Mazhar<sup>b</sup> and Matthias Zeller<sup>c\*</sup>

<sup>a</sup>National Center for Physics, Quaid-i-Azam University, Islamabad 45320, Pakistan,

<sup>b</sup>Department of Chemistry, Faculty of Science, University of Malaya, Lembah Pantai, 50603-Kuala Lumpur, Malaysia, and <sup>c</sup>STaRBURSTT Cyber Instrumentation Consortium, Youngstown State University, Department of Chemistry, One University Plaza, Youngstown, Ohio 44555-3663, USA

Correspondence e-mail: dr\_tajammul@yahoo.ca, mzeller@ysu.edu


Received 10 May 2010; accepted 14 June 2010

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å; disorder in main residue;  $R$  factor = 0.043;  $wR$  factor = 0.103; data-to-parameter ratio = 22.0.

The title compound,  $[\text{Cu}_4(\text{C}_2\text{F}_3\text{O}_2)_3(\text{C}_4\text{H}_{10}\text{NO})_3\text{Cl}(\text{OH})] \cdot \text{C}_4\text{H}_8\text{O}$  or  $[\text{Cu}_4(\text{TFA})_3(\text{dmae})_3\text{Cl}(\text{OH})] \cdot \text{THF}$  (dmae is dimethylaminoethanolate, TFA is trifluoroacetate and THF is tetrahydrofuran), has an approximate molecular threefold symmetry with three equivalent  $\{\text{Cu}(\text{dmae})(\text{TFA})\}$  units bridging between a Cu—Cl and a hydroxide unit, with the latter two lying on the molecular threefold axis. However, in the solid state, the tetranuclear complex has  $C_i$  symmetry. The Cu atom bonded to the Cl atom has a distorted tetrahedral geometry. The other three Cu atoms have distorted square-pyramidal geometries with an  $\text{NO}_4$  coordination environment. The bonds within the  $\text{CuNO}_3$  base of the pyramid range from 1.953 (2) to 2.033 (3) Å, while the apical Cu—O bonds are significantly longer, ranging from 2.286 (2) to 2.377 (2) Å. The square-pyramidal geometries are augmented by weak interactions towards a sixth O atom, forming a highly distorted octahedral coordination environment [long Cu—O distances = 2.712 (2)–2.824 (2) Å]. The hydroxide group is hydrogen bonded to the tetrahydrofuran solvent molecule. One of the  $-\text{CF}_3$  groups shows minor disorder over two positions, with a refined occupancy ratio of 0.894 (4):0.106 (5).

## Related literature

For the synthesis of  $[\text{Cu}(\text{dmae})\text{Cl}]_4$ , used as starting material for title compound, see: Anwander *et al.* (1997). For general background to copper(II) complexes, see: Coastamagna *et al.* (1992). For related structures, see: Tahir *et al.* (2008); Shahid *et al.* (2009).



## Experimental

### Crystal data

$[\text{Cu}_4(\text{C}_2\text{F}_3\text{O}_2)_3(\text{C}_4\text{H}_{10}\text{NO})_3\text{Cl}(\text{OH})] \cdot \text{C}_4\text{H}_8\text{O}$   
 $M_r = 982.21$   
 Monoclinic,  $C2/c$   
 $a = 16.4353$  (14) Å  
 $b = 12.1893$  (12) Å  
 $c = 35.547$  (3) Å

$\beta = 94.678$  (2)°  
 $V = 7097.7$  (11) Å<sup>3</sup>  
 $Z = 8$   
 Mo  $K\alpha$  radiation  
 $\mu = 2.54$  mm<sup>-1</sup>  
 $T = 100$  K  
 $0.41 \times 0.38 \times 0.28$  mm

### Data collection

Bruker SMART APEX CCD diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 2008)  
 $T_{\text{min}} = 0.673$ ,  $T_{\text{max}} = 0.746$

20465 measured reflections  
 10267 independent reflections  
 7515 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.035$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$   
 $wR(F^2) = 0.103$   
 $S = 1.01$   
 10267 reflections  
 467 parameters

15 restraints  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.92$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.69$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{O10} \cdots \text{H10} \cdots \text{O11}$	1.00	1.73	2.723 (3)	174

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: SHELXTL and publCIF (Westrip, 2010).

The authors are grateful to the Higher Education Commission of Pakistan for financial support. The diffractometer was funded by NSF grant 0087210, by Ohio Board of Regents grant CAP-491, and by YSU.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2302).

## References

- Anwander, R., Munck, F. C., Priermeier, T., Scherer, W., Runte, O. & Herrmann, W. A. (1997). *Inorg. Chem.* **36**, 3545–3552.
- Bruker (2008). *APEX2, SAINT and SADABS*. Bruker AXS Inc, Madison, Wisconsin, USA.
- Coastamagna, J., Vargas, J., Latorre, R. & Alvarado, A. (1992). *Coord. Chem. Rev.* **119**, 67–88.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Shahid, M., Mazhar, M., Zeller, M. & Hunter, A. D. (2009). *Acta Cryst.* **E65**, m345–m346.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Tahir, A. A., Hamid, M., Mazhar, M., Zeller, M., Hunter, A. D., Nadeem, M. & Akhtar, M. J. (2008). *Dalton Trans.* pp. 1224–1232.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**. Submitted.

**supplementary materials**

*Acta Cryst.* (2010). E66, m814-m815 [ doi:10.1107/S1600536810022877 ]

## Chloridotris[ $\mu_2$ -2-(dimethylamino)ethanolato]- $\mu_3$ -hydroxido-tri- $\mu_2$ -trifluoroacetato-tetracopper(II) tetrahydrofuran solvate

S. T. Hussain, S. A. Bakar, M. Mazhar and M. Zeller

### Comment

In recent years, there has been a considerable interest towards the synthesis of copper complexes; these complexes are extensively used in catalysis, enzymatic reactions, magnetism and molecular architecture (Coastamagna, Vargas *et al.*, 1992). The present work is a continuation of earlier studies for the preparation and structure elucidation of copper (II) complexes (Shahid *et al.*, 2009). The motivation behind the synthesis of the title compound was to use it as a starting material for the synthesis of single source precursors for the deposition of thin films of copper oxides using aerosol assisted chemical vapor deposition (AACVD). We present here the synthesis and crystal structure of the title compound,  $[\text{Cu}_4((\text{CH}_3)_2\text{NCH}_2\text{CH}_2\text{O})_3(\text{F}_3\text{CCOO})_3(\text{OH})\text{Cl}]$  or  $[\text{Cu}_4(\text{dmae})_3(\text{TFA})_3(\text{OH})\text{Cl}]$  (dmae = dimethylaminoethanolate, TFA = trifluoroacetate), which crystallized from THF as the mono solvate with the THF molecule tightly hydrogen bonded to the hydroxyl group.

The title compound has a slightly distorted molecular three fold symmetry with three equivalent  $\text{Cu}(\text{dmae})(\text{TFA})$  units bridging *via* their alcoholate oxygen atoms between a  $\text{Cu}-\text{Cl}$  and an hydroxyl unit with the latter two lying on a molecular pseudo threefold axis. The Cu atom bonded to the chlorine has a distorted tetrahedral geometry. The other three copper atoms have distorted square pyramidal geometries with a  $\text{CNO}_4$  coordination environment from the dmae O and N atoms, the hydroxyl O atom and two TFA anions. The TFA anions are bridging between two neighboring copper ions with one of the oxygen atoms being part of the base of the pyramid of one copper ion, and the other being in the apical position of the neighboring copper ion. The bonds within the  $\text{CuNO}_3$  bases of the pyramids are strong and quite similar in length with distances between 1.953 (2) and 2.033 (3) Å. The apical  $\text{Cu}-\text{O}$  bonds are significantly longer and between 2.286 (2) and 2.377 (2) Å, thus rendering the  $\mu_2$ -bridge of the TFA ions asymmetric. The square pyramidal geometries are augmented by weak interactions towards a fifth oxygen atom to form a highly distorted octahedral coordination environment ( $\text{Cu}-\text{O}$  distances:  $\text{O}_3-\text{Cu}_3 = 2.712$  (2),  $\text{O}_2-\text{Cu}_2 = 2.780$  (2),  $\text{O}_1-\text{Cu}_4 = 2.8240$  (2) Å).

A similar motif as in the title compound was previously observed for two mixed metal copper-titanium complexes (Tahir *et al.*, 2008). In these complexes the TFA anions were replaced by benzoate or 2-methyl-benzoate ligands, and the  $\text{Cu}-\text{Cl}$  unit was replaced by a titanium atom, which in turn was bonded to another larger  $\text{Cu}-\text{Ti}$  cluster. The  $[\text{Cu}_3(\text{dmae})_3(\text{TFA})_3(\text{OH})]$  unit in the title compound and the  $[\text{Cu}_3((\text{CH}_3)_2\text{NCH}_2\text{CH}_2\text{O})_3(\text{O}_2\text{C}-\text{C}_6\text{H}_5\text{R})_3(\text{OH})]$  units in the  $\text{Cu}-\text{Ti}$  complexes ( $R = \text{H}, \text{Me}$ ) are quite similar. In the 2-methyl-benzoate complexes the  $[\text{Cu}_3((\text{CH}_3)_2\text{NCH}_2\text{CH}_2\text{O})_3(\text{O}_2\text{C}-\text{C}_6\text{H}_5\text{Me})_3(\text{OH})]$  unit is located on an actual crystallographic three fold axis. The carboxylate anions show coordination modes differing slightly from those observed in the title compound with some of the oxygen atoms being detached from the copper ions and interactions to the fifth oxygen atom, which are very weak in the title compound, being strengthened instead. The overall coordination environment - distorted square pyramidal  $\text{CNO}_4$  geometries with an additional weak interaction towards a fifth oxygen atom - is however the same in all three compounds, which shows the idiosyncrasy commonly observed for copper(II) to form strongly distorted and highly flexible octahedral geometries with a set of four strong bonds in a square planar arrangement and two apical ligands at variable distances. Indi-

## supplementary materials

---

vidual ligand atoms in these kinds of complexes can easily switch from tightly bound to only weakly coordinated as long as the overall coordination environment of the metal center is retained, and energy differences and activation barriers between the different arrangements that can be achieved that way are quite small. The difference in bonding arrangement in the three complexes in the solid state does thus probably not translate into a different chemical nature for the three complexes as the bonding environment around Cu(II) is very flexible and it can be assumed that in solution (*i.e.* upon release of packing effects) all complexes will attain the same connectivity pattern.

In the title compound the hydroxyl group is O—H···O hydrogen bonded to a tetrahydrofuran molecule (Table 1), which is embedded in a bowl shaped cleft of the complex formed by the three TFA ligands. No such host–guest behavior was observed for the other two related compounds (Tahir *et al.*, 2008).

### Experimental

Tetrameric *N,N*-dimethylaminoethanolato copper(II) chloride, [Cu(dmae)Cl]<sub>4</sub> was prepared according to a literature method (Anwander *et al.*, 1997). The title compound was prepared as follows: 1.25 g (1.67 mmole) of [Cu(dmae)Cl]<sub>4</sub> in 20 ml THF were combined with 1.77 g (6.66 mmole) of Cu(F<sub>3</sub>CCOO)<sub>2</sub> in 10 ml THF followed by the addition of 0.297 g (3.33 mmole) *N,N*-dimethylaminoethanol. The reaction mixture was stirred for 3 h and filtered through a cannula to remove any undissolved species. The filtrate was evaporated to dryness under vacuum, the solid was re-dissolved in 5 ml THF and placed in a vial with rubber seal at room temperature for one week to give blue crystals suitable for single-crystal X-ray diffraction analysis. Yield: 86% m.p. 393–394 K. Elemental Analysis for Cu<sub>4</sub>((CH<sub>3</sub>)<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>O)<sub>3</sub>(F<sub>3</sub>CCOO)<sub>3</sub>(OH)Cl % calc: C, 21.99 H, 3.97 N, 4.27, % found: C, 22.10 H, 3.90 N, 4.53.

### Refinement

The fluorine atoms bonded to C14 were refined as disordered over two mutually exclusive positions with a refined occupancy ratio of 0.894 (4) to 0.106 (5). C—F bond distances within this CF<sub>3</sub> group were restrained to be the same within a standard uncertainty of 0.02 Å and ADPs of the minor F atoms were constrained to be identical to those of the major moiety F atom opposite their position.

All hydrogen atoms were added in calculated positions with a C—H bond distances of 0.97 (methylene), 0.96 (methyl) and 1.00 Å (OH). They were refined with isotropic displacement parameters  $U_{\text{iso}}$  of 1.5 (methyl, OH) or 1.2 times  $U_{\text{eq}}$  (methylene) of the adjacent carbon or oxygen atom.

### Figures




Fig. 1. Perspective view of the title compound with the atom numbering scheme. The displacement ellipsoids are at the 50% probability level and H atoms are drawn as small spheres of arbitrary radii.




Fig. 2. Perspective view of the title compound, view down the pseudo three fold axis. The displacement ellipsoids are at the 50% probability level and H atoms are drawn as small spheres of arbitrary radii.

**Chloridotris[ $\mu_2$ -2-(dimethylamino)ethanolato]- $\mu_3$ -hydroxido-tri- $\mu_2$ - trifluoroacetato-tetracopper(II) tetrahydrofuran solvate**

*Crystal data*

[Cu<sub>4</sub>(C<sub>2</sub>F<sub>3</sub>O<sub>2</sub>)<sub>3</sub>(C<sub>4</sub>H<sub>10</sub>NO)<sub>3</sub>Cl(OH)]·C<sub>4</sub>H<sub>8</sub>O

$M_r = 982.21$

Monoclinic, *C2/c*

Hall symbol: -C 2yc

$a = 16.4353$  (14) Å

$b = 12.1893$  (12) Å

$c = 35.547$  (3) Å

$\beta = 94.678$  (2)°

$V = 7097.7$  (11) Å<sup>3</sup>

$Z = 8$

$F(000) = 3952$

$D_x = 1.838$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 1722 reflections

$\theta = 2.4$ – $30.1$ °

$\mu = 2.54$  mm<sup>-1</sup>

$T = 100$  K

Block, blue

$0.41 \times 0.38 \times 0.28$  mm

*Data collection*

Bruker SMART APEX CCD  
diffractometer

Radiation source: fine-focus sealed tube  
graphite

$\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Bruker, 2008)

$T_{\min} = 0.673$ ,  $T_{\max} = 0.746$

20465 measured reflections

10267 independent reflections

7515 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.035$

$\theta_{\text{max}} = 31.6$ °,  $\theta_{\text{min}} = 1.2$ °

$h = -12$ → $22$

$k = -15$ → $17$

$l = -37$ → $50$

*Refinement*

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.043$

$wR(F^2) = 0.103$

$S = 1.01$

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0396P)^2 + 12.2423P]$

where  $P = (F_o^2 + 2F_c^2)/3$

## supplementary materials

---

10267 reflections	$(\Delta/\sigma)_{\max} = 0.001$
467 parameters	$\Delta\rho_{\max} = 0.92 \text{ e } \text{\AA}^{-3}$
15 restraints	$\Delta\rho_{\min} = -0.69 \text{ e } \text{\AA}^{-3}$

### Special details

**Experimental.** The fluorine atoms bonded to C14 were refined as disordered over two mutually exclusive positions with a refined occupancy ratio of 0.894 (4) to 0.106 (5). C-F bond distances within this CF<sub>3</sub> group were restrained to be the same within a standard uncertainty of 0.02 Angstrom and ADPs of the minor F atoms were constrained to be identical to those of the major moiety F atom opposite their position.

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.3487 (2)	0.6333 (3)	0.27202 (8)	0.0213 (7)	
H1A	0.2955	0.6699	0.2659	0.026*	
H1B	0.3614	0.5884	0.2500	0.026*	
C2	0.41488 (18)	0.7186 (3)	0.28039 (9)	0.0188 (7)	
H2A	0.4693	0.6841	0.2796	0.023*	
H2B	0.4097	0.7771	0.2610	0.023*	
C3	0.3331 (2)	0.8365 (3)	0.31874 (10)	0.0253 (8)	
H3A	0.3358	0.8958	0.3003	0.038*	
H3B	0.3303	0.8679	0.3440	0.038*	
H3C	0.2843	0.7919	0.3123	0.038*	
C4	0.4799 (2)	0.8341 (3)	0.32982 (10)	0.0240 (7)	
H4A	0.4845	0.8938	0.3117	0.036*	
H4B	0.5289	0.7881	0.3305	0.036*	
H4C	0.4745	0.8649	0.3550	0.036*	
C5	0.16045 (19)	0.6242 (3)	0.37865 (9)	0.0203 (7)	
H5A	0.1270	0.6612	0.3580	0.024*	
H5B	0.1718	0.6772	0.3995	0.024*	
C6	0.11552 (19)	0.5262 (3)	0.39231 (10)	0.0234 (7)	
H6A	0.0666	0.5508	0.4045	0.028*	
H6B	0.0974	0.4787	0.3706	0.028*	
C7	0.1697 (2)	0.5093 (3)	0.45818 (10)	0.0283 (8)	
H7A	0.1147	0.5027	0.4668	0.042*	
H7B	0.1853	0.5868	0.4578	0.042*	
H7C	0.2085	0.4690	0.4754	0.042*	
C8	0.1435 (2)	0.3471 (3)	0.42013 (11)	0.0274 (8)	

H8A	0.0861	0.3437	0.4256	0.041*	
H8B	0.1769	0.3067	0.4396	0.041*	
H8C	0.1498	0.3141	0.3954	0.041*	
C9	0.24516 (18)	0.2700 (3)	0.31588 (9)	0.0180 (6)	
H9A	0.2240	0.2915	0.2901	0.022*	
H9B	0.2000	0.2358	0.3285	0.022*	
C10	0.31478 (18)	0.1882 (3)	0.31392 (9)	0.0175 (6)	
H10A	0.3235	0.1482	0.3381	0.021*	
H10B	0.3008	0.1341	0.2937	0.021*	
C11	0.3849 (2)	0.2949 (3)	0.26740 (9)	0.0238 (7)	
H11A	0.3777	0.2355	0.2489	0.036*	
H11B	0.4350	0.3355	0.2634	0.036*	
H11C	0.3380	0.3448	0.2644	0.036*	
C12	0.46108 (19)	0.1720 (3)	0.31080 (10)	0.0237 (7)	
H12A	0.4524	0.1108	0.2930	0.036*	
H12B	0.4661	0.1437	0.3367	0.036*	
H12C	0.5112	0.2110	0.3058	0.036*	
C13	0.37061 (18)	0.7122 (3)	0.42640 (9)	0.0166 (6)	
C14	0.3775 (2)	0.8151 (3)	0.45127 (10)	0.0294 (8)	
C15	0.36846 (19)	0.2779 (3)	0.42436 (9)	0.0194 (6)	
C16	0.3779 (2)	0.1874 (3)	0.45482 (10)	0.0252 (7)	
C17	0.55106 (18)	0.4851 (3)	0.34328 (8)	0.0157 (6)	
C18	0.6447 (2)	0.4702 (3)	0.34677 (10)	0.0244 (7)	
C19	0.5583 (2)	0.3809 (3)	0.44361 (11)	0.0290 (8)	
H19A	0.5641	0.3255	0.4236	0.035*	
H19B	0.5265	0.3485	0.4633	0.035*	
C20	0.6412 (2)	0.4186 (4)	0.46031 (11)	0.0331 (9)	
H20A	0.6634	0.3684	0.4805	0.040*	
H20B	0.6805	0.4245	0.4407	0.040*	
C21	0.6219 (2)	0.5295 (4)	0.47598 (13)	0.0426 (11)	
H21A	0.6701	0.5785	0.4767	0.051*	
H21B	0.6036	0.5231	0.5017	0.051*	
C22	0.5532 (2)	0.5721 (3)	0.44827 (11)	0.0330 (9)	
H22A	0.5109	0.6091	0.4620	0.040*	
H22B	0.5747	0.6252	0.4305	0.040*	
Cl1	0.13236 (5)	0.50928 (7)	0.27516 (2)	0.02051 (16)	
Cu1	0.23798 (2)	0.50916 (3)	0.317137 (10)	0.01044 (8)	
Cu2	0.39464 (2)	0.63387 (3)	0.350823 (10)	0.01304 (8)	
Cu3	0.28392 (2)	0.48008 (3)	0.401847 (10)	0.01366 (9)	
Cu4	0.39472 (2)	0.37149 (3)	0.343178 (10)	0.01343 (8)	
F1	0.38789 (19)	0.2286 (2)	0.48959 (7)	0.0616 (8)	
F2	0.44062 (16)	0.1223 (2)	0.45106 (7)	0.0499 (7)	
F3	0.31145 (16)	0.1262 (2)	0.45306 (9)	0.0618 (8)	
F4	0.45195 (17)	0.8352 (4)	0.46438 (13)	0.0738 (15)	0.894 (5)
F5	0.3505 (3)	0.9028 (2)	0.43088 (9)	0.0678 (12)	0.894 (5)
F6	0.3305 (2)	0.8121 (3)	0.47923 (9)	0.0526 (10)	0.894 (5)
F4B	0.418 (2)	0.8966 (17)	0.4377 (8)	0.0526 (10)	0.106 (5)
F5B	0.3162 (13)	0.855 (3)	0.4664 (11)	0.0738 (15)	0.106 (5)
F6B	0.420 (2)	0.788 (2)	0.4834 (6)	0.0678 (12)	0.106 (5)



## supplementary materials

---

F7	0.68329 (13)	0.5493 (2)	0.36747 (8)	0.0452 (7)
F8	0.67280 (13)	0.4751 (2)	0.31263 (7)	0.0474 (7)
F9	0.66948 (11)	0.37600 (18)	0.36251 (6)	0.0329 (5)
N1	0.40701 (15)	0.7666 (2)	0.31829 (7)	0.0158 (5)
N2	0.17040 (16)	0.4632 (2)	0.41990 (7)	0.0195 (6)
N3	0.39102 (15)	0.2481 (2)	0.30606 (7)	0.0163 (5)
O1	0.34445 (12)	0.56503 (18)	0.30425 (6)	0.0159 (4)
O2	0.23492 (12)	0.58621 (18)	0.36547 (6)	0.0155 (4)
O3	0.27429 (12)	0.36447 (17)	0.33646 (6)	0.0141 (4)
O4	0.40721 (13)	0.72338 (18)	0.39660 (6)	0.0187 (5)
O5	0.33254 (13)	0.6340 (2)	0.43810 (6)	0.0211 (5)
O6	0.32578 (13)	0.35798 (19)	0.43373 (6)	0.0192 (5)
O7	0.40130 (13)	0.25998 (19)	0.39509 (6)	0.0189 (5)
O8	0.51284 (12)	0.39551 (19)	0.34027 (6)	0.0196 (5)
O9	0.52741 (13)	0.58004 (19)	0.34305 (7)	0.0213 (5)
O10	0.38614 (12)	0.49500 (16)	0.37767 (6)	0.0122 (4)
H10	0.4325	0.4905	0.3976	0.018*
O11	0.51928 (15)	0.4780 (2)	0.42819 (8)	0.0318 (6)

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0238 (17)	0.0301 (18)	0.0103 (14)	-0.0031 (14)	0.0035 (12)	0.0015 (13)
C2	0.0169 (15)	0.0227 (17)	0.0180 (15)	0.0011 (12)	0.0087 (12)	0.0044 (13)
C3	0.0224 (17)	0.0220 (18)	0.0321 (19)	0.0076 (14)	0.0060 (14)	0.0069 (15)
C4	0.0229 (17)	0.0194 (17)	0.0296 (18)	-0.0061 (13)	0.0021 (14)	0.0040 (14)
C5	0.0201 (16)	0.0214 (17)	0.0199 (15)	0.0065 (13)	0.0046 (12)	0.0013 (13)
C6	0.0168 (15)	0.0282 (19)	0.0250 (17)	0.0009 (13)	0.0007 (13)	-0.0003 (14)
C7	0.0271 (17)	0.040 (2)	0.0189 (17)	-0.0023 (16)	0.0088 (14)	-0.0038 (15)
C8	0.0244 (18)	0.028 (2)	0.0312 (19)	-0.0064 (14)	0.0074 (15)	0.0034 (15)
C9	0.0167 (15)	0.0161 (15)	0.0212 (16)	-0.0021 (12)	0.0008 (12)	-0.0019 (13)
C10	0.0178 (15)	0.0175 (16)	0.0171 (15)	-0.0017 (12)	0.0009 (12)	-0.0018 (12)
C11	0.0274 (18)	0.0251 (18)	0.0197 (16)	-0.0027 (14)	0.0075 (14)	-0.0045 (14)
C12	0.0168 (16)	0.0243 (18)	0.0297 (18)	0.0062 (13)	0.0010 (13)	-0.0089 (15)
C13	0.0161 (15)	0.0173 (15)	0.0156 (15)	0.0034 (12)	-0.0033 (12)	-0.0047 (12)
C14	0.036 (2)	0.0260 (19)	0.0273 (19)	-0.0043 (16)	0.0099 (16)	-0.0086 (16)
C15	0.0195 (16)	0.0184 (16)	0.0197 (16)	-0.0030 (13)	-0.0031 (12)	0.0024 (13)
C16	0.035 (2)	0.0217 (18)	0.0199 (17)	0.0028 (15)	0.0053 (14)	0.0050 (14)
C17	0.0125 (13)	0.0211 (16)	0.0139 (14)	0.0018 (12)	0.0036 (11)	0.0025 (12)
C18	0.0166 (15)	0.0246 (18)	0.0324 (19)	0.0038 (13)	0.0054 (14)	0.0113 (15)
C19	0.0284 (19)	0.029 (2)	0.0283 (19)	0.0025 (15)	-0.0032 (15)	0.0047 (16)
C20	0.0233 (18)	0.048 (3)	0.0270 (19)	0.0037 (17)	-0.0060 (15)	0.0058 (18)
C21	0.032 (2)	0.049 (3)	0.044 (3)	-0.0038 (19)	-0.0155 (19)	-0.007 (2)
C22	0.029 (2)	0.030 (2)	0.037 (2)	0.0000 (16)	-0.0118 (16)	-0.0089 (17)
Cl1	0.0180 (3)	0.0238 (4)	0.0185 (4)	0.0001 (3)	-0.0054 (3)	0.0021 (3)
Cu1	0.01002 (16)	0.01276 (18)	0.00839 (16)	-0.00009 (13)	-0.00024 (12)	0.00033 (13)
Cu2	0.01405 (18)	0.01354 (18)	0.01150 (17)	-0.00088 (14)	0.00094 (13)	0.00082 (14)
Cu3	0.01306 (17)	0.01724 (19)	0.01080 (17)	0.00010 (14)	0.00167 (13)	0.00157 (14)

Cu4	0.01185 (18)	0.01471 (18)	0.01367 (17)	0.00029 (14)	0.00071 (13)	-0.00261 (14)
F1	0.115 (2)	0.0484 (17)	0.0214 (13)	0.0182 (16)	0.0036 (13)	0.0092 (12)
F2	0.0589 (17)	0.0486 (16)	0.0429 (15)	0.0237 (13)	0.0085 (12)	0.0191 (13)
F3	0.0555 (17)	0.0442 (17)	0.086 (2)	-0.0120 (13)	0.0071 (15)	0.0327 (16)
F4	0.0287 (16)	0.099 (3)	0.093 (3)	-0.0140 (17)	0.0005 (16)	-0.075 (3)
F5	0.121 (4)	0.0276 (16)	0.056 (2)	0.0143 (18)	0.012 (2)	-0.0113 (15)
F6	0.061 (2)	0.056 (2)	0.0442 (19)	-0.0053 (16)	0.0264 (16)	-0.0278 (16)
F4B	0.061 (2)	0.056 (2)	0.0442 (19)	-0.0053 (16)	0.0264 (16)	-0.0278 (16)
F5B	0.0287 (16)	0.099 (3)	0.093 (3)	-0.0140 (17)	0.0005 (16)	-0.075 (3)
F6B	0.121 (4)	0.0276 (16)	0.056 (2)	0.0143 (18)	0.012 (2)	-0.0113 (15)
F7	0.0210 (11)	0.0334 (13)	0.0787 (19)	-0.0063 (10)	-0.0113 (11)	0.0017 (13)
F8	0.0272 (12)	0.075 (2)	0.0427 (15)	0.0169 (12)	0.0203 (10)	0.0258 (13)
F9	0.0180 (10)	0.0303 (12)	0.0505 (14)	0.0051 (8)	0.0039 (9)	0.0172 (11)
N1	0.0137 (12)	0.0168 (13)	0.0173 (13)	0.0012 (10)	0.0034 (10)	0.0015 (10)
N2	0.0181 (13)	0.0260 (15)	0.0148 (13)	-0.0005 (11)	0.0039 (10)	0.0009 (11)
N3	0.0154 (12)	0.0155 (13)	0.0180 (13)	0.0011 (10)	0.0011 (10)	-0.0029 (11)
O1	0.0160 (11)	0.0204 (11)	0.0114 (10)	-0.0034 (9)	0.0026 (8)	0.0016 (9)
O2	0.0151 (10)	0.0186 (11)	0.0128 (10)	0.0018 (9)	0.0018 (8)	0.0002 (9)
O3	0.0128 (10)	0.0139 (10)	0.0153 (10)	0.0000 (8)	-0.0006 (8)	-0.0019 (8)
O4	0.0204 (11)	0.0191 (12)	0.0168 (11)	-0.0024 (9)	0.0022 (9)	-0.0016 (9)
O5	0.0227 (12)	0.0260 (13)	0.0147 (11)	-0.0046 (10)	0.0018 (9)	-0.0039 (10)
O6	0.0211 (12)	0.0200 (12)	0.0165 (11)	0.0013 (9)	0.0021 (9)	0.0037 (9)
O7	0.0222 (11)	0.0179 (11)	0.0167 (11)	0.0023 (9)	0.0020 (9)	0.0009 (9)
O8	0.0138 (11)	0.0213 (12)	0.0242 (12)	0.0000 (9)	0.0048 (9)	-0.0032 (10)
O9	0.0155 (11)	0.0201 (12)	0.0285 (13)	0.0027 (9)	0.0030 (9)	0.0045 (10)
O10	0.0122 (9)	0.0135 (10)	0.0109 (9)	-0.0005 (8)	0.0002 (7)	-0.0007 (8)
O11	0.0296 (14)	0.0249 (14)	0.0373 (15)	0.0007 (11)	-0.0193 (11)	-0.0005 (12)

*Geometric parameters (Å, °)*

C1—O1	1.422 (4)	C14—F6	1.308 (4)
C1—C2	1.516 (5)	C14—F4B	1.310 (14)
C1—H1A	0.9900	C14—F6B	1.333 (15)
C1—H1B	0.9900	C14—F5	1.346 (5)
C2—N1	1.484 (4)	C15—O7	1.230 (4)
C2—H2A	0.9900	C15—O6	1.263 (4)
C2—H2B	0.9900	C15—C16	1.545 (5)
C3—N1	1.485 (4)	C16—F2	1.316 (4)
C3—H3A	0.9800	C16—F3	1.319 (4)
C3—H3B	0.9800	C16—F1	1.332 (4)
C3—H3C	0.9800	C17—O9	1.221 (4)
C4—N1	1.484 (4)	C17—O8	1.260 (4)
C4—H4A	0.9800	C17—C18	1.545 (4)
C4—H4B	0.9800	C18—F9	1.326 (4)
C4—H4C	0.9800	C18—F8	1.334 (4)
C5—O2	1.423 (3)	C18—F7	1.340 (4)
C5—C6	1.505 (5)	C19—O11	1.434 (4)
C5—H5A	0.9900	C19—C20	1.512 (5)
C5—H5B	0.9900	C19—H19A	0.9900

## supplementary materials

---

C6—N2	1.490 (4)	C19—H19B	0.9900
C6—H6A	0.9900	C20—C21	1.506 (6)
C6—H6B	0.9900	C20—H20A	0.9900
C7—N2	1.473 (4)	C20—H20B	0.9900
C7—H7A	0.9800	C21—C22	1.528 (5)
C7—H7B	0.9800	C21—H21A	0.9900
C7—H7C	0.9800	C21—H21B	0.9900
C8—N2	1.483 (4)	C22—O11	1.439 (4)
C8—H8A	0.9800	C22—H22A	0.9900
C8—H8B	0.9800	C22—H22B	0.9900
C8—H8C	0.9800	C11—Cu1	2.1948 (8)
C9—O3	1.426 (4)	Cu1—O2	1.962 (2)
C9—C10	1.523 (4)	Cu1—O1	1.966 (2)
C9—H9A	0.9900	Cu1—O3	1.968 (2)
C9—H9B	0.9900	Cu2—O10	1.954 (2)
C10—N3	1.496 (4)	Cu2—O4	1.956 (2)
C10—H10A	0.9900	Cu2—O1	1.975 (2)
C10—H10B	0.9900	Cu2—N1	2.009 (3)
C11—N3	1.484 (4)	Cu2—O9	2.317 (2)
C11—H11A	0.9800	Cu3—O2	1.956 (2)
C11—H11B	0.9800	Cu3—O10	1.957 (2)
C11—H11C	0.9800	Cu3—O6	1.961 (2)
C12—N3	1.478 (4)	Cu3—N2	2.032 (3)
C12—H12A	0.9800	Cu3—O5	2.377 (2)
C12—H12B	0.9800	Cu4—O10	1.954 (2)
C12—H12C	0.9800	Cu4—O8	1.974 (2)
C13—O5	1.231 (4)	Cu4—O3	1.976 (2)
C13—O4	1.267 (4)	Cu4—N3	1.998 (3)
C13—C14	1.533 (5)	Cu4—O7	2.287 (2)
C14—F5B	1.278 (15)	O10—H10	0.9990
C14—F4	1.296 (4)		
O1—C1—C2	109.0 (2)	F8—C18—C17	109.7 (3)
O1—C1—H1A	109.9	F7—C18—C17	112.5 (3)
C2—C1—H1A	109.9	O11—C19—C20	105.1 (3)
O1—C1—H1B	109.9	O11—C19—H19A	110.7
C2—C1—H1B	109.9	C20—C19—H19A	110.7
H1A—C1—H1B	108.3	O11—C19—H19B	110.7
N1—C2—C1	109.5 (2)	C20—C19—H19B	110.7
N1—C2—H2A	109.8	H19A—C19—H19B	108.8
C1—C2—H2A	109.8	C21—C20—C19	102.0 (3)
N1—C2—H2B	109.8	C21—C20—H20A	111.4
C1—C2—H2B	109.8	C19—C20—H20A	111.4
H2A—C2—H2B	108.2	C21—C20—H20B	111.4
N1—C3—H3A	109.5	C19—C20—H20B	111.4
N1—C3—H3B	109.5	H20A—C20—H20B	109.2
H3A—C3—H3B	109.5	C20—C21—C22	103.5 (3)
N1—C3—H3C	109.5	C20—C21—H21A	111.1
H3A—C3—H3C	109.5	C22—C21—H21A	111.1
H3B—C3—H3C	109.5	C20—C21—H21B	111.1

N1—C4—H4A	109.5	C22—C21—H21B	111.1
N1—C4—H4B	109.5	H21A—C21—H21B	109.0
H4A—C4—H4B	109.5	O11—C22—C21	106.5 (3)
N1—C4—H4C	109.5	O11—C22—H22A	110.4
H4A—C4—H4C	109.5	C21—C22—H22A	110.4
H4B—C4—H4C	109.5	O11—C22—H22B	110.4
O2—C5—C6	107.8 (3)	C21—C22—H22B	110.4
O2—C5—H5A	110.1	H22A—C22—H22B	108.6
C6—C5—H5A	110.1	O2—Cu1—O1	97.18 (9)
O2—C5—H5B	110.1	O2—Cu1—O3	98.75 (9)
C6—C5—H5B	110.1	O1—Cu1—O3	98.13 (9)
H5A—C5—H5B	108.5	O2—Cu1—Cl1	121.31 (6)
N2—C6—C5	109.6 (3)	O1—Cu1—Cl1	120.73 (7)
N2—C6—H6A	109.8	O3—Cu1—Cl1	116.00 (6)
C5—C6—H6A	109.8	O10—Cu2—O4	94.81 (9)
N2—C6—H6B	109.8	O10—Cu2—O1	89.98 (9)
C5—C6—H6B	109.8	O4—Cu2—O1	160.49 (9)
H6A—C6—H6B	108.2	O10—Cu2—N1	173.56 (9)
N2—C7—H7A	109.5	O4—Cu2—N1	91.20 (10)
N2—C7—H7B	109.5	O1—Cu2—N1	85.10 (10)
H7A—C7—H7B	109.5	O10—Cu2—O9	85.32 (8)
N2—C7—H7C	109.5	O4—Cu2—O9	102.75 (9)
H7A—C7—H7C	109.5	O1—Cu2—O9	96.48 (8)
H7B—C7—H7C	109.5	N1—Cu2—O9	91.08 (9)
N2—C8—H8A	109.5	O2—Cu3—O10	88.29 (8)
N2—C8—H8B	109.5	O2—Cu3—O6	172.03 (9)
H8A—C8—H8B	109.5	O10—Cu3—O6	92.93 (9)
N2—C8—H8C	109.5	O2—Cu3—N2	86.38 (10)
H8A—C8—H8C	109.5	O10—Cu3—N2	172.39 (10)
H8B—C8—H8C	109.5	O6—Cu3—N2	91.62 (10)
O3—C9—C10	109.3 (2)	O2—Cu3—O5	86.47 (9)
O3—C9—H9A	109.8	O10—Cu3—O5	84.38 (8)
C10—C9—H9A	109.8	O6—Cu3—O5	101.48 (9)
O3—C9—H9B	109.8	N2—Cu3—O5	100.69 (10)
C10—C9—H9B	109.8	O10—Cu4—O8	92.30 (9)
H9A—C9—H9B	108.3	O10—Cu4—O3	89.22 (8)
N3—C10—C9	109.5 (3)	O8—Cu4—O3	168.40 (9)
N3—C10—H10A	109.8	O10—Cu4—N3	173.88 (9)
C9—C10—H10A	109.8	O8—Cu4—N3	93.10 (10)
N3—C10—H10B	109.8	O3—Cu4—N3	84.91 (9)
C9—C10—H10B	109.8	O10—Cu4—O7	87.29 (8)
H10A—C10—H10B	108.2	O8—Cu4—O7	98.53 (9)
N3—C11—H11A	109.5	O3—Cu4—O7	93.03 (8)
N3—C11—H11B	109.5	N3—Cu4—O7	94.73 (9)
H11A—C11—H11B	109.5	C4—N1—C2	110.0 (2)
N3—C11—H11C	109.5	C4—N1—C3	108.8 (3)
H11A—C11—H11C	109.5	C2—N1—C3	111.6 (2)
H11B—C11—H11C	109.5	C4—N1—Cu2	113.88 (19)
N3—C12—H12A	109.5	C2—N1—Cu2	103.03 (19)

## supplementary materials

---

N3—C12—H12B	109.5	C3—N1—Cu2	109.54 (19)
H12A—C12—H12B	109.5	C7—N2—C8	109.5 (3)
N3—C12—H12C	109.5	C7—N2—C6	111.2 (3)
H12A—C12—H12C	109.5	C8—N2—C6	109.3 (3)
H12B—C12—H12C	109.5	C7—N2—Cu3	109.3 (2)
O5—C13—O4	130.9 (3)	C8—N2—Cu3	112.3 (2)
O5—C13—C14	117.0 (3)	C6—N2—Cu3	105.17 (19)
O4—C13—C14	112.1 (3)	C12—N3—C11	109.9 (2)
F4—C14—F6	109.3 (4)	C12—N3—C10	109.2 (3)
F5B—C14—F4B	108 (2)	C11—N3—C10	111.5 (2)
F5B—C14—F6B	96 (2)	C12—N3—Cu4	114.56 (19)
F4B—C14—F6B	105 (2)	C11—N3—Cu4	108.6 (2)
F4—C14—F5	107.6 (4)	C10—N3—Cu4	103.00 (18)
F6—C14—F5	104.0 (3)	C1—O1—Cu1	119.59 (18)
F4—C14—C13	112.7 (3)	C1—O1—Cu2	112.44 (19)
F6—C14—C13	113.2 (3)	Cu1—O1—Cu2	105.70 (9)
F4B—C14—C13	115.1 (10)	C5—O2—Cu3	108.43 (17)
F6B—C14—C13	107.4 (12)	C5—O2—Cu1	121.87 (18)
F5—C14—C13	109.4 (3)	Cu3—O2—Cu1	102.93 (10)
O7—C15—O6	130.7 (3)	C9—O3—Cu1	117.73 (17)
O7—C15—C16	116.1 (3)	C9—O3—Cu4	112.83 (17)
O6—C15—C16	113.2 (3)	Cu1—O3—Cu4	106.00 (10)
F2—C16—F3	107.9 (3)	C13—O4—Cu2	127.6 (2)
F2—C16—F1	106.5 (3)	C13—O5—Cu3	125.9 (2)
F3—C16—F1	107.2 (3)	C15—O6—Cu3	127.6 (2)
F2—C16—C15	113.2 (3)	C15—O7—Cu4	125.6 (2)
F3—C16—C15	109.6 (3)	C17—O8—Cu4	127.6 (2)
F1—C16—C15	112.2 (3)	C17—O9—Cu2	124.8 (2)
O9—C17—O8	131.7 (3)	Cu2—O10—Cu4	110.45 (10)
O9—C17—C18	115.2 (3)	Cu2—O10—Cu3	113.15 (10)
O8—C17—C18	113.1 (3)	Cu4—O10—Cu3	108.20 (9)
F9—C18—F8	107.8 (3)	Cu2—O10—H10	108.2
F9—C18—F7	106.0 (3)	Cu4—O10—H10	108.3
F8—C18—F7	106.5 (3)	Cu3—O10—H10	108.4
F9—C18—C17	113.9 (3)	C19—O11—C22	109.0 (3)
O1—C1—C2—N1	-45.7 (3)	O10—Cu3—O2—C5	-165.02 (19)
O2—C5—C6—N2	52.7 (3)	N2—Cu3—O2—C5	20.4 (2)
O3—C9—C10—N3	-42.2 (3)	O5—Cu3—O2—C5	-80.55 (19)
O5—C13—C14—F5B	-46 (3)	O10—Cu3—O2—Cu1	64.62 (9)
O4—C13—C14—F5B	135 (3)	N2—Cu3—O2—Cu1	-109.95 (11)
O5—C13—C14—F4	114.5 (4)	O5—Cu3—O2—Cu1	149.09 (9)
O4—C13—C14—F4	-64.8 (5)	O1—Cu1—O2—C5	146.6 (2)
O5—C13—C14—F6	-10.3 (5)	O3—Cu1—O2—C5	-113.9 (2)
O4—C13—C14—F6	170.4 (3)	Cl1—Cu1—O2—C5	13.8 (2)
O5—C13—C14—F4B	-179.5 (19)	O1—Cu1—O2—Cu3	-91.71 (10)
O4—C13—C14—F4B	1.2 (19)	O3—Cu1—O2—Cu3	7.71 (10)
O5—C13—C14—F6B	63.9 (19)	Cl1—Cu1—O2—Cu3	135.49 (6)
O4—C13—C14—F6B	-115.4 (19)	C10—C9—O3—Cu1	138.4 (2)
O5—C13—C14—F5	-125.8 (4)	C10—C9—O3—Cu4	14.3 (3)

O4—C13—C14—F5	54.9 (4)	O2—Cu1—O3—C9	146.95 (19)
O7—C15—C16—F2	21.6 (4)	O1—Cu1—O3—C9	-114.43 (19)
O6—C15—C16—F2	-160.1 (3)	Cl1—Cu1—O3—C9	15.7 (2)
O7—C15—C16—F3	-98.9 (4)	O2—Cu1—O3—Cu4	-85.68 (10)
O6—C15—C16—F3	79.5 (4)	O1—Cu1—O3—Cu4	12.93 (11)
O7—C15—C16—F1	142.1 (3)	Cl1—Cu1—O3—Cu4	143.02 (6)
O6—C15—C16—F1	-39.5 (4)	O10—Cu4—O3—C9	-171.33 (19)
O9—C17—C18—F9	153.1 (3)	O8—Cu4—O3—C9	91.0 (5)
O8—C17—C18—F9	-28.5 (4)	N3—Cu4—O3—C9	10.4 (2)
O9—C17—C18—F8	-86.0 (4)	O7—Cu4—O3—C9	-84.08 (19)
O8—C17—C18—F8	92.4 (3)	O10—Cu4—O3—Cu1	58.43 (10)
O9—C17—C18—F7	32.4 (4)	O8—Cu4—O3—Cu1	-39.2 (5)
O8—C17—C18—F7	-149.2 (3)	N3—Cu4—O3—Cu1	-119.84 (11)
O11—C19—C20—C21	36.9 (4)	O7—Cu4—O3—Cu1	145.67 (9)
C19—C20—C21—C22	-33.5 (4)	O5—C13—O4—Cu2	17.2 (5)
C20—C21—C22—O11	19.0 (4)	C14—C13—O4—Cu2	-163.6 (2)
C1—C2—N1—C4	169.3 (3)	O10—Cu2—O4—C13	-39.4 (3)
C1—C2—N1—C3	-69.9 (3)	O1—Cu2—O4—C13	64.3 (4)
C1—C2—N1—Cu2	47.6 (3)	N1—Cu2—O4—C13	142.9 (3)
O4—Cu2—N1—C4	51.0 (2)	O9—Cu2—O4—C13	-125.7 (3)
O1—Cu2—N1—C4	-148.2 (2)	O4—C13—O5—Cu3	-15.7 (5)
O9—Cu2—N1—C4	-51.8 (2)	C14—C13—O5—Cu3	165.1 (2)
O4—Cu2—N1—C2	170.02 (17)	O2—Cu3—O5—C13	-51.2 (3)
O1—Cu2—N1—C2	-29.17 (17)	O10—Cu3—O5—C13	37.4 (3)
O9—Cu2—N1—C2	67.24 (17)	O6—Cu3—O5—C13	129.2 (3)
O4—Cu2—N1—C3	-71.1 (2)	N2—Cu3—O5—C13	-136.9 (3)
O1—Cu2—N1—C3	89.7 (2)	O7—C15—O6—Cu3	8.0 (5)
O9—Cu2—N1—C3	-173.9 (2)	C16—C15—O6—Cu3	-170.1 (2)
C5—C6—N2—C7	84.5 (3)	O10—Cu3—O6—C15	-41.7 (3)
C5—C6—N2—C8	-154.5 (3)	N2—Cu3—O6—C15	132.2 (3)
C5—C6—N2—Cu3	-33.7 (3)	O5—Cu3—O6—C15	-126.6 (3)
O2—Cu3—N2—C7	-111.5 (2)	O6—C15—O7—Cu4	-1.2 (5)
O6—Cu3—N2—C7	76.2 (2)	C16—C15—O7—Cu4	176.9 (2)
O5—Cu3—N2—C7	-25.8 (2)	O10—Cu4—O7—C15	30.6 (3)
O2—Cu3—N2—C8	126.7 (2)	O8—Cu4—O7—C15	122.6 (3)
O6—Cu3—N2—C8	-45.5 (2)	O3—Cu4—O7—C15	-58.4 (3)
O5—Cu3—N2—C8	-147.5 (2)	N3—Cu4—O7—C15	-143.6 (3)
O2—Cu3—N2—C6	7.9 (2)	O9—C17—O8—Cu4	-12.4 (5)
O6—Cu3—N2—C6	-164.4 (2)	C18—C17—O8—Cu4	169.5 (2)
O5—Cu3—N2—C6	93.6 (2)	O10—Cu4—O8—C17	-30.2 (3)
C9—C10—N3—C12	170.1 (3)	O3—Cu4—O8—C17	67.2 (6)
C9—C10—N3—C11	-68.2 (3)	N3—Cu4—O8—C17	146.9 (3)
C9—C10—N3—Cu4	48.0 (3)	O7—Cu4—O8—C17	-117.8 (3)
O8—Cu4—N3—C12	41.2 (2)	O8—C17—O9—Cu2	15.4 (5)
O3—Cu4—N3—C12	-150.3 (2)	C18—C17—O9—Cu2	-166.5 (2)
O7—Cu4—N3—C12	-57.7 (2)	O10—Cu2—O9—C17	24.9 (3)
O8—Cu4—N3—C11	-82.1 (2)	O4—Cu2—O9—C17	118.8 (3)
O3—Cu4—N3—C11	86.44 (19)	O1—Cu2—O9—C17	-64.6 (3)
O7—Cu4—N3—C11	179.06 (19)	N1—Cu2—O9—C17	-149.8 (3)

## supplementary materials

---

O8—Cu4—N3—C10	159.62 (18)	O4—Cu2—O10—Cu4	-168.32 (10)
O3—Cu4—N3—C10	-31.84 (18)	O1—Cu2—O10—Cu4	30.61 (10)
O7—Cu4—N3—C10	60.78 (18)	O9—Cu2—O10—Cu4	-65.89 (10)
C2—C1—O1—Cu1	144.4 (2)	O4—Cu2—O10—Cu3	70.20 (11)
C2—C1—O1—Cu2	19.6 (3)	O1—Cu2—O10—Cu3	-90.86 (11)
O2—Cu1—O1—C1	-115.4 (2)	O9—Cu2—O10—Cu3	172.64 (11)
O3—Cu1—O1—C1	144.7 (2)	O8—Cu4—O10—Cu2	72.73 (11)
Cl1—Cu1—O1—C1	17.8 (2)	O3—Cu4—O10—Cu2	-95.76 (10)
O2—Cu1—O1—Cu2	12.61 (11)	O7—Cu4—O10—Cu2	171.17 (10)
O3—Cu1—O1—Cu2	-87.35 (10)	O8—Cu4—O10—Cu3	-162.91 (10)
Cl1—Cu1—O1—Cu2	145.78 (6)	O3—Cu4—O10—Cu3	28.60 (10)
O10—Cu2—O1—C1	-170.14 (19)	O7—Cu4—O10—Cu3	-64.47 (10)
O4—Cu2—O1—C1	85.4 (3)	O2—Cu3—O10—Cu2	23.58 (11)
N1—Cu2—O1—C1	5.7 (2)	O6—Cu3—O10—Cu2	-164.31 (11)
O9—Cu2—O1—C1	-84.9 (2)	O5—Cu3—O10—Cu2	-63.05 (10)
O10—Cu2—O1—Cu1	57.71 (10)	O2—Cu3—O10—Cu4	-99.15 (10)
O4—Cu2—O1—Cu1	-46.8 (3)	O6—Cu3—O10—Cu4	72.97 (11)
N1—Cu2—O1—Cu1	-126.46 (11)	O5—Cu3—O10—Cu4	174.22 (10)
O9—Cu2—O1—Cu1	143.00 (10)	C20—C19—O11—C22	-25.9 (4)
C6—C5—O2—Cu3	-44.1 (3)	C21—C22—O11—C19	4.2 (4)
C6—C5—O2—Cu1	74.9 (3)		

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
O10—H10 $\cdots$ O11	1.00	1.73	2.723 (3)	174

Fig. 1






Fig. 2

